

admission. We developed the Italian AES through translation back-translation and administered it to acutely hospitalized psychiatric patients.

Objectives/Aims To verify psychometric characteristics of the Italian AES. To Examine the AES factor structure.

Methods $n = 156$ acutely hospitalized patients (48% women, 69% voluntary) were recruited in two university hospitals in Rome (Umberto I Policlinic, Sant'Andrea Hospital) and were administered the Italian AES. We conducted a principal component analysis (PCA) with equamax rotation.

Results Socio-demographic and clinical characteristics of the sample are reported in Table 1. The Italian AES had good internal consistency (Cronbach's $\alpha = 0.90$); Guttman split-half reliability coefficient was 0.90. AES total score significantly differed between voluntary and involuntary patients (5.08 ± 4.1 vs. 8.1 ± 4.9 , $P < 0.05$). PCA disclosed a three-factor solution explaining 59.3 of the variance. Significant correlations emerged between AES total score and clinical variables (Table 2). Pearson's correlation coefficient disclosed a significant correlation between perceived coercion and psychiatric symptoms severity (BPRS total score).

Conclusions The Italian version of AES and proposed new factor structure proved reliable.

Table 1

Age, years, M (SD)	40.5 (12.7)
Women	48%
Education, years, M (SD)	12.1 (4.0)
Disease duration, years, M (SD)	12.1 (9.4)
Diagnosis	
Schizophrenia spectrum disorders	35.9%
Bipolar disorders	35.3%
Depressive disorders	19.2%
Others	9.6%
Number of previous psychiatric hospitalizations, M (SD)	1.4 (2.9)
Number of previous involuntarily psychiatric hospitalization, M (SD)	0.7 (2.6)
MMSE total score, M (SD)	25.6 (2.7)
BPRS total score, M (SD)	54.2 (13.0)

Table 2

Admission Experience Survey items	Perceived Coercion (Cronbach's $\alpha = 0.84$)	External pressure (Cronbach's $\alpha = 0.79$)	Choice expression (Cronbach's $\alpha = 0.71$)
7. It was my idea to come into the hospital	0.78		
15. I had more influence than anyone else on whether I came into the hospital	0.73		
4. I chose to come into the hospital	0.69		
1. I felt free to do what I wanted about coming into the hospital	0.63		
13. My opinion about coming into the hospital didn't matter	0.60		
14. I had lot of control over whether I went into the hospital	0.59		
2. People tried to force me to come into the hospital	0.46	0.51	
11. they said they would make me come into the hospital	0.50		
10. I was threatened with commitment		0.86	
6. Someone threatened me to get me to come into the hospital		0.79	
8. Someone physically tried to make me come into the hospital		0.63	
12. No one tried to force me to come into the hospital	0.46	0.53	
5. I got to say what I wanted about coming into the hospital			0.75
3. I had enough of a chance to say whether I wanted to come into the hospital			0.67
9. No one seemed to want to know whether I wanted to come into the hospital			0.60

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.304>

EW0691

Sexual dysfunctions and treatment compliance in individuals with psychotic disorder

G. Unsal^{1,*}, S. Karaca¹, N. Onan², Y. Can Oz¹, S. Aydın³, H. Aydoğan⁴, H. Gulec⁵, E. Ongun¹, N. Eren⁶

¹ Marmara university, faculty of health sciences, psychiatric nursing department, Istanbul, Turkey

² Karabuk university school of health science, psychiatric nursing department, Karabuk, Turkey

³ Erenköy psychiatric and neurological diseases training and research hospital, training psychiatric nursing, Istanbul, Turkey

⁴ Erenköy psychiatric and neurological diseases training and research hospital, management nursing, Istanbul, Turkey

⁵ Erenköy psychiatric and neurological diseases training and research hospital, community mental health, Istanbul, Turkey

⁶ Istanbul university Istanbul, faculty of medicine, psychiatry, Istanbul, Turkey

* Corresponding author.

Introduction Sexual dysfunctions are more common in individuals with psychotic disorders and has a major impact on both quality of life and compliance.

Objectives The purpose of this study is to investigate whether a relationship between sexual dysfunction and level of treatment compliance in individuals with psychotic disorders.

Methods The sample group of the study consisted of 173 inpatients who agreed to participate were selected by random sampling method. The permission was obtained from the hospital's ethics committee. In this study, to assess the sexual functionality Golombok-Rust inventory of sexual satisfaction male and female form and to assess the treatment compliance; medical treatment compliance rate scale is used.

Results When sexual problems and treatment compliance compared to gender, subscales of satisfaction ($t = 4.423$, $P = 0.000$), avoidance ($t = 3.348$, $P = 0.001$), touch ($t = 2.165$, $P = 0.032$) and overall total ($t = 4.015$, $P = 0.000$), although a statistically significant difference was found, there were no differences in treatment compliance. Additionally, there is no relation between sexual problems and treatment compliance in men. It is also found that there is a weak negative statistical relation amongst treatment compliance and communication ($r = -0.244$, $P = 0.027$), avoidance ($r = -0.270$, $P = 0.014$), anorgasmia ($r = -0.253$, $P = 0.022$) and overall total ($r = -0.249$, $P = 0.024$) in women. According to these findings while sexual problems increase, treatment compliance decreases.

Conclusions The level of compliance to the treatment and subscales of sexual problems; satisfaction, avoidance, and touch mean scores differ from each other. There was a weak negative correlation in between Women's compliance and sexual problems.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.305>

EW0692

Post-traumatic stress disorder screening among Syrian war victims

A. Kutluoğlu¹, B.Ö. Ünsalver^{2,*}, A. Evrensel¹

¹ Uskudar university, psychology, İstanbul, Turkey

² Uskudar university, medical documentation and secretariat, İstanbul, Turkey

* Corresponding author.

Introduction War in Syria and related refugee crisis has caught worldwide attention for the past few years. The war is still continuing by the time of the writing of this abstract. War is one of the reasons of posttraumatic stress disorder (PTSD). Syrian people experience many traumatic events like witnessing death, torture and rape.

Aims The aim of this research is to screen for PTSD among Syrian war victims.

Methods A total of 150 (women = 52, men = 98) war victims between 18–65 years, inhabiting a Syrian camp near the Turkish border were screened with trauma response checklist for PTSD between March 11 and April 11 2015.

Results Significant amount of the interviewed participants were found to show symptoms of PTSD. Seventy-six percent ($n = 114$) of the participants had experienced a traumatic event. In total, 80.6% ($n = 121$) were experiencing distress. Seventy-eight percent ($n = 117$) had avoidance. Eighty-four percent ($n = 126$) had negative

interfering thoughts. Eighty-eight percent ($n = 132$) were experiencing trauma associated disturbance. Answers from 78.6% ($n = 118$) of the participants indicated that they should be further referred to a specialist. The findings were not affected by gender or age.

Conclusion We have found that most of the participants showed signs of PTSD. Our findings highlight the psychological impact of war on Syrian people. A definite diagnosis of PTSD can be made with detailed psychiatric examination, however given the amount of victims and available staff a brief screening instrument may help identify potential cases to be further evaluated. PTSD has life-long consequences and trauma can be passed through generations. International support for war victims should include psychological support and interventions.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.306>

EW0693

Program for the use of antipsychotics with metabolic monitoring in North Carolina medicaid children

S. Wegner^{1,2,*}, J. McKee³, T. Trygstad⁴, L. Wegner⁵, A. Stiles⁶

¹ AccessCare, Morrisville, USA

² University of North Carolina-Chapel Hill, department of pediatrics, Chapel Hill, USA

³ Community care of North Carolina, pharmacy operations and payer programs, Raleigh, USA

⁴ Community care of North Carolina, pharmacy programs, Raleigh, USA

⁵ University of North Carolina-Chapel Hill, UNC development and behavioral pediatrics, Chapel Hill, USA

⁶ University of North Carolina-Chapel Hill, Sr. VP - network development and strategic affiliations, Chapel Hill, USA

* Corresponding author.

Introduction Children are at greater risk than adults for weight gain and metabolic disorders including hyperlipidemia and diabetes with newer antipsychotics. A web-based safety-monitoring program using a prior documentation model required submission of patient safety data (prior documentation) for insurance coverage at the pharmacy point of sale. This program launched in April of 2011, covering all NC Medicaid and Health Choice recipients under age 18. Clinical monitoring parameters and interactive educational features were developed with pediatric psychiatric experts and key mental health stakeholder groups.

Objectives Using a four-year run in period and a full 9 months of post implementation claims data, evaluate the rates of antipsychotic prescribing and safety monitoring before and after the implementation of the A + KIDS program.

Results Implementation of this program was associated with a consistent monthly decrease in overall antipsychotic use and increases in patient monitoring of glucose and lipid (Figure. 1, Table 1).

Conclusions The prior documentation registry was effective in decreasing antipsychotic use and increasing safety monitoring. The impact of changing to more traditional prior authorization on the same clinical endpoints is currently under evaluation.




Fig. 1 A+KIDS Population Antipsychotic Prescription Fills per 1000 Medicaid Enrollees Per Day

Table 1 Percent of A+KIDS Patients on an Antipsychotic with Metabolic Monitoring Recorded in Claims.

Year Ending	Glucose Screening Percent	Cholesterol Screening Percent
June 2010	52%	27%
June 2011	55%	32%
June 2012	60%	41%

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.307>

e-Poster Walk: Neuroimaging and neuroscience in psychiatry

EW0694

Effort-based reward task, a behavioral measure to study negative symptoms in schizophrenia

A. Arcos¹, D. Berge^{1,2,*}, C. Pretus¹, A. Pous¹, C. Diez-Aja³, L. Gomez³, O. Vilarroya^{1,2}

¹ Universitat autonoma de Barcelona, psychiatry, Barcelona, Spain

² IMIM, hospital del MAR research institute, neuroscience, psychiatry, Barcelona, Spain

³ Parc de Salut Mar, psychiatry, INAD, Barcelona, Spain

* Corresponding author.

Negative symptoms in schizophrenia, and specifically amotivation/apathy, have been correlated with impaired general functioning. Its neurobiological basis are thought to rely on an aberrant reward system. To study the association of reward deficits and negative symptoms, 25 schizophrenia patients and 35 controls underwent a new reward behavioral task. Briefly, patients had to choose a level of effort (1 to 3), each one corresponding to a progressively increasing number of required button presses and 3 different probabilities to win an economic reward. We compared the chosen effort between groups and correlated this output with the score of the Brief negative symptoms scale in the group of patients. Patients chose less effort than controls but without reaching significance level (mean patients effort: 2.49 vs controls: 2.76, $P = 0.064$). A negative correlation was found between BNSS score and effort chosen for the maximum reward corrected by sex ($t: -0.021, P = 0.045$). When the group of patients was split according to negative symptoms score, patients with more negative symptoms (BNSS score > 23) chose significantly less effort than patients