

# Using citizen science to study a mesocarnivore: the jungle cat *Felis chaus* in Sri Lanka

SRIYANIE MITHTHAPALA, JEREMY DERTIEN, NIROSHA LIYANAGE  
NIROSHAN MIRANDO, ANYA AVANTHI WEERAWARDANA RATNAYAKA  
ASHAN THUDUGALA, DARSHANI WIJESINGHE and SAMPATH DE ALWIS GOONATILAKE

**Abstract** We used citizen science and inexpensive methodology to assess the distribution of the jungle cat *Felis chaus*, a relatively common species in Sri Lanka but the least studied of the four wild cat species occurring in the country. We obtained three types of records of the jungle cat: georeferenced photographs of the species from the public; sightings obtained from print and social media, and provided via an online sighting form; and sightings by field biologists. We combined the 112 unique records obtained in this way with the 21 records from the 2012 National Red List distribution map of the species, and used *MaxEnt* to predict habitat suitability for the species. The new sightings were primarily in drier regions, expanding the known extent of occurrence for this species in Sri Lanka. Of the new sightings, 7.1% were road kills. Distance to nearest riverine forest, annual precipitation and distance to the nearest reservoir were the most important variables explaining habitat suitability. These findings validate our hypotheses that the species is more widespread than demonstrated previously and also ranges in human-dominated landscapes outside protected areas. Our study provides a model for how ecological and behavioural information for common species can be obtained inexpensively and incorporated into species distribution models. Studies of species such as the jungle cat, which are neither threatened nor charismatic, will help ensure that we keep common species common.

**Keywords** Citizen science, *Felis chaus*, jungle cat, habitat suitability modelling, common, inexpensive, Sri Lanka

Supplementary material for this article is available at [doi.org/10.1017/S0030605321000764](https://doi.org/10.1017/S0030605321000764)

## Introduction

Although the importance of ecosystems is now central to discussions about biodiversity (Millennium Ecosystem Assessment, 2005; United Nations, 2015; Schultz et al., 2016), the distribution and habitat use of the many species that are essential components of these ecosystems remain poorly known (Meyer et al., 2015; Minin et al., 2016). Within this context, an estimated 1 million species of animals and plants are threatened with extinction (Díaz et al., 2019). Mammalian carnivores have lost > 75% of their original range (Faurby & Svenning, 2015; Díaz et al., 2019): e.g. the red wolf *Canis rufus* (> 99%), tiger *Panthera tigris* (95%), lion *Panthera leo* (94%) and leopard *Panthera pardus* (79.4%; Wolf & Ripple, 2017). Approximately 25% of known mammalian species are believed to be at risk of extinction (Minin et al., 2016).

Because of the urgent need for conservation action for threatened species, research has predominantly been based on proxies such as charismatic, endemic or threatened species (Ducarme et al., 2013; Brum et al., 2017), with large-bodied, charismatic species studied preferentially (Brooke et al., 2014; Albert et al., 2018). For example, of 4,351 peer-reviewed articles on canids and felids published during 2013–2017 there were 359 publications on the Endangered tiger and only one on the small, Near Threatened rusty-spotted cat *Prionailurus rubiginosus* (Tenson, 2018). Small wild cats receive only 1.2% of all funding for research on wild cats (Global Wildlife Conservation, 2020).

This focus on large, charismatic species ignores many medium-sized, common species. There are many common, little studied mesocarnivores that are important as predators of species such as rodents, as competitors with other medium-sized predators, and as omnivorous seed dispersers, and ‘may be fundamentally important drivers of ecosystem function, structure, or dynamics’ (Gittleman & Gompper, 2005; Roemer et al., 2009, p. 171). In response to this concern there have been appeals to ensure common species remain common (Ramesh & McGowan, 2009;

SRIYANIE MITHTHAPALA (Corresponding author, [orcid.org/0000-0003-1585-5581](https://orcid.org/0000-0003-1585-5581), [sriyanie.miththapala@gmail.com](mailto:sriyanie.miththapala@gmail.com)) Independent researcher, 165/12-5/1 Park Road, Colombo 00500, Sri Lanka

JEREMY DERTIEN ([orcid.org/0000-0003-4011-0258](https://orcid.org/0000-0003-4011-0258)) Department of Forestry and Environmental Conservation, Clemson University, Clemson, USA

NIROSHA LIYANAGE ([orcid.org/0000-0003-4332-4235](https://orcid.org/0000-0003-4332-4235)) Independent researcher, Pannipitiya, Sri Lanka

NIROSHAN MIRANDO ([orcid.org/0000-0002-6116-0839](https://orcid.org/0000-0002-6116-0839)) Independent researcher, Winnipeg, Canada

ANYA AVANTHI WEERAWARDANA RATNAYAKA ([orcid.org/0000-0001-8818-8471](https://orcid.org/0000-0001-8818-8471)) and ASHAN THUDUGALA ([orcid.org/0000-0003-2853-3098](https://orcid.org/0000-0003-2853-3098)) Small Cat Advocacy and Research, Bowalawatta, Kandy, Sri Lanka

DARSHANI WIJESINGHE ([orcid.org/0000-0003-2651-984X](https://orcid.org/0000-0003-2651-984X)) Independent researcher, Athurugiriya, Sri Lanka

SAMPATH DE ALWIS GOONATILAKE ([orcid.org/0000-0001-7859-3844](https://orcid.org/0000-0001-7859-3844)) Independent researcher, Kawdana, Dehiwala, Sri Lanka

Received 26 August 2020. Revision requested 3 February 2021.

Accepted 24 May 2021. First published online 30 May 2022.

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

*Oryx*, 2022, 56(5), 782–788 © The Author(s), 2022. Published by Cambridge University Press on behalf of Fauna & Flora International doi:10.1017/S0030605321000764

Redford et al., 2013; Frimpong, 2018; USGS, 2018; Baker et al., 2019).

Here we describe how we used citizen science, with minimal financial investment, to document the occurrence of the jungle cat *Felis chaus*, a relatively common species in Sri Lanka but the least studied of the four wild cat species occurring in the country (Kittle & Watson, 2018; Miththapala, 2018). The other three species are the fishing cat *Prionailurus viverrinus*, rusty-spotted cat and leopard *Panthera pardus kotiya*. The jungle cat is categorized as Least Concern on the IUCN Red List, although more information is needed on the general ecology of the species and its status across its range, and the population trend is decreasing (Gray et al., 2021). In Sri Lanka the jungle cat is categorized as Near Threatened (Ministry of Environment Sri Lanka, 2012), but little is known about its habitat use and distribution (Miththapala, 2018). We aimed to model habitat suitability for this species in Sri Lanka and improve knowledge of its distribution. Based on our observations that the species may have a greater tolerance for, and perhaps benefit from, human activities than was previously believed, we hypothesized that the jungle cat is more widespread than described in the National Red List assessment.

## Study area

Sri Lanka is a highly biodiverse tropical island nation in the Indian Ocean (Ministry of Mahaweli Development & Environment, 2016). The topography of the land mass of 65,610 km<sup>2</sup> is a result of geological uplift and erosion that have led to the formation of three peneplains, with the highest (1,501–2,524 m), forming a central hill massif. The combination of these hills and the winds of the south-west and north-east monsoons result in three main climatic zones: the wet zone in the south-west (low and mid elevation and montane) with a mean annual rainfall of 2,500–5,000 mm; the dry zone extending over much of the lowlands, with a mean annual rainfall of 1,250–1,900 mm; and the intermediate zone, a narrow strip in the lowlands and on the eastern slopes of the central hills, between the wet and dry zones with a mean annual rainfall of 1,900–2,500 mm (Ministry of Mahaweli Development & Environment, 2016; Fig. 1a). Land cover is a mosaic of forests, agriculture, plantations (tea, rubber and coconut), homesteads, roads and scattered towns. Nearly 10,000 reservoirs, some up to 820 years old, provide perennial and seasonal sources of water to the large expanse of the lowland dry zone (Ministry of Mahaweli Development & Environment, 2016).

## Methods

We conducted this study during 8 November 2016–18 March 2019, using three methods to obtain records of jungle cats.

Firstly, we took advantage of increased interest in wildlife and wildlife photography, and the popularization of sharing wildlife-related photographs on social media. We created a website for this project, and through an e-mail campaign and advertisements and posts on Facebook (Meta Platforms, Menlo Park, USA), we invited wildlife and photography enthusiasts to submit geo-referenced photographs of the jungle cat and, if possible, to identify the habitat from among a given list of habitats (primary collection in Fig. 1; Miththapala, 2016). Each submission was verified as a jungle cat by SM or SdAG, and uploaded onto an online map (Miththapala, 2016). Secondly, we collected data on sightings from newspapers and social media pages (secondary collection in Fig. 1). We searched Instagram (Meta Platforms, Menlo Park, USA) for '#junglecat', recording any sightings in Sri Lanka. We also used personal contacts made during fieldwork and through awareness programmes to request information (by phone or text) of any jungle cat sightings. We asked these contacts to use an online small wild cat sighting form, and verified records from submitted photographs. To enlist a wide audience, we made announcements on the organizational pages of Small Cat Advocacy & Research (2017), Urban Fishing Cat Conservation Project (2020), the Save Fishing Cats Conservation Project (2015) and other pages with an interest in wildlife news. Thirdly, we communicated with other field biologists, seeking additional sightings, and also included SdAG's own records of the jungle cat.

Using *Arcmap 10.3.1* (Esri, Redlands, USA), we combined all new records for the jungle cat with those from the National Red List assessment for this species (Ministry of Environment Sri Lanka, 2012), obtained from the Biodiversity Secretariat of the Ministry of Mahaweli Development and Environment (Fig. 1a). We used these records to model habitat suitability for the species, using *MaxEnt* (Phillips et al., 2006). *MaxEnt* is a machine learning approach to modelling species habitat suitability using presence-only data and environmental covariates. *MaxEnt* models the probability density of environmental features at the locations of species records and at a large set of background points across the study area, and then attempts to find the broadest probability distribution to describe habitat suitability within the constraints of the environmental features (Elith et al., 2011).

We generated 13 environmental covariate layers to model habitat suitability (Table 1). We used a raster land-use layer of agricultural forest plantations, home gardens, grasslands, open forests/scrub forests, dense forests, marshes and paddy fields (Surveyor General, Survey Department, 2016), and a water features layer for reservoirs and streams/rivers (Department of Agrarian Development, 2011). To identify riverine forests, we trained a colleague to identify and digitize riverine forest habitat in Sri Lanka, using *Google Earth* (Google, Mountain View, USA), and we then visually checked the digitized areas (we were not able to ground-




FIG. 1 (a) Water bodies in the study area, and records of the jungle cat *Felis chaus* from the Ministry of Environment Sri Lanka (2012), primary and secondary collections (see text for details), SdAG's sightings, other biologists and road kill. (b) Our model of predicted habitat suitability for the jungle cat in Sri Lanka. Protected areas from Ministry of Mahaweli Development and Environment (2016), and water bodies from Department of Agrarian Development (2011).

truth the data because of the lack of financial resources to do so). We calculated the Euclidean distance from each jungle cat record to the nearest riverine forest, marsh, paddy field, stream/river, and reservoir.

We transformed the land-use raster layer into a point layer and used a kernel density estimator to calculate the densities of plantation forests, home gardens, grasslands, open forests (which included arid-mixed evergreen forests, commonly called scrublands) and dense forests. We used a 45 km<sup>2</sup> moving window with a radius (3.78 km) based on the approximate distance moved by individual jungle cats in a night (Sunquist & Sunquist, 2002), the only information available on the species' home range. We followed a similar procedure to calculate road density using the Survey Department's database of Sri Lanka's roads (Surveyor General, Survey Department, 2016) but used linear rather than point density. All densities were transformed into km<sup>2</sup>, from which we calculated the per cent surface area of each land cover surrounding a point within a 45 km<sup>2</sup> moving window. We used elevation data from the ASTER Global Digital Elevation Model (DEM) Version 2 to create a 30-m resolution raster layer of Sri Lanka. We obtained annual rainfall and temperature data for Sri Lanka from *WorldClim 2.0* (Hijmans et al., 2005). Spatial correlations were assessed to reduce the multicollinearity of

environmental covariates; one covariate was removed from a covariate pair if  $|r| \leq 0.80$ . To reduce potential bias in the presence-only data as a result of unequal sampling effort, we used a spatial rarefaction tool (*SDMToolbox 2.4*; Brown et al., 2017) to truncate one sighting from any pair of points that were within 1 km of each other. We used *ArcGIS 2.2* (Esri, Redlands, USA) and *R 3.6* (R Core Team, 2019) for processing spatial data and preparation for model input.

We ran *MaxEnt* models for a maximum of 1,000 iterations, including linear, quadratic, product and hinge features, and used default regularization settings (Elith et al., 2011). We used a 10-fold cross-validation method to estimate model error and predictive performance (Rodriguez et al., 2010; Pascoe et al., 2019). This procedure runs 10 iterations of the same set of model variable inputs, but with different training and testing subsets per iteration. We created and ran models with different sets of variable inputs. We assessed model performance and selected a model structure by using the highest mean area under the receiver operating characteristic curve (AUC) and if all variables had a positive influence on the regularized training gain of the model; AUC values range from 0 to 1, with 0.5 indicating that the model is no better than random at predicting habitat suitability and 1 indicating a perfect ability of the model to estimate species presence (Phillips et al., 2006). Regularized

TABLE 1 Per cent contribution, in decreasing order, of 13 environmental covariates to the *MaxEnt* model of habitat suitability for the jungle cat *Felis chaus* in Sri Lanka. Density refers to the per cent coverage of a land cover within a 3.78 km radius (45 km<sup>2</sup>) of a particular location.

Covariate	Contribution (%)
Distance from nearest riverine forest (km)	24.62
Distance from nearest reservoir (km)	16.77
Annual rainfall (mm)	13.84
Density of agricultural forests (%)	13.81
Distance from nearest stream/river (km)	7.50
Elevation (m)	5.17
Density of roads (%)	4.66
Distance from nearest paddy field (km)	3.63
Mean annual temperature (°C)	3.32
Density of open forests (%)	3.10
Density of home gardens (%)	2.02
Distance from nearest marsh (km)	1.03
Density of dense forests (%)	0.54

training gain is a measure of the increased model fit of the *MaxEnt* model distribution to the species locations versus the fit of a uniform distribution to the species locations, with a penalty factor. Higher values of regularized training gain for a model with one specific variable is an indication that variable is of greater importance in modelling habitat suitability (Kissel & Poserina, 2017).

## Results

We collated 112 jungle cat records spanning 2016–2019, a substantial increase compared to the 21 sightings used for the species assessment on the National Red List (Ministry of Environment Sri Lanka, 2012; Fig. 1a, Plate 1). We verified records from photographs received from the public and from photographs in newspapers and on social media, and accepted some records without supporting photographic evidence if the submitter's field identification knowledge and experience was known to us. These records expand the known range for this species in Sri Lanka and also show that it occurs outside protected areas (Fig. 1a). The jungle cat appears to be a habitat generalist, with records in a range of open habitats, including human-dominated areas such as paddy fields and near human-made water reservoirs, and in forests (S. Miththapala et al., unpubl. data, 2022). Eight of the 112 records (7.1%) were road kills.

The best *MaxEnt* model had a mean AUC of 0.873, indicating the model had a relatively strong diagnostic ability to predict habitat suitability (Fig. 1b, Supplementary Fig. 1). The model included 13 environmental covariates, with six of these having > 80% cumulative contribution to the model (Table 1, Supplementary Fig. 2). Distances to nearest riverine forest and reservoir contributed the most to the

model and were both negatively associated with habitat suitability for jungle cats (i.e. increasing distances from these habitats resulted in decreased habitat suitability). The highest probability of presence was within 5 km of riverine forests and in proximity to a reservoir (Supplementary Fig. 2). Annual rainfall and the density of agricultural forests (rubber, palmyra, coconut, cashew, mixed-tree and other perennial crops) were also high contributors to the model (Table 1, Supplementary Fig. 3). The highest probability of presence was in areas that received 500–1,000 mm of rainfall per year and had the lowest densities of plantation forests.

Relative to other covariates, the densities of roads, open forest and home gardens contributed little to the final model. Road density showed a strong negative relationship with habitat suitability when road density was the only variable considered in the model but this became a positive relationship when all other variables in the model were included. Habitat suitability was negatively correlated with the density of home gardens. When the density of home gardens was > 33%, habitat suitability was < 0.50.

## Discussion

Our findings validate our hypotheses that the jungle cat is more widespread than shown in the current Sri Lanka National Red List and than reported by Kittle & Watson (2018), and that the species appears to have a tolerance for human activities as it was sighted in human-dominated landscapes. Figure 1a shows the jungle cat's preference for the drier parts of Sri Lanka, with most sightings in the dry zone. The habitat suitability model also indicated the species' preference for the dry zone (Fig. 1b, Table 1, Supplementary Fig. 2).

The jungle cat has also been referred to as the swamp cat or reed cat because of its occurrence in swamps and among reeds (Sunquist & Sunquist, 2002; Priszahnyuk & Belousova, 2007; Sanei et al., 2016), and it has also been recorded near rivers or wetlands (Hatt, 1959; Harrison, 1968; Mendelssohn, 1989; Dayan et al., 1990; Heptner & Sludskii, 1992; Sunquist & Sunquist, 2002; Priszahnyuk & Belousova, 2007; Avgan, 2009; Gerngross, 2014; Sanei et al., 2016). We only recorded one sighting in a riverine habitat, but the habitat suitability modelling indicated a high probability of presence within 5 km of riverine forests and in proximity to a reservoir (Table 1, Fig. 1b, Supplementary Fig. 2). However, the habitat suitability model prediction must be interpreted with caution, as we were not able to ground-truth the digitized satellite data. The jungle cat also appears to use a range of open habitats, including human-dominated areas in Sri Lanka (S. Miththapala et al., unpubl. data, 2022), as shown in other parts of its range (Patel, 2011; Kalle et al., 2013; Sanei et al., 2016; Mishra et al., 2020; Shrestha et al., 2020).



PLATE 1 A sample photograph, provided by a member of the public, taken within Udawalawe National Park, away from any human habitation (Photo: Chitral Jayatilake, 2016). We identified this individual as a jungle cat by its long legs, ear tufts, tawny head and part of the legs, most of the body grey, black and white speckled, few stripes on the legs and tail, and plain coat. The practice of hybridizing domestic with wild cats is not known in Sri Lanka (Phillips, 1935; Yapa, 2013) and therefore we are confident this is not a hybrid. Furthermore, this photograph was taken deep within the National Park, where it is unlikely that natural hybridization would occur.

Our finding that the jungle cat occurs both within and outside protected areas parallels studies in Sri Lanka of other carnivores such as the leopard (Kittle et al., 2018), fishing cat (Urban Fishing Cat Conservation Project, 2020) and elephant *Elephas maximus* (Fernando et al., 2021). However, the presence of these species in human-dominated landscapes can result in negative interactions with people. In 2019, for example, 405 elephants (Prakash et al., 2020), 12 leopards (A. Kittle, pers. comm., 2021) and 22 fishing cats (Small Cat Advocacy & Research, 2020) were killed as a result of human–wildlife interactions. Research is required to investigate whether the jungle cat is also affected in this way.

Eight of our records were road kills. Of these, six were along the south-east coastline, probably because there are good roads and little traffic in this area, facilitating fast driving. Road kills, and irresponsible driving within national parks, are threats to wildlife in Sri Lanka (Karunaratna et al., 2017) and elsewhere (Filius et al., 2020; Schwartz et al., 2020; Ascensão et al., 2021). An additional threat to the jungle cat is the decline of its prey as a result of deforestation and consequent killing by farmers in retaliation for preying on livestock (de Alwis, 1973). Habitat destruction

has not abated since 1973 and remains the prime driver of ecosystem change in Sri Lanka (Ministry of Mahaweli Development & Environment, 2016). An additional threat is the killing of jungle cats by feral dogs (Bambaradeniya & Amerasinghe, 2001).

Globally, funding for research on biodiversity is inadequate. In Sri Lanka most of the limited funds available are spent on exploration for new species (e.g. fish: Sudasinghe, 2018; amphibians: Senevirathne et al., 2018; reptiles: Mendis Wickramasinghe et al., 2020; mammals: Meegaskumbura, 2007) or studies of threatened species (e.g. leopards: Kittle et al., 2018; elephants: Fernando et al., 2021). Studying common species is not a priority, even though by studying such species and planning for their needs before they become rare, future conservation funds can be conserved (Scott et al., 1993; Redford et al., 2013). Our study provides a model for how ecological and behavioural information for common species could be obtained inexpensively and incorporated into species distribution models. From our findings we will be able to develop more precise hypotheses about habitat dependencies, and employ camera traps and GPS collars on individuals in specific locations (S. Miththapala et al., unpubl. data, 2022) to investigate the species' ecology and behaviour. Studies of species such as the jungle cat, which are neither threatened nor charismatic, will help ensure that we are 'keep [...] common species common' (Ramesh & McGowan, 2009, p. 108; Frimpong, 2018; USGS, 2018; Baker et al., 2019).

**Acknowledgements** We thank the Mohamed bin Zayed Species Conservation Fund (grant number 160513505) for financial support; the Biodiversity Secretariat for providing us with the 2012 Red List map for *Felis chaus*; Isuru Dayananda, Saranga Dissanayake Suresh Ellawala, Wasiri Rasu Gajaman, Anton Gihan, Shelley Glass, Medhisha Pasan Gunawardena, Chaminda Jayasekera, Chitral Jayatileke, Eric Joseph, Namal Kamalgoda, Sunitha Kanthi, Moditha Hiranya Kodikara Arachchi, Dineth Mallikarachchy, Srilal Miththapala, Luxshmanan Nadaraja, Abhaya Indika Nettigama, Madduma Wellalage Tiron Premajith, Stuart Peoples, Madura Priyankara, Phil Rekret, Oliver Rook, Danushka Senadeera, Pasan Senevirathne, Aisha Uduman, Milinda Wattedegedera, Mithra Weerakone, Rajiv Welikala Romesh Wijesiri and Christopher Young for data collection; Chandima Fernando, Manori Gunawardena, Andrew Kittle, Enoka Kudawidanage and Sampath Seneviratne for sharing their sighting data; Virginia Hayssen and Angie Appel for sharing references; Rob Baldwin, Janaki Galappatti and John Seidensticker for reviews; and Rob Baldwin for providing technical support for habitat suitability modelling.

**Author contributions** Funding acquisition: SM, NM; study design, supervision: SM; data collection: SM, AAWR, AT, SdAG; data analysis: NL, JD, DW, SdAG; writing: SM, JD, AAWR, SdAG; website construction and maintenance: NM.

**Conflicts of interest** None.

**Ethical standards** No specimens were collected during this research, which otherwise abided by the *Oryx* guidelines on ethical standards.

## References

- ALBERT, C., LUQUE, G.M. & COURCHAMP, F. (2018) The twenty most charismatic species. *PLOS ONE*, 13, e0199149.
- ASCENSAO, F., YOGUI, D.R., ALVES, M.H., ALVES, A.C., ABRA, F. & DESBIEZ, A.L.J. (2021) Preventing wildlife roadkill can offset mitigation investments in short-medium term. *Biological Conservation*, 253, 108902.
- AVGAN, B. (2009) Sighting of a jungle cat and the threats of its habitat in Turkey. *CAT news*, 50, 16.
- BAKER, D.J., GARNETT, S.T., O'CONNOR, J., EHMKE, G., CLARKE, R.H., WOJNARSKI, J.C.Z. & MCGEOCH, M.A. (2019) Conserving the abundance of nonthreatened species. *Conservation Biology*, 33, 319–328.
- BAMBARADENIYA, C.N.B. & AMARASINGHE, S. (2001) Wild cats in Sri Lanka. *CAT news*, 35, 18–19.
- BROOKE, Z.M., BIELBY, J., NAMBIAR, K. & CARBONE, C. (2014) Correlates of research effort in carnivores: body size, range size and diet matter. *PLOS ONE*, 9, e93195.
- BROWN, J.L., BENNETT, J.R. & FRENCH, C.M. (2017) SDMtoolbox 2.0: the next generation Python-based GIS toolkit for landscape genetic, biogeographic and species distribution model analyses. *PeerJ*, 5, e4095.
- BRUM, F.T., GRAHAM, C.H., COSTA, G.C., HEDGES, S.B., PENONE, C., RADELOFF, V.C. et al. (2017) Global priorities for conservation across multiple dimensions of mammalian diversity. *Proceedings of the National Academy of Sciences of the United States of America*, 114, 7641–7646.
- DAYAN, T., SIMBERLOFF, D., TCHERNOV, E. & YOM-TOV, Y. (1990) Feline canines: community-wide character displacement among the small cats of Israel. *The American Naturalist*, 136, 39–60.
- DE ALWIS, W.L.E. (1973) Status of Southeast Asia's small cats. In *The World's Cats Volume 1* (ed. R.L. Eaton), pp. 198–208. World Wildlife Safari, Winston, USA.
- DEPARTMENT OF AGRARIAN DEVELOPMENT (2011) *Watersheds of Sri Lanka*. 1st edition. Department of Agrarian Development, Colombo, Sri Lanka.
- DEUTZ, A., HEAL, G.M., NIU, R., SWANSON, E., TOWNSHEND, T., ZHU, T. et al. (2020) *Financing Nature: Closing the Global Biodiversity Financing Gap*. The Paulson Institute, The Nature Conservancy, and the Cornell Atkinson Center for Sustainability, USA.
- DIAZ, S., SETTELE, J., BRONDIZIO, E., NGO, H., AGARD, J., ARNETH, A. et al. (2019) Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science*, 366, 1327.
- DUCARME, F., LUQUE, G. & COURCHAMP, F. (2013) What are 'charismatic species' for conservation biologists? *BioSciences Master Reviews*, 1, 1–8.
- ELITH, J., PHILLIPS, S.J., HASTIE, T., DUDIK, M., CHEE, Y.E. & YATES, C.J. (2011) A statistical explanation of *MaxEnt* for ecologists (Report). *Diversity and Distributions*, 17, 43–57.
- FAURBY, S. & SVENNING, J.-C. (2015) Historic and prehistoric human-driven extinctions have reshaped global mammal diversity patterns. *Diversity and Distributions*, 21, 1155–1166.
- FERNANDO, P., SILVA, M.K.C.R.D., JAYASINGHE, L.K.A., JANAKA, H.K. & PASTORINI, J. (2021) First country-wide survey of the Endangered Asian elephant: towards better conservation and management in Sri Lanka. *Oryx*, 55, 46–55.
- FILIUS, J., VAN DER HOEK, Y., JARRIN, V.P. & VAN HOOFT, P. (2020) Wildlife roadkill patterns in a fragmented landscape of the Western Amazon. *Ecology and Evolution*, 10, 6623–6635.
- FRIMPONG, E.A. (2018) A case for conserving common species. *PLOS Biology*, 16, e2004261.
- GERNGROSS, P. (2014) Recent records of jungle cat in Turkey. *CAT News*, 61, 10–11.
- GITTLEMAN, J.L. & GOMPPER, M.E. (2005) Plight of predators: the importance of carnivores for understanding patterns of biodiversity and extinction risk. In *Ecology of Predator–Prey Interactions* (eds P. Barbosa & I. Castellanos), pp. 370–388. Oxford University Press, Oxford, UK.
- GLOBAL WILDLIFE CONSERVATION (2020) *Awe-Inspiring Small Cats*. Global Wildlife Conservation. [globalwildlife.org/project/small-cats](https://globalwildlife.org/project/small-cats) [accessed 6 June 2020].
- GRAY, T.N.E., TIMMINS, R.J., JATHANA, D., DUCKWORTH, J.W., BARAL, H. & MUKHERJEE, S. (2021) *Felis chaus* (amended version of 2016 assessment). In *The IUCN Red List of Threatened Species 2021*. [dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T8540A50651463.en](https://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T8540A50651463.en).
- HARRISON, D.L. (1968) *Felis chaus*. In *The Mammals of Arabia Volume II: Carnivora, Artiodactyla and Hyracoidea*, pp. 291–295. Ernest Benn Limited, London, UK.
- HATT, R.T. (1959) *Felis chaus furax*. In *The Mammals of Iraq*, p. 47. Museum of Zoology, University of Michigan, Ann Arbor, USA.
- HEPTNER, V.G. & SLUDSKII, A.A. (1992) *Mammals of the Soviet Union*. Volume 2 Part 2 Carnivora (Hyenas and Cats). Smithsonian Institution Libraries and The National Science Foundation, Washington, DC, USA.
- HIJMANS, R.J., CAMERON, S.E., PARRA, J.L., JONES, P.G. & JARVIS, A. (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25, 1965–1978.
- IUCN (2012) *IUCN Red List Categories and Criteria Version 3.1*. IUCN Species Survival Commission, Gland, Switzerland and Cambridge, UK.
- KALLE, R., RAMESH, T., QURESHI, Q. & SANKAR, K. (2013) The occurrence of small felids in Mudumalai Tiger Reserve, Tamil Nadu, India. *CAT news*, 58, 32–35.
- KARUNARATHNA, S., RANWALA, S., SURASINGHE, T. & MADAWALA, M. (2017) Impact of vehicular traffic on vertebrate fauna in Horton plains and Yala national parks of Sri Lanka: some implications for conservation and management. *Journal of Threatened Taxa*, 9, 9928–9939.
- KISSEL, R. & POSERINA, J. (eds) (2017) *Advanced math and statistics. In Optimal Sports Math, Statistics, and Fantasy*, pp. 103–135. Academic Press Books—Elsevier, Cambridge, USA.
- KITTLE, A.M. & WATSON, A.C. (2018) Small wildcats of Sri Lanka – some recent records. *CAT news*, 68, Autumn 2018, 9–12.
- KITTLE, A.M., WATSON, A.C., CUSHMAN, S. & MACDONALD, D. (2018) Forest cover and level of protection influence the island-wide distribution of an apex carnivore and umbrella species, the Sri Lankan leopard (*Panthera pardus kotiya*). *Biodiversity and Conservation*, 27, 235–263.
- MEEGASKUMBURA, S., MEEGASKUMBURA, M., PETHIYAGODA, R., MANAMENDRA-ARACHCHI, K. & SCHNEIDER, C.J. (2007) *Crociodura hikmiya*, a new shrew (Mammalia: Soricomorpha: Soricidae) from Sri Lanka. *Zootaxa*, 1665, 19–30.
- MENDELSSOHN, H. (1989) Felids in Israel. *CAT news*, 10, 2–4.
- MENDIS WICKRAMASINGHE, L.J., VIDANAPATHIRANA, D.R. & WICKRAMASINGHE, N. (2020) A new species of *Lankascincus* Greer, 1991 (Reptilia: Scincidae) from the Rakwana hills of Sri Lanka. *Taprobanica*, 9, 23–30.
- MEYER, C., KREFT, H., GURALNICK, R. & JETZ, W. (2015) Global priorities for an effective information basis of biodiversity distributions. *Nature Communications*, 6, 8221.
- MILLENNIUM ECOSYSTEM ASSESSMENT (2005) *Ecosystems and Human Well-being: Biodiversity Synthesis*. World Resources Institute, Washington, DC, USA.
- MININ, E.D., SLOTOW, R., HUNTER, L.T.B., POUZOLS, F.M., TOIVONEN, T., PETER, H.V. et al. (2016) Global priorities for national carnivore conservation under land use change. *Scientific Reports*, 6, 23814.

- MINISTRY OF ENVIRONMENT SRI LANKA (2012) *The National Red List 2012 of Sri Lanka; Conservation Status of the Fauna and Flora*. Ministry of Environment, Colombo, Sri Lanka.
- MINISTRY OF MAHAWELI DEVELOPMENT AND ENVIRONMENT (2016) *National Biodiversity Strategic Action Plan 2016–2022*. Biodiversity Secretariat, Ministry of Mahaweli Development and Environment, Colombo, Sri Lanka.
- MISHRA, R., GAUTAM, B., SHAH, S.K., SUBEDI, N., POKHERAL, C.P. & LAMICHHANE, B.R. (2020) Opportunistic records of jungle cat (*Felis chaus*) and their activity pattern in Koshi Tappu Wildlife Reserve, Nepal. *Nepalese Journal of Zoology*, 4, 50–55.
- MITHTHAPALA, S. (2016) *Obtaining a better distribution map of the Jungle Cat in Sri Lanka*. [junglecatsrilanka.com](http://junglecatsrilanka.com) [accessed 14 February 2021].
- MITHTHAPALA, S. (2018) An overview of wild cats of Sri Lanka Part II. *Loris*, 28, 37–47.
- MUTHUWATTA, L.P. & LIYANAGE, P.K.N.C. (2013) Impact of rainfall change on the agro-ecological regions of Sri Lanka. In *Proceedings of the International Conference on Climate Change Impacts and Adaptations for Food and Environment Security on Sustaining Agriculture Under Changing Climate*. Colombo, Sri Lanka, 30–31 July 2013. (eds H.P.M. Gunasena, H.A.J. Gunathilake, J.M.D. T. Everard, C.S. Ranasinghe & A.D. Nainanayake), pp. 59–66. Ministry of Environment and Renewable Energy, Lunuwila, Sri Lanka, and Ministry of Environment and Renewable Energy, New Delhi, India.
- PASCOE, E.L., MARCANTONIO, M., CAMINADE, C. & FOLEY, J.E. (2019) Modeling potential habitat for tick species in California. *Insects*, 10, 201.
- PATEL, K. (2011) Preliminary survey of small cats in Eastern Gujarat, India. *CAT news*, 54, 8–11.
- PHILLIPS, S.J., ANDERSON, R.P. & SCHAPIRE, R.E. (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190, 231–259.
- PHILLIPS, W.A. (1935) *Manual of the Mammals of Sri Lanka*. Colombo Museum, Colombo, Sri Lanka.
- PRAKASH, T.G.S., WIJERATNE, A. & FERNANDO, P. (2020) Human–elephant conflict in Sri Lanka: patterns and extent. *Gajah*, 51, 16–25.
- PRISAZHNYUK, B.E. & BELOUSOVA, A.E. (2007) Species account for *Felis chaus*. [biodat.ru/db/rb/rb.php?src=1&vid=389](http://biodat.ru/db/rb/rb.php?src=1&vid=389) [accessed 9 July 2019]. [In Russian]
- RAMESH, K. & MCGOWAN, P. (2009) On the current status of Indian Peafowl *Pavo cristatus* (Aves: Galliformes: Phasianidae): keeping the common species common. *Journal of Threatened Taxa*, 1, 106–108.
- R CORE TEAM (2019) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. [r-project.org](http://r-project.org) [accessed 7 February 2019].
- REDFORD, K.H., BERGER, J. & ZACK, S. (2013) Abundance as a conservation value. *Oryx*, 47, 157–158.
- RODRIGUEZ, J.D., PEREZ, A. & LOZANO, J.A. (2010) Sensitivity analysis of k-fold cross validation in prediction error estimation. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 32, 569–575.
- ROEMER, G., GOMPPER, M. & VAN VALKENBURGH, B. (2009) The ecological role of the mammalian mesocarnivore. *BioScience*, 59, 165–173.
- SANEI, A., MOUSAVI, M., RABIEE, K., KHOSRAVI, M.S., JULAAE, L., GUDARZI, F. et al. (2016) Distribution, characteristics and conservation of the jungle cat in Iran. *CAT news Special Issue*, 10, 51–55.
- SAVE FISHING CATS CONSERVATION PROJECT (2015) *Save Fishing Cats Conservation Project*. [savefishingcats.org](http://savefishingcats.org) [accessed 30 June 2019].
- SCHULTZ, M., TYRRELL, T.D. & EBENHARD, T. (2016) *The 2030 Agenda and Ecosystems – A Discussion Paper on the Links Between the Aichi Biodiversity Targets and the Sustainable Development Goals*. SwedBio at Stockholm Resilience Centre, Stockholm, Sweden.
- SCHWARTZ, A.L.W., SHILLING, F.M. & PERKINS, S.E. (2020) The value of monitoring wildlife roadkill. *European Journal of Wildlife Research*, 66, 18.
- SCOTT, J.M., DAVIS, F., CSUTI, B., NOSS, R., BUTTERFIELD, B., GROVES, C. et al. (1993) Gap analysis: a geographic approach to protection of biological diversity. *Wildlife Monographs*, 123, 3–41.
- SENEVIRATHNE, G., SAMARAWICKRAMA, V.A.M.P.K., WIJAYATHILAKA, N., MANAMENDRA-ARACHCHI, K., BOWATTE, G. et al. (2018) A new frog species from rapidly dwindling cloud forest streams of Sri Lanka–*Lankanectes pera* (Anura, Nyctibatrachidae). *Zootaxa*, 4461, 519–538.
- SHRESTHA, B.S., SUBEDI, N. & KANDEL, R.C. (2020) Jungle cat *Felis chaus* Schreber, 1777 (Mammalia: Carnivora: Felidae) at high elevations in Annapurna Conservation Area, Nepal. *Journal of Threatened Taxa*, 12, 15267–15271.
- SMALL CAT ADVOCACY AND RESEARCH (2017) *Small Cat Advocacy and Research*. [scar.lk](http://scar.lk) [accessed 6 October 2021].
- SMALL CAT ADVOCACY AND RESEARCH (2020) Each of the 25 squares represents a Sri Lankan small wild cat that was hit by a vehicle, snared, killed by dogs or killed by people in 2020. [instagram.com/p/CJeBsvZhEJl](https://www.instagram.com/p/CJeBsvZhEJl) [accessed 6 October 2021].
- SUDASINGHE, H. (2018) A new species of *Schistura* (Teleostei: Nemacheilidae) from the south-western lowlands of Sri Lanka. *Zootaxa*, 4422, 478–492.
- SUNQUIST, M. & SUNQUIST, F. (eds) (2002) Jungle cat (*Felis chaus*). In *Wild Cats of the World*, pp. 60–66. University of Chicago Press, Chicago, USA.
- SURVEYOR GENERAL, SURVEY DEPARTMENT (2016) *Sri Lanka land Use: Forests, Marshes, Home Gardens, Paddy Fields*. Surveyor General, Survey Department, Colombo, Sri Lanka.
- TENSEN, L. (2018) Biases in wildlife and conservation research, using felids and canids as a case study. *Global Ecology And Conservation*, 15, e00423.
- UNITED NATIONS (2015) *Transforming Our World: the 2030 Agenda for Sustainable Development*. [sdgs.un.org/2030agenda](https://www.sdg.un.org/2030agenda) [accessed 21 July 2020].
- USGS (UNITED STATES GEOLOGICAL SURVEY) (2018) *U.S. Geological Survey – Gap Analysis Project Species Habitat Maps CONUS\_2001*. [sciencebase.gov/catalog/item/527do83e4b0850ea0518326](https://sciencebase.gov/catalog/item/527do83e4b0850ea0518326) [accessed 18 April 2021].
- URBAN FISHING CAT CONSERVATION PROJECT (2020) *Urban Fishing Cat Conservation Project*. [fishingcats.lk](http://fishingcats.lk) [accessed 30 June 2019].
- WOLF, C. & RIPPLE, W.J. (2017) Range contractions of the world's large carnivores. *Royal Society Open Science*, 4, 170052.
- YAPA, A. & RATNAVIRA, G. (2013) *The Mammals of Sri Lanka*. Field Ornithology Group of Sri Lanka, Department of Zoology, University of Colombo, Colombo, Sri Lanka.