## VOL. XII.—1897.

THE

# JOURNAL OF LARYNGOLOGY,

RHINOLOGY, AND OTOLOGY;

AN ANALYTICAL RECORD OF CURRENT LITERATURE

RELATING TO

THE THROAT, NOSE, AND EAR.

PUBLISHED MONTHLY.

#### Kondon:

THE REBMAN PUBLISHING COMPANY, LIMITED,

SUCCESSORS TO F. J. REBMAN, 11, ADAM STREET, STRAND.

#### AMERICAN AGENTS:

THE F. A. DAVIS COMPANY, 1914-16, CHERRY STREET, PHILADELPHIA, PA.

ENTERED AT STATIONERS' HALL.

# THE JOURNAL OF LARYNGOLOGY, RHINOLOGY, AND OTOLOGY.

Founded in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN

#### EDITORS:

JOHN MACINTYRE, M.B., F.R.S.E.,
DUNDAS GRANT, M.A., M.D., F.R.C.S.Eng.,
ARTHUR SANDFORD, M.D., M.Ch.R.U.I.

W. MILLIGAN, M.D. (Manchester), Sub-Ed.
R. LAKE, F.R.C.S. (London), Managing Sub-Ed.

WITH THE CO-OPERATION OF

DRS. BARON (Bristol), BETTMAN (Rome), BOTEY (Barcelona),

CARTAZ (Paris), BRYSON DELAVAN (New York), DODD (Chicago),

DRAISPUL (St. Petersburg), GRAZZI (Florence), GUYE (Amsterdam),

HICGUET (Brussels), MIDDLEMASS HUNT (Liverfool), HUTCHISON (Glaszow),

JOAL (Paris), KARWOWSKI (Warsaw), KELLY (Glaszow),

LACOARRET (Toulouse), LICHTWITZ (Bordeaux),

LIEVEN (Aix-la-Chapelle), PROF. JOHN N. MACKENZIE (Baltimore),

PROF. MASSEI (Naples), MEYJES (Amsterdam), E. MEYER (Berlin),

HOLGER MYGIND (Copenhagen), PORTER (St. Louis),

STGEORGE REID (London), RUEDA (Madrid), SAJOUS (Paris),

SENDZIAK (Warsaw), SOTA (Seville), STCLAIR THOMSON (London),

WAGGETT (London), WOODS (Dublin), AND ZIEM (Dantzic).

## CONTENTS.

Dr. Norris Wolfenden
Societies' Meetings-
BRITISH LARYNGOLOGICAL ASSOCIATION.—Stenosis of External Auditory Meatus, Web Formation in the External Auditory Meatus, Syphilis of Nose, Lupus of Nose, Tresilian.—Syphilitic Laryngitis, Wilkin.—Frontal Empyema, Milligan.—Esthiomenic Ulcer, Jones.—Laryngeal Growth, Whistler.—Lupus of Larynx and Nose, Lake.—Discussion on Indications for, and Methods of, Operation for Post-Nasal Adenoids, Greville MacDonald and Dundas Grant Pages 298 to 315
BRADSHAW LECTURE (concluded) Pages 315 to 320  LETTSOMIAN LECTURE (concluded) Pages 321 to 330  INTERNATIONAL MEDICAL CONGRESS Page 331  BRITISH MEDICAL ASSOCIATION Page 332
Abstracts-
Diphtheria.—Infectiousness of, Dennig——Serum, Dobezynoski, Gottstein——Blood in, Engel ——Intubation in, Watson
Mouth.—Foot-and-Mouth Disease, Bussenius and Siegel.—Taste Fibres, Dixon—Ludwig's Angina, Ombridanne.—Tubercle of Tonsil, Pluder.—Saliva in Quantity, Sticher. Pages 335 to 337
Nose.—Pictures, Bergengruen—Aural Reflex, Collet.—Sinusitis, Combé.—Acute Coryza, Courtade.—Congenital Tumour of Face, Ewald.—Meyer's Tonsil, Goure.—Adenoids, Kahn.—Naso-Pharyngeal Autoscopy, Katzenstein.—Frontal Sinusitis, Thomas.—Phenacetin in Catarrh, Woodruff
Larynk.—Intubation, Bayeux.—Anatomical Examination, Benda — Ligno-Sulphite in Tubercle, Brannsfeld.—Neuroses, Collet.—Syphilis, Fink.—Sarcomata, Grand.—Innervation, Grossmann.—Primary Edema, Liaras.—Publica Laryngologica, Semon.—Iodide Edema, Stankowski
Ear.—Paralysis of Chorda, Alderton — Intra-Cranial Disease, Barr — Acute Syphilis, Crockett —Osteoma, Friedewald — Ocular Reflex, Laurens — Lumbar Puncture, Leutert — Traumatic Perforation, Somers
Reviews-
Dalby.—Contributions to Aural Surgery
MOURE.—Diseases of Naso-Pharynx and Pharynx
Literary NotePage 356
GUIDE TO ADVERTISEMENTS.
Bullock & Co.         vii.           Burroughs, Wellcome & Co.         vi.           Central London Throat Hospital         xiv.           Ingram & Royle.         vii., xi.           Lambert Pharmacal Co.         ii.           Mayer & Meltzer         xiii.           Pearl Life Assurance Company         iv.           Popelreuter         iv.           Rebman Publishing Company         vii., viii., ix.           Slater, Bros. & Crossley         x.           Woodruff         xiii

# "TABLOIDS" and "SOLOIDS"

For Laryngological Use.

A "TABLOID," when intended for the treatment of the throat and voice, is compressed hard so as to dissolve slowly and produce a prolonged action on the mucous membrane. There are no sugary or gummy substances present to interfere with the direct application of the medicament to the affected surfaces. Placed upon the tongue, a "Tabloid" is immediately acted upon by the saliva forming a highly-concentrated solution, which remains in continuous contact with the affected mucous membrane until the whole of the "Tabloid" is dissolved.

#### RED GUM "TABLOIDS."



Red Gum, an exudation from the Australian *Eucalyptus rostrata*, is a mild styptic and antiseptic, and an excellent local and general astringent. The *Practitioner* reports:—"When a Red Gum 'Tabloid' is allowed to dissolve in the mouth, the continuous action of the drug on the throat is fully secured. Great care has been exercised so to adjust the relative portions of the ingredients that the full therapeutic effect of the drug may be obtained without offending the palate or affecting the appetite."

Supplied in boxes of 30 and bottles of 100 "Tabloids" at 5d. and 1s. each.

#### TAR "TABLOIDS."

In a paper, entitled "A Mode of Administering Tar," which appeared in the Medical Press and Circular, a distinguished physician reports as follows:—

"I have recently had my attention called to 'Tabloids' of Tar, containing one grain; and these, I think, constitute by far the best mode of administering the drug. They contain the whole of the constituents of tar and are palatable and speedily disintegrate. The new mode of administration is three or four every four hours; but a better plan is to suck one frequently—in fact, every time the cough is troublesome."

Supplied in bottles of 50 and 100 at 5d. and 9d. each.

#### EUCAINE HYDROCHLORIDE "SOLOIDS." (Gr. 1 and Gr. 5.

**Eucaine** is a synthetic compound allied to cocaine and having a similar chemical composition. Its local anæsthetic action is not unlike that of cocaine, and it has not been found to produce untoward after-effects. The anæsthesia of eucaine develops more slowly than that of cocaine, but lasts longer. In nose and throat surgery, where a large quantity of anæsthetic has occasionally to be used, it is found to act efficiently, and to be without the poisonous effect previously produced in the same patients by similar doses of cocaine.

Owing to the sparing solubility of eucaine in cold water, and to the fact that it freely crystallizes from hot water on cooling, it is necessary that solutions for anæsthetic purposes should be made only at the moment they are required. For this purpose Eucaine Hydrochloride "Soloids" are most convenient. By dissolving one or more of these "Soloids" in definite quantities of warm water, solutions of any desired strength can be instantly prepared, and the operator is assured of obtaining the full anæsthetic effect of the drug. In laryngological practice eucaine may be used, either as a spray or as an injection, in the same strength as cocaine.

Eucaine Hydrochloride "Soloids" are supplied in bottles of 25 "Soloids" (1 gr. and 5 gr.) at 1s. 8d. and 7s. per bottle.

## BURROUGHS, WELLCOME & Co.,

SNOW HILL BUILDINGS, LONDON.

Australasian Branch: 108, PITT STREET, SYDNEY, N.S.W.

Cable and Telegraphic Address: "BURCOME, LONDON."