

BULLETIN

OF

ENTOMOLOGICAL RESEARCH

ISSUED BY THE
ENTOMOLOGICAL RESEARCH COMMITTEE
APPOINTED BY THE COLONIAL OFFICE.

EDITOR: THE SCIENTIFIC SECRETARY.

VOL. III.

LONDON:

SOLD BY

DULAU & Co., Ltd., 37, SOHO SQUARE, W.

1912.

CONTENTS.

ORIGINAL ARTICLES.

| | PAGE. |
|--|-------|
| AUSTEN, ERNEST E. | |
| A new species of <i>Stygeromyia</i> (Family Muscidae, Sub-family Stomoxydinae) from the East Africa Protectorate | 97 |
| Notes on African blood-sucking midges (Family Chironomidae, Sub-family Ceratopogoninae) with descriptions of new species... .. | 99 |
| New African Tabanidae.—Part I | 113 |
| New African Tabanidae.—Part II | 329 |
| New African Tabanidae.—Part III | 399 |
| A new species of <i>Hippobosca</i> from Northern Rhodesia | 417 |
| CARPENTER, PROF. GEORGE H. | |
| A new West African Spring-tail | 79 |
| CUMMINGS, BRUCE F. | |
| Anoplura from African hosts | 393 |
| DEWITZ, DR. J. | |
| The bearing of physiology on economic entomology | 343 |
| DURRANT, JOHN HARTLEY. | |
| Notes on Tineina bred from cotton-bolls | 203 |
| EDWARDS, F. W. | |
| A synopsis of the species of African Culicidae other than <i>Anopheles</i> ... | 1 |
| A key for determining the African species of <i>Anopheles</i> (sensu lato) ... | 241 |
| Revised keys to the known larvæ of African Culicinae | 373 |
| FELL, T. E. | |
| Notes on Tsetse-flies and on prophylactic measures against Sleeping Sickness in the Western Province of Ashanti | 227 |
| (28784—2.) Wt. P 11—21 1000. 3/13. D & S. | |

| | PAGE. |
|---|-------|
| GOWDEY, C. C. | |
| On the utilisation of an indigenous African silkworm (<i>Anaphe infracta</i> , Wlsm.) in Uganda | 269 |
| HIRST, STANLEY. | |
| On a new species of mite (<i>Tarsonemus</i>) injurious to sugar-cane in Barbados | 325 |
| On two new parasitic Acari of the genus <i>Leiognathus</i> , Cn. (Gamasidae) ... | 369 |
| HOWARD, C. W. | |
| Insects directly or indirectly injurious to man and animals in Mozambique, East Africa | 211 |
| INGRAM, DR. A. | |
| Notes on the mosquitos observed at Bole, Northern Territories, Gold Coast | 73 |
| KING, HAROLD H. | |
| Observations on the occurrence of <i>Glossina</i> in the Mongalla Province of the Anglo-Egyptian Sudan | 89 |
| LLOYD, LL. | |
| Notes on <i>Glossina morsitans</i> in Northern Rhodesia | 95 |
| Notes on <i>Glossina morsitans</i> , Westw., in the Luangwa Valley, Northern Rhodesia | 233 |
| McCONNELL, DR. ROBERT E. | |
| Notes on the occurrence and habits of <i>Glossina fuscipes</i> in Uganda... .. | 55 |
| MACFIE, DR. J. W. SCOTT. | |
| Experiments and observations upon <i>Glossina palpalis</i> | 61 |
| MOISER, DR. BERNARD. | |
| Notes on the haunts and habits of <i>Glossina tachinoides</i> , near Geidam, Bornu Province, Northern Nigeria | 195 |
| NEAVE, S. A. | |
| Notes on the blood-sucking insects of Eastern Tropical Africa | 275 |
| NEWSTEAD, PROF. R. | |
| On the characteristics of the newly-discovered tsetse-fly, <i>Glossina austeni</i> , Newstead; with descriptions of the genital armature of <i>Glossina fusi-</i> <i>pleuris</i> , Austen, and <i>Glossina longipennis</i> , Corti | 355 |
| Notes on <i>Phlebotomus</i> , with descriptions of new species.—Part I. | 361 |
| NICHOLLS, LUCIUS. | |
| The transmission of pathogenic micro-organisms by flies in St. Lucia ... | 81 |
| Some observations on the bionomics and breeding-places of <i>Anopheles</i> in Saint Lucia, British West Indies | 251 |

CONTENTS.

v

| | PAGE. |
|--|-------|
| POLLARD, DR. J. | |
| Notes on the tsetse-flies of Muri Province, Northern Nigeria | 219 |
| SIMPSON, DR. JAS. J. | |
| Entomological Research in British West Africa. III.—Southern Nigeria ... | 137 |
| STANTON, A. T. | |
| On the changes which occur in certain characters of <i>Anopheles</i> larvae in the course of their growth | 387 |
| SUMMERS, SOPHIA L. M. | |
| A new species of <i>Phlebotomus</i> from South America | 209 |

MISCELLANEOUS.

| | |
|---|--------------------|
| Collections received | 109, 225, 339, 419 |
| Tabanidae attracted by Scale-insects | 223 |

PLATES.

| | PAGE. |
|--|-------------|
| I. Wings of African <i>Culicoides</i> | to face 108 |
| II. to V. Cattle and Views in Southern Nigeria | " 194 |
| VI. to IX. Breeding-places of <i>Anopheles</i> in St. Lucia, B.W.I. | " 268 |
| X. and XI. African Tabanidae | " 324 |

MAPS.

| | PAGE. |
|---|------------|
| Map showing distribution of <i>Glossina palpalis</i> and <i>G. morsitans</i> in Southern Sudan | to face 92 |
| Map of S. Nigeria showing distribution of <i>Glossina</i> | " 194 |
| Sketch-map of R. Wobe, N. Nigeria, showing localisation of <i>Glossina tachinoides</i> | " 202 |
| Sketch-map of Muri Province, N. Nigeria, showing distribution of <i>Glossina</i> | " 221 |

ILLUSTRATIONS IN THE TEXT.

| | PAGE. |
|--|-------|
| Terminal segments of female of: <i>Culex tigripes</i> var. <i>fuscus</i> ; <i>Mimomyia hispida</i> ; <i>Ochlerotatus cumminsi</i> | 5 |
| <i>Stegomyia fraseri</i> , sp. n., head, thorax and apex of abdomen of ♀ | 11 |
| <i>Ochlerotatus longipalpis</i> , Grünb., head, thorax and apex of abdomen of ♀ | 18 |
| <i>Ochlerotatus fascipalpis</i> , sp. n. | 19 |
| <i>Culex ager</i> , Giles, var. n. <i>ethiopicus</i> , wing | 30 |
| Head and thorax of ♂ of:— | |
| <i>Uranotaenia connali</i> , sp. n. | 40 |
| ,, <i>ornata</i> , Theo. | 43 |
| <i>Eretmopodites quinquevittatus</i> , Theo. | 46 |
| <i>Eretmopodites chrysogaster</i> , Graham, larva and pupa | 47 |
| <i>Chaoborus pallidipes</i> , Theo., head and thorax of ♀ | 48 |
| <i>Glossina palpalis</i> , diagram of vesica | 56 |
| <i>Mansonioides uniformis</i> , Theo., early stages | 76 |
| <i>Isotoma fasciata</i> , sp. n., side view, ocelli and post-antennal organ, mandible, tip of fore-foot with claw, tibio-tarsus of hind leg, catch, dentes and mucrones | 80 |
| <i>Drosophila melanogaster</i> , Mg. | 82 |
| <i>Sarcophaga aurifinis</i> , Walk. | 85 |
| <i>Oscinis pallipes</i> , Lw. | 86 |
| <i>Limosina punctipennis</i> , Wd. | 87 |
| Chalcid ovipositing in larva of <i>Sarcophaga</i> | 88 |
| <i>Glossina morsitans</i> : preparation of salivary glands and proboscis (semi- diagrammatic) | 96 |
| <i>Silvius fallax</i> , Aust., ♀ | 114 |
| <i>Cadicera speciosa</i> , Aust., ♀ | 120 |
| <i>Pangonia comata</i> , Aust., ♂ | 123 |
| <i>Subpangonia grahami</i> , Aust. ♀ | 127 |
| <i>Tabanus canofasciatus</i> , Aust., ♀ | 129 |
| ,, <i>tenuipalpis</i> , Aust., ♀ | 131 |
| <i>Hippocentrum murphyi</i> , Aust., ♀ | 135 |
| <i>Gelechia gossypiella</i> , Sndrs., ♂ | 204 |
| <i>Pyroderces simplex</i> , Wlsm., ♀ | 206 |
| ,, <i>rileyi</i> , Wlsm., ♀ | 207 |
| <i>Phlebotomus rostrans</i> , Summers; wing; genitalia, ♂; maxillary palp, ♀ | 209 |
| Larvae of <i>Anopheles (Nyssorhynchus)</i> | 252 |
| Pupa of <i>Anopheles albimanus</i> | 254 |
| Nest of <i>Anopheles infracta</i> | 270 |
| Soaking nests of <i>Anopheles</i> to soften hairs | 272 |

| | Page. |
|--|-------|
| <i>Tarsonemus spinipes</i> , Hirst, ♂ ; dorsal and ventral views | 326 |
| " " " ♀ ; ventral view | 327 |
| Male armatures of:— | |
| <i>Glossina austeni</i> , Newst. | 357 |
| " <i>fuscipleuris</i> , Aust. | 358 |
| " <i>longipennis</i> , Corti | 360 |
| Palpi of <i>Phlebotomus</i> | 361 |
| Antennae of <i>Phlebotomus</i> | 362 |
| Wings of <i>Phlebotomus</i> | 364 |
| <i>Leiognathus creightoni</i> , Hirst, ♀ ; dorsal and ventral aspects | 370 |
| " <i>liberiensis</i> , Hirst, ♀ ; ventral aspect | 371 |
| <i>Stegomyia sugens</i> , Wied. ; larva | 375 |
| <i>Ochlerotatus nigeriensis</i> , Theo. ; larva | 377 |
| <i>Mansonioides africanus</i> , Theo. ; larva | 377 |
| <i>Aëdomyia catasticta</i> , Knab ; larva | 378 |
| " " " " | 379 |
| <i>Culex ager</i> , Giles, var. <i>ethiopicus</i> , Edw. ; larva | 380 |
| " <i>annulioris</i> , Theo. ; larva | 382 |
| " <i>invidiosus</i> , Theo. ; larva | 382 |
| <i>Ingramia malfeyti</i> , Newst. ; larva | 384 |
| Head of <i>Anopheles</i> larva showing clypeal and frontal hairs | 388 |
| <i>Anopheles albirostris</i> ; larva ; head and thorax, showing hairs, 388 ; abdominal segments, showing leaflets and papilla, 389 ; head, showing change after first moult, 389 ; cockade or "palmate hair," 389 ; head and thorax of full-grown larva | 390 |
| <i>Neohaematopinus heliosciuri</i> , Cummings, ♂ | 394 |
| <i>Polyplax otomydis</i> , Cummings, ♀ | 396 |
| <i>Haematopota grahami</i> , Aust., ♀ | 405 |
| " <i>rubens</i> , Aust., ♀ | 410 |
| " <i>beringeri</i> , Aust., ♀ | 413 |
| " <i>crudelis</i> , Aust., ♀ | 415 |