

## SHORT CONTRIBUTIONS

### MARTINGALES AND TAIL PROBABILITIES

BY HANS U. GERBER

At the twenty-eighth Actuarial Research Conference of the Society of Actuaries, WILLMOT and LIN (1993) presented a paper whose central result is a bound on the tail probability of a random sum. In the subsequent discussion, Professor Bühlmann raised the question, if this bound could be derived by martingale methods. The purpose of this note is to show how it can be done.

We consider a random variable of the form

$$S = X_1 + \dots + X_N.$$

Here the random variables  $N, X_1, X_2, \dots$  are independent, and the  $X_k$ 's are assumed to be positive and identically distributed; their common distribution function is denoted by  $F(x)$ .

Let

$$p_k = \Pr(N = k), \quad k = 0, 1, \dots$$

We assume the existence of a number  $\phi, 0 < \phi < 1$ , with

$$(1) \quad \Pr(N > k | N \geq k) \leq \phi \quad \text{for } k = 1, 2, \dots$$

and a positive number  $r$  with

$$(2) \quad \phi \cdot \int_0^\infty e^{rx} dF(x) \leq 1$$

(if  $F(x)$  is sufficiently regular, we might choose the value of  $r$  for which equality holds). Then the result of Willmot and Lin is that

$$\Pr(S \geq x) \leq \frac{1 - p_0}{\phi} \cdot e^{-rx}$$

for any  $x > 0$ .

For the following proof we introduce

$$S_k = X_1 + \dots + X_k$$

and

$$Y_k = \begin{cases} e^{rS_k} & \text{if } N \geq k \\ 0 & \text{if } N < k. \end{cases}$$

We note the recursive relationship

$$Y_k = Z_k \cdot Y_{k-1}, \quad k = 1, 2, \dots$$

with

$$Z_k = \begin{cases} e^{rX_k} & \text{if } N \geq k \\ 0 & \text{if } N < k. \end{cases}$$

According to (1) and (2), the conditional expectation of  $Z_{k+1}$  (given  $N \geq k$ ) is less than or equal to 1, which shows that the sequence  $Y_1, Y_2, \dots$  is a supermartingale.

If we stop it at time

$$T = \min \{k : S_k \geq x \text{ or } N < k\}$$

it follows that, given  $N \geq 1$  and  $X_1$ ,

$$Y_1 \geq E[Y_T | N \geq 1, X_1]$$

or

$$e^{rX_1} \geq E[e^{rS_T} 1_{\{S \geq x\}} | N \geq 1, X_1] \geq e^{rx} \cdot \Pr(S \geq x | N \geq 1, X_1).$$

Then we get

$$\begin{aligned} \Pr(S \geq x) &= (1 - p_0) \cdot E[\Pr(S \geq x | N \geq 1, X_1)] \\ &\leq (1 - p_0) \cdot E[e^{rX_1} \cdot e^{-rx}] \\ &\leq \frac{1 - p_0}{\phi} \cdot e^{-rx}, \end{aligned}$$

which completes the proof.

#### REFERENCE

- WILMOT, G. E. and LIN X. (1993) *Lundberg bounds on the tails of compound distributions*. Research Report 93-14, Institute of Insurance and Pension Research. University of Waterloo. Forthcoming in the Journal of Applied Probability.

HANS U. GERBER

*Ecole des HEC, University of Lausanne, CH-1015 Lausanne.*