



COMPOSITIO MATHEMATICA

Cremona transformations and derived equivalences of K3 surfaces

Brendan Hassett and Kuan-Wen Lai

Compositio Math. **154** (2018), 1508–1533.

[doi:10.1112/S0010437X18007145](https://doi.org/10.1112/S0010437X18007145)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



Cremona transformations and derived equivalences of K3 surfaces

Brendan Hassett and Kuan-Wen Lai

ABSTRACT

We exhibit a Cremona transformation of \mathbb{P}^4 such that the base loci of the map and its inverse are birational to K3 surfaces. The two K3 surfaces are derived equivalent but not isomorphic to each other. As an application, we show that the difference of the two K3 surfaces annihilates the class of the affine line in the Grothendieck ring of varieties.

Introduction

Let X be a smooth complex projective variety that is *rational*, that is, admits a birational map $\varrho : \mathbb{P}^r \dashrightarrow X$ where $r = \dim(X)$. The map ϱ blows up various subvarieties of \mathbb{P}^r – to what extent are these determined by X ? We can always precompose ϱ by a birational automorphism of \mathbb{P}^r (i.e., a *Cremona transformation*) so we must take this into account.

For small dimensions these subvarieties are determined by X . When $r = 1$, ϱ extends to an isomorphism; if $r = 2$, ϱ is resolved by blowing up points in \mathbb{P}^2 . The case of 3-folds was analyzed by Clemens and Griffiths [CG72]. We may assume that ϱ (respectively, ϱ^{-1}) is resolved by blowing up a finite number of points and non-singular irreducible curves; let C_1, \dots, C_k (respectively, D_1, \dots, D_l) denote those of positive genus. Comparing the Hodge structures on middle cohomology groups using the blow-up formula, we obtain an isomorphism of principally polarized abelian varieties:

$$J(C_1) \times \cdots \times J(C_k) \simeq J(X) \times J(D_1) \times \cdots \times J(D_l).$$

The factors are Jacobians of curves and the intermediate Jacobian of X . Principally polarized abelian varieties admit unique decompositions into irreducible factors and the Jacobian of a curve is irreducible with respect to the natural polarization. It follows that $J(X)$ can be expressed as a product of Jacobians of curves C_{i_1}, \dots, C_{i_t} , $\{i_1, \dots, i_t\} \subset \{1, \dots, k\}$, and these curves are determined up to isomorphism by the Torelli theorem.

Therefore, we focus on 4-folds and their middle cohomology. Suppose that a smooth projective surface Σ is contained in the base locus of ϱ . The blow-up formula gives a homomorphism of Hodge structures

$$\beta : H^2(\Sigma, \mathbb{Z})(-1) \rightarrow H^4(X, \mathbb{Z});$$

can we recover Σ from its image? Keeping track of divisor classes of Σ is complicated, as they might disappear under subsequent blow-downs. Thus all we can expect to recover is the transcendental cohomology $T(\Sigma) \subset H^2(\Sigma, \mathbb{Z})$.

Received 14 February 2017, accepted in final form 28 November 2017, published online 28 May 2018.

2010 Mathematics Subject Classification 14E07, 14J28 (primary).

Keywords: Cremona transformations, derived equivalences, K3 surfaces.

This journal is © [Foundation Compositio Mathematica](#) 2018.

Mukai and Orlov [Orl97] have shown that K3 surfaces offer many examples of non-birational surfaces R and \hat{R} with $T(R) \simeq T(\hat{R})$ as integral Hodge structures. These are explained through the notion of *derived equivalence*. There are cubic 4-folds [Has16] whose Hodge structures show the trace of several derived equivalent K3 surfaces. However, these are not known to be rational. Nevertheless, this raises the following question.

Question 0.1. Let R and \hat{R} be derived equivalent K3 surfaces. Do there exist smooth projective 4-folds X , P , and \hat{P} and birational maps

$$\varrho : P \dashrightarrow X, \quad \hat{\varrho} : \hat{P} \dashrightarrow X,$$

such that R and \hat{R} are birational to components of the base loci of ϱ and $\hat{\varrho}$ respectively, and the induced

$$\beta : H^2(R, \mathbb{Z})(-1) \rightarrow H^4(X, \mathbb{Z}), \quad \hat{\beta} : H^2(\hat{R}, \mathbb{Z})(-1) \rightarrow H^4(X, \mathbb{Z})$$

induce an isomorphism $T(R) \simeq T(\hat{R})$?

In other words, are derived equivalences of K3 surfaces induced by birational maps? It makes sense to start with the case where $P \simeq \hat{P} \simeq \mathbb{P}^4$. Are derived equivalences of K3 surfaces induced by Cremona transformations?

This last question may be too ambitious, as the base loci of Cremona transformations are highly constrained. According to Crauder and Katz [CK89], the Cremona transformation of \mathbb{P}^4 which can be resolved by blowing up along a smooth and irreducible surface $S \subset \mathbb{P}^4$ occurs as one of the following two cases:

- (1) S is a quintic elliptic scroll $S = \mathbb{P}_C(E)$, where C is an elliptic curve and E is a rank-two vector bundle with $e = -\deg(\wedge^2 E) = -1$;
- (2) S is a degree-10 determinantal surface given by the vanishing of the 4×4 minors of a 4×5 matrix of linear forms.

Here we present an example where derived equivalences of K3 surfaces are explained through Cremona transformations, and offer further evidence that such examples are quite rare. We can explain derived equivalences among degree-12 K3 surfaces in this way; however, we do not know how to realize derived equivalences of higher-degree K3 surfaces.

Our construction gives new examples of zero-divisors in the Grothendieck ring of complex algebraic varieties. The difference of each derived equivalent pair is non-vanishing in the ring and annihilated by the class of the affine line. The first example in this direction is given by the Pfaffian–Grassmannian Calabi–Yau 3-folds [Bor15, Mar16]. Other examples include Calabi–Yau 3-folds from Grassmannians of type G_2 [IMOU16a, Kuz16]. Kuznetsov and Shinder [KS17] have formulated general conjectures relating derived equivalence to zero-divisors in the Grothendieck ring; our example is an instance of [KS17, Conjecture 1.6]. The relationship between zero-divisors in the Grothendieck ring and approaches to the rationality of cubic 4-folds is discussed in [GS14].

Section 1 presents preliminary results on Cremona transformations with singular base loci. The construction of our rational map is in § 2 and we verify the non-trivial derived equivalence in § 3. We apply the results to study the Grothendieck ring in § 4. Section 5 shows these constructions do not admit obvious extensions through a generalization of the classification of Crauder and Katz; the underlying computations are also used to analyze the maps defined in § 2.

1. Cremona transformation with singular base locus

1.1 Terminology and notation

A Cremona transformation of \mathbb{P}^n is a birational map $f : \mathbb{P}^n \dashrightarrow \mathbb{P}^n$. Its base locus $\text{Bs}(f)$ is the subscheme where f is undefined.


Throughout this paper, we consider the Cremona transformation $f : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$ with base locus resolved by blowing up an irreducible surface S , with singular locus consisting of *transverse double points*, which means a point where the surface has two smooth branches meeting transversally.

Suppose S has δ transverse double points which form a subset $\Delta \subset \mathbb{P}^4$. The blow-up of \mathbb{P}^4 along S can be factored as follows.

- (1) Blow up \mathbb{P}^4 along Δ , introducing δ exceptional divisors E_1, \dots, E_δ isomorphic to \mathbb{P}^3 . Let P denote the resulting 4-fold and S' the proper transform of S , which is now smooth.
- (2) Blow up P along S' to obtain P' . Let E denote the resulting exceptional divisor and E'_1, \dots, E'_δ the proper transforms of the first group of exceptional divisors. Each E'_i is isomorphic to $E_i \simeq \mathbb{P}^3$ blown up along two skew lines $Q'_i, Q''_i \subset E_i$.
- (3) Each E'_i is a \mathbb{P}^1 -bundle over $\mathbb{P}^1 \times \mathbb{P}^1$. Indeed, through each $p \in E_i$ not on Q'_i and Q''_i passes a unique line l intersecting Q'_i and Q''_i . The bundle map is given by $p \mapsto (l \cap Q'_i, l \cap Q''_i) \in Q'_i \times Q''_i$. Blow down each E'_i to $\mathbb{P}^1 \times \mathbb{P}^1$. The resulting X is isomorphic to $\text{Bl}_S \mathbb{P}^4$.

Remark. The blow-up $X \rightarrow \mathbb{P}^4$ has a quadric surface $Q_i, i = 1, \dots, \delta$, over each transverse double point of S . Then P' is obtained as the blow-up of X along these quadrics.

Let $\pi_1 : X \rightarrow \mathbb{P}^4$ be the blow-up along S and $\pi_2 : X \rightarrow \mathbb{P}^4$ the resolution of f so that $\pi_2 = \pi_1 \circ f$. We organize these maps into a diagram:



Note that, by the definition of blow-up, X is exactly the graph of f . Let L (respectively, M) denote the divisor of the hyperplane class of the left (respectively, right) \mathbb{P}^4 . We also use L (respectively, M) to denote its pullbacks to X, P and P' (respectively, X and P').

It's clear that $L^4 = 1$. We have

$$M^4 = 1 \tag{1.1}$$

on X as f is birational. We define n by

$$L^3 M = n \tag{1.2}$$

and ξ by

$$LM^3 = \xi. \tag{1.3}$$

We may interpret n and ξ as the degrees of the homogeneous forms inducing f and f^{-1} , respectively. Define m as the multiplicity of S in the base locus. It is clear that

$$M = nL - m(E + 2\sum_{i=1}^{\delta} E'_i) \quad \text{on } P'.$$

Since a non-degenerate subvariety in projective space has degree greater than one, the linear system in $|M|$ inducing $P' \rightarrow \mathbb{P}^4$ must be complete. Thus we have

$$h^0(P', M) = 5. \tag{1.4}$$

We use these equations in our classification of Cremona transformations below.

1.2 Computing the intersection numbers

Let Σ denote the normalization of S and let K_Σ be its canonical class. Then the blow-up of Σ along the preimage of the transverse double points is isomorphic to S' . We denote by C a general sectional curve of S and also its preimages in Σ and S' . Let $d = C^2 = \text{deg } S$. Note that $E_i \cap S' = Q'_i \cup Q''_i$ are exactly the exceptional curves on S' over the i th double point.

LEMMA 1.1. *We have $LE'_i = 0$. We also have $E^3E'_i = -4$, $E^2E'^2_i = 2$, $EE'^3_i = 0$ and $E'^4_i = -1$.*

Proof. First, $LE'_i = 0$ since their intersection is empty.

Recall that E'_i is isomorphic to $E_i \simeq \mathbb{P}^3$ blown up at skew lines Q'_i and Q''_i . Write $\text{Pic}(E'_i) = \langle H, \tilde{Q}', \tilde{Q}'' \rangle$ where H is the polarization from \mathbb{P}^3 , while \tilde{Q}' and \tilde{Q}'' are the exceptional divisors over the lines. We clearly have $\tilde{Q}'\tilde{Q}'' = 0$ and $\tilde{Q}'H^2 = \tilde{Q}''H^2 = 0$. Since $N_{Q'_i/\mathbb{P}^3} = \mathcal{O}_{Q'_i}(1) \oplus \mathcal{O}_{Q'_i}(1)$, writing $\zeta = c_1(\mathcal{O}_{\mathbb{P}(N_{Q'_i/\mathbb{P}^3})}(1))$, we obtain $\zeta^2 + 2H\zeta = 0$ in the Chow group of $\tilde{Q}' = \mathbb{P}(N_{Q'_i/\mathbb{P}^3})$. We have $\tilde{Q}'|_{\tilde{Q}'} = -\zeta$, so that

$$\tilde{Q}'^2H = -\zeta H = -1, \quad \tilde{Q}'^3 = \zeta^2 = -2H\zeta = -2.$$

We have $N_{E'_i/P'} = \mathcal{O}(-H)$ and $E|_{E'_i} = \tilde{Q}' + \tilde{Q}''$. Thus we obtain

$$\begin{aligned} E^3E'_i &= (\tilde{Q}' + \tilde{Q}'')^3 = \tilde{Q}'^3 + \tilde{Q}''^3 = -4, \\ E^2E'^2_i &= (\tilde{Q}' + \tilde{Q}'')^2(-H) = 2, \\ EE'^3_i &= (\tilde{Q}' + \tilde{Q}'')(-H)^2 = 0, \\ E'^4_i &= (-H)^3 = -1. \end{aligned} \tag{□}$$

LEMMA 1.2. *The intersection numbers involving L and E are:*

- (1) $L^3E = 0$ and $L^2E^2 = -d$;
- (2) $LE^3 = -5d - K_\Sigma C$;
- (3) $E^4 = -15d - 5K_\Sigma C - c_2(\Sigma) + 6\delta$;
- (3') $E^4 = d^2 - 25d - 10K_\Sigma C - K_\Sigma^2 + 4\delta$.

Proof. $L^3E = 0$ since a general line does not intersect S . We have $L^2E^2 = -\text{deg } S' = -d$.

Assume that $C = S \cap L$ for some hyperplane $L \simeq \mathbb{P}^3$. Then $LE^3 = s(C, L)_0$, the zeroth Segre class of C in L , which equals $[c(N_{C/L})^{-1}]_0 = [c(C)c(\iota^*T_{\mathbb{P}^3})^{-1}]_0 = [([C] - K_\Sigma C - C^2)([C] - 4d)]_0 = -5d - K_\Sigma C$.

We have $E^4 = -s(S', P)_0 = -[c(N_{S'/P})^{-1}]_0 = -[c(S')c(P)|_{S'}^{-1}]_0$. Let $\epsilon : P \rightarrow \mathbb{P}^4$ be the blow-up. The blow-up formula for Chern classes gives

$$\begin{aligned} c(P) &= \epsilon^*c(\mathbb{P}^4) + (1 + \sum_i E_i)(1 - \sum_i E_i)^4 - 1 \\ &= ([P] + L)^5 + \sum_i (-3E_i + 2E_i^2 + 2E_i^3 - 3E_i^4). \end{aligned}$$

Thus we have

$$\begin{aligned} c(P)|_{S'} &= ([S'] + 5C + 10C^2) + \sum_i[-3(Q'_i + Q''_i) + 2(Q'^2_i + Q''^2_i)] \\ &= [S'] + 5C - 3\sum_i(Q'_i + Q''_i) + 10d - 4\delta \end{aligned}$$

and also

$$c(P)|_{S'}^{-1} = [S'] - 5C + 3\sum_i(Q'_i + Q''_i) + 15d - 14\delta.$$

Let $\tau : S' \rightarrow \Sigma$ be the blow-up. Then we have

$$c(S') = [S'] - \tau^*K_\Sigma - \sum_i(Q'_i + Q''_i) + c_2(\Sigma) + 2\delta.$$

Multiply the results to get $E^4 = -15d - 5K_\Sigma C - c_2(\Sigma) + 6\delta$.

Another expression for E^4 is derived from

$$-[c(N_{S'/P})^{-1}]_0 = c_2(N_{S'/P}) - c_1(N_{S'/P})^2.$$

We have $c_2(N_{S'/P}) = d^2 - 4\delta$. On the other hand,

$$\begin{aligned} c_1(N_{S'/P}) &= c_1(T_P)|_{S'} - c_1(T_{S'}) \\ &= -(-5L + 3\sum_i E_i)|_{S'} - (-K_{S'}) \\ &= 5C + \tau^*K_\Sigma - 2\sum_i(Q'_i + Q''_i), \end{aligned}$$

hence we deduce

$$c_1(N_{S'/P})^2 = 25d + 10K_\Sigma C + K_\Sigma^2 - 8\delta$$

and also

$$\begin{aligned} E^4 &= (d^2 - 4\delta) - (25d + 10K_\Sigma C + K_\Sigma^2 - 8\delta) \\ &= d^2 - 25d - 10K_\Sigma C - K_\Sigma^2 + 4\delta. \end{aligned} \quad \square$$

2. Construction of our example

In this section, we use Mukai's construction [Muk88] to produce an explicit example of a degree-12 K3 surface $R \subset \mathbb{P}^7$ together with three points $p_1, p_2, p_3 \in R$. This example helps us prove the following theorem.

THEOREM 2.1. *Let $R \subset \mathbb{P}^7$ be a generic K3 surface of degree 12 and $\Pi := \{p_1, p_2, p_3\} \subset R$ a generic triple of points.*

- (1) *Projection from Π maps R to a surface $S \subset \mathbb{P}^4$ with three transverse double points.*
- (2) *The complete linear system M of quartics vanishing along S cuts out S scheme-theoretically.*
- (3) *M induces a birational map $f : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$.*
- (4) *The base locus of the inverse f^{-1} is also a projection of a degree-12 K3 surface from three points.*

2.1 Orthogonal Grassmannian

Let V be a 10-dimensional vector space equipped with a non-degenerate quadratic form q . The five-dimensional subspaces of V isotropic with respect to q form a subvariety \mathcal{S} of the Grassmannian $G(5, V)$. It has two components \mathcal{S}^+ and \mathcal{S}^- which are isomorphic to each other. They are called *orthogonal Grassmannians* and are denoted by $OG(5, V)$.

Fix a 5-subspace $W \in OG(5, V)$ and let W^* be its orthogonal complement with respect to q . Then $OG(5, V)$ can be identified scheme-theoretically as the zero locus in

$$\mathbb{P}\left(\mathbb{C} \oplus \bigwedge^2 W \oplus \bigwedge^4 W\right) \simeq \mathbb{P}^{15}$$

of the quadratic form [IM04, § 2]

$$\begin{aligned} \det W \oplus \bigwedge^2 W \oplus W^* &\longrightarrow \bigwedge^4 W \oplus W \\ (x, \Omega, v) &\longmapsto (x(v) + \frac{1}{2}\Omega \wedge \Omega, \Omega(v)). \end{aligned} \tag{2.1}$$

Here we choose an isomorphism $\mathbb{C} \simeq \det W$. This induces an isomorphism $\bigwedge^4 W \simeq W^*$.

Let $\mathbf{x} = (x_0, \dots, x_{15})$ be the homogeneous coordinate for \mathbb{P}^{15} . Then (2.1) can be explicitly written down as ten quadrics:

$$\begin{array}{l|l} x_0x_{11} + x_5x_{10} - x_6x_9 + x_7x_8 & -x_1x_{12} + x_2x_{13} - x_3x_{14} + x_4x_{15} \\ x_0x_{12} + x_2x_{10} - x_3x_9 + x_4x_8 & x_1x_{11} - x_5x_{13} + x_6x_{14} - x_7x_{15} \\ x_0x_{13} + x_1x_{10} - x_3x_7 + x_4x_6 & -x_2x_{11} + x_5x_{12} - x_8x_{14} + x_9x_{15} \\ x_0x_{14} + x_1x_9 - x_2x_7 + x_4x_5 & x_3x_{11} - x_6x_{12} + x_8x_{13} - x_{10}x_{15} \\ x_0x_{15} + x_1x_8 - x_2x_6 + x_3x_5 & -x_4x_{11} + x_7x_{12} - x_9x_{13} + x_{10}x_{14}. \end{array} \tag{2.2}$$

2.2 An explicit example

Mukai [Muk88, § 3] proves that a generic K3 surface of degree 12 appears as a linear section of $OG(5, V)$ and vice versa.

For example, the $\mathbb{P}^7 \subset \mathbb{P}^{15}$ spanned by the rows of the 8×16 matrix

$$\mathbf{H} = \begin{pmatrix} -1 & 3 & 2 & 0 & 2 & -3 & -1 & 0 & 3 & 1 & 0 & 3 & 0 & 2 & 0 & -3 \\ 1 & 0 & -3 & 0 & -2 & 1 & 1 & 0 & -2 & -1 & -1 & -1 & 0 & 4 & 0 & 2 \\ -1 & -3 & -2 & 0 & -3 & 0 & 3 & 2 & -1 & -3 & -1 & 2 & -1 & 2 & 0 & 3 \\ 3 & 0 & 0 & 2 & 2 & 3 & 0 & 1 & 2 & -1 & 0 & 2 & -1 & -2 & 2 & 3 \\ 0 & -1 & 1 & -1 & 0 & 1 & -3 & 3 & 2 & 2 & 1 & 3 & 0 & -3 & 0 & -3 \\ 1 & 0 & 0 & 0 & 0 & -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 3 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

cuts out a degree-12 K3 surface R on $OG(5, V)$. More explicitly, let $\mathbf{z} = (z_0, \dots, z_7)$ be homogeneous coordinates for \mathbb{P}^7 . We define the inclusion $\iota : \mathbb{P}^7 \hookrightarrow \mathbb{P}^{15}$ by

$$\mathbf{x} = \mathbf{z} \cdot \mathbf{H}.$$

Then we get $R = \iota^{-1}(OG(5, V))$.

The last three rows of \mathbf{H} are chosen as solutions of (2.2) so that they form a triple of points $\Pi = \{p_1, p_2, p_3\} \subset R$. With this choice the projection from Π is exactly the map

$$\begin{aligned} \pi : \mathbb{P}^7 &\dashrightarrow \mathbb{P}^4 \\ (z_0, \dots, z_7) &\mapsto (z_0, \dots, z_4) \end{aligned}$$

which takes R to $S = \pi(R)$.

We manipulate this example in a computer algebra system¹ over the finite field \mathbb{F}_7 . We compute that S is singular along three transverse double points $\{a_1, a_2, a_3\}$ and is the base locus of a Cremona transformation

$$\bar{f} : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4.$$

Moreover, the base locus of the inverse $(\bar{f})^{-1}$ is again a surface \bar{T} singular along three transverse double points $\{b_1, b_2, b_3\}$. The matrix \mathbf{H} is chosen such that the preimage of $\{a_1, a_2, a_3\}$ on R and the preimage of $\{b_1, b_2, b_3\}$ on the normalization of \bar{T} are \mathbb{F}_7 -rational points. This is the smallest field where our computer could quickly find such an \mathbf{H} .

2.3 Proof of Theorem 2.1

We prove Theorem 2.1 for our example first.

We confirm the following properties by computer over \mathbb{F}_7 .

- (1) S is singular along three points. The preimage of each singular point on R has two points outside Π . So they are transverse double points.
- (2) The ideal of S is generated by five quartics $\bar{f}_0, \dots, \bar{f}_4$.

The double-point formula [Ful98, Theorem 9.3] indicates that the three transverse double points of (1) exist over characteristic zero. Indeed, let $\epsilon : \Sigma \rightarrow S$ be the normalization. Then Σ is isomorphic to R blown up at three points. The *double-point class* $\mathbb{D}(\epsilon) \in \text{CH}_0(\Sigma)$ is given by the formula

$$\begin{aligned} \mathbb{D}(\epsilon) &= \epsilon^* \epsilon_* [\Sigma] - [\epsilon^* c(\mathbb{P}^4) \cdot c(\Sigma)^{-1}]_0 \\ &= \langle S, S \rangle_{\mathbb{P}^4} - (\epsilon^* c_2(\mathbb{P}^4) - \epsilon^* c_1(\mathbb{P}^4) \cdot c_1(\Sigma) - c_1(\Sigma)^2 + c_2(\Sigma)). \end{aligned}$$

It is easy to verify that $\mathbb{D}(\epsilon) = 6$. The quantity $\frac{1}{2}\mathbb{D}(\epsilon) = 3$ counts the number of singularities on S with multiplicity if the singular locus is a finite set. Therefore (1) implies that the singular locus of S consists of three transverse double points. This proves Theorem 2.1(1).

The five quartics $\bar{f}_0, \dots, \bar{f}_4$ lift to a basis f_0, \dots, f_4 for the ideal of S over characteristic zero. In particular, Theorem 2.1(2) holds. The forms f_0, \dots, f_4 define a rational map

$$f := (f_0, \dots, f_4) : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$$

which reduces to

$$\bar{f} := (\bar{f}_0, \dots, \bar{f}_4) : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$$

over \mathbb{F}_7 . The degree of f is computed by the self-intersection M^4 , which can be expanded as the right-hand side of equation (5.2). It is easy to check that our example satisfies

$$(n, m, d, \delta) = (4, 1, 9, 3), \quad K_\Sigma C = 3, \quad \text{and} \quad c_2(\Sigma) = 27.$$

Inserting these data into (5.2), we get $M^4 = 1$, that is, the map f is birational. Thus Theorem 2.1(3) holds.

¹The main program we use in this work is SINGULAR [DGPS15].


The inverse $(\bar{f})^{-1}$ can be calculated by computer. It consists of five quartics also and the base locus is a surface \bar{T} singular along three points. These are transverse double points since each point has two preimage points on the normalization. By the same reasons as above, the base locus of f^{-1} is again a surface cut by five quartics and singular along three transverse double points. Then Theorem 2.1(4) follows from Theorem 5.1.

Next, we prove Theorem 2.1 in the generic case.

It is clear that (1), (2), and (3) of the theorem are open conditions, so they hold for a generic example. As a consequence of Theorem 5.1, property (4) holds once $\text{Bs}(f^{-1})$ is a surface cut out by five quartics and singular along three transverse double points. These are open conditions again, so Theorem 2.1 holds for a generic example.

2.4 Some geometry of the construction

Let f be a Cremona transformation of Theorem 2.1. It has a resolution



Let S_L and S_M be the base locus of f and its inverse f^{-1} , respectively. Then π_1 is the blow-up along S_L . Recall that X coincides with the graph of f as well as f^{-1} . Therefore, π_2 is the blow-up along S_M .

This example has $d = 9$, $K_\Sigma C = 3$, $c_2(\Sigma) = 27$, $\delta = 3$, and

$$M = 4L - E - 2\sum_{i=1}^3 E'_i.$$

Evaluating Lemmas 1.1 and 1.2 with this data yields the following corollary.

COROLLARY 2.2. *We have*

- (1) $LE'_i = 0$, $E^3 E'_i = -4$, $E^2 E'^2_i = 2$, $EE'^3_i = 0$, $E'^4_i = -1$;
- (2) $L^3 E = 0$, $L^2 E^2 = -9$, $LE^3 = -48$, $E^4 = -159$.

Thus consequently,

- (3) $L^3 M = 4$, $L^2 M^2 = 7$, $LM^3 = 4$, $M^4 = 1$.

Let X_0, \dots, X_4 be the homogeneous coordinates for \mathbb{P}^4 . The Cremona transformation f is ramified along the locus Θ where the Jacobian matrix

$$Df = \left(\frac{\partial f_i}{\partial X_j} \right)_{5 \times 5}$$

is degenerate. So Θ is a degree-15 hypersurface in \mathbb{P}^4 defined by

$$\det(Df) = 0.$$

This locus is called the *P-locus*, which is classically defined as the image of the exceptional divisor of the blow-up π_2 [Dol12, § 7.1.4]. In particular, Θ is irreducible. It also follows that Θ is the locus contracted by f and its image is the base locus S_M .

PROPOSITION 2.3. *The locus $\Theta \subset \mathbb{P}^4$ contracted by f is an irreducible hypersurface of degree 15. It has multiplicity four along S_L . Moreover, it equals the union of all of the 4-secant lines to S_L . The analogous statement holds for the inverse f^{-1} by symmetry.*

Proof. Let m be the multiplicity of Θ along S_L . Then the divisor class of its pullback to X equals

$$\Theta_X = 15L - mE_X.$$

Here we use E_X to denote the exceptional divisor of the blow-up π_1 . Because Θ is contracted onto a surface, we have

$$0 = M^3\Theta_X = M^3(15L - mE_X) = 60 - mM^3E_X.$$

By definition, E_X is mapped onto the P -locus of the inverse map f^{-1} . In particular, it is again a degree-15 hypersurfaces in \mathbb{P}^4 by symmetry. So $M^3E_X = 15$, which implies $m = 4$.

Let F_X be the exceptional locus of the blow-up π_2 . We have $L = 4M - F_X$ by symmetry, hence

$$F_X = 4M - L.$$

(Note that this equals $15L - 4E_X = \Theta_X$.) The fiber of the map $F_X \rightarrow S_M$ over a smooth point is represented by the class

$$l_X = \frac{F_X M^2}{\deg S_M} = \frac{1}{9}F_X M^2.$$

The image $l = \pi_1(l_X)$ is a rational curve of degree

$$L \cdot l_X = \frac{1}{9}LF_X M^2 = \frac{1}{9}L(4M - L)M^2 = \frac{1}{9}(16 - 7) = 1.$$

The intersection number between l and S_L can be computed by

$$\begin{aligned} E_X \cdot l_X &= \frac{1}{9}E_X F_X M^2 = \frac{1}{9}(4L - M)(4M - L)M^2 \\ &= \frac{1}{9}(64 - 28 - 4 + 4) = 4. \end{aligned}$$

Hence the fibers of $F_X \rightarrow S_M$ away from the double points is mapped by π_1 to 4-secant lines to S_L . In other words, S_L admits a family of 4-secant lines parametrized by the smooth locus of S_M .

Conversely, every 4-secant line l to S_L satisfies

$$l \cdot M = l \cdot (4L - E_X) = 4 - 4 = 0.$$

So l is contracted to a point by f . Hence the union of the 4-secant lines to S_L forms a 3-fold contained in Θ and thus coincides with Θ . □

3. Derived equivalences of K3 surfaces

Let us retain the notation of § 2.4. By Theorem 2.1, there exists two degree-12 K3 surfaces R_L and R_M projected onto the base loci S_L and S_M , respectively. This section is devoted to the following results.

THEOREM 3.1. *The two K3 surfaces R_L and R_M are derived equivalent. They are non-isomorphic if they have Picard number one.*

COROLLARY 3.2. *There is a birational map $\sigma : R_L^{[3]} \dashrightarrow R_M^{[3]}$ between the Hilbert schemes of length three subschemes.*

3.1 Derived equivalences and general strategy

Let R and \hat{R} denote K3 surfaces and $T(R)$ and $T(\hat{R})$ the corresponding transcendental lattices. Recall that R and \hat{R} are derived equivalent if and only if $T(R)$ and $T(\hat{R})$ are Hodge isometric [Orl97]. Suppose that R has Picard rank one and degree $2n$. Let $\tau(n)$ be the number of prime factors of n . Then the number of isomorphism classes of K3 surfaces derived equivalent to R is equal to $2^{\tau(n)-1}$ [HLOY03]. Thus a general degree-12 K3 surface admits a unique such partner.

Our general approach is to prove that $T(R_L)$ is isometric to $T(R_M)$ by showing that both of them can be identified as the transcendental sublattice of $H^4(X, \mathbb{Z})$. Then we show that the induced isomorphism on the discriminant groups is non-trivial, which implies that R_L and R_M are not isomorphic to each other.

3.2 The middle cohomology of X

Retain the notation of § 1. Let H_L be the polarization of R_L . Let $F_1, F_2,$ and F_3 be the exceptional curves from the projection $R_L \dashrightarrow S_L$. We consider $H_L, F_1, F_2,$ and F_3 as curves on S_L . Their strict transforms $\tilde{H}_L, \tilde{F}_1, \tilde{F}_2, \tilde{F}_3$ on X together with L^2 and the quadrics Q_1, Q_2, Q_3 form a rank-eight sublattice $A_L(X) \subset H^4(X, \mathbb{Z})$. We have

$$\tilde{H}_L^2 = -H_L^2 = -12, \quad \tilde{F}_i^2 = -F_i^2 = 1, \quad \text{and} \quad Q_i^2 = -E_i'^4 = 1,$$

where $i = 1, 2, 3$. These classes are mutually disjoint, so the intersection matrix for $A_L(X)$ is

$$\begin{array}{c|cccc} & L^2 & \tilde{H}_L & \tilde{F}_{1,2,3} & Q_{1,2,3} \\ \hline L^2 & 1 & & & \\ \tilde{H}_L & & -12 & & \\ \tilde{F}_{1,2,3} & & & I_{3 \times 3} & \\ Q_{1,2,3} & & & & I_{3 \times 3} \end{array} \tag{3.1}$$

where $I_{3 \times 3}$ is the identity matrix of rank 3.

LEMMA 3.3. *There is a decomposition*

$$H^4(X, \mathbb{Z}) \simeq H^4(X, \mathbb{Z})_{\text{alg}} \oplus_{\perp} T(R_L)(-1),$$

where $H^4(X, \mathbb{Z})_{\text{alg}}$ is the sublattice spanned by algebraic classes. We have

$$H^4(X, \mathbb{Z})_{\text{alg}} = A_L(X)$$

when R_L has Picard number one. Here we use $\Lambda(-1)$ to denote a lattice Λ equipped with the negative of its original product.

Proof. We apply the blow-up formula for cohomology to the composition $P' \rightarrow P \rightarrow \mathbb{P}^4$ and the map $P' \rightarrow X$ to obtain two decompositions for $H^4(P', \mathbb{Z})$. Then we compare them to get our result.

Let $S'_L \subset P$ be the strict transform of S_L . Recall that S'_L is isomorphic to R_L blown up at $3+6=9$ points, where three are from the projection $R_L \dashrightarrow S_L$ while six are from the resolution $S'_L \rightarrow S_L$. Thus we have

$$H^2(S'_L, \mathbb{Z}) \simeq \langle F_i, Q'_i, Q''_i \rangle_{i=1,2,3} \oplus H^2(R_L, \mathbb{Z}).$$

Let \tilde{Q}'_i and \tilde{Q}''_i be the strict transforms of Q'_i and Q''_i on P' . Since $P' \rightarrow P$ is the blow-up along S'_L , we have

$$\begin{aligned} H^4(P', \mathbb{Z}) &\simeq H^4(P, \mathbb{Z}) \oplus H^2(S'_L, \mathbb{Z})(-1) \\ &\simeq \langle L^2, E_i'^2, \tilde{F}_i, \tilde{Q}'_i, \tilde{Q}''_i \rangle_{i=1,2,3} \oplus H^2(R_L, \mathbb{Z})(-1). \end{aligned} \tag{3.2}$$

For every i , we have

$$\tilde{Q}_i'^2 = -Q_i'^2 = 1, \quad \tilde{Q}_i''^2 = -Q_i''^2 = 1, \quad E_i'^2 \tilde{Q}'_i = E_i''^2 \tilde{Q}''_i = 0,$$

and $E_i'^4 = -1$. With these it is straightforward to prove the isometry

$$\langle E_i'^2, \tilde{Q}'_i, \tilde{Q}''_i \rangle \simeq \langle E_i'^2 + \tilde{Q}'_i, E_i'^2 + \tilde{Q}''_i, E_i'^2 + \tilde{Q}'_i + \tilde{Q}''_i \rangle,$$

whence (3.2) equals

$$H^4(P', \mathbb{Z}) \simeq \langle E_i'^2 + \tilde{Q}'_i, E_i''^2 + \tilde{Q}''_i, E_i'^2 + \tilde{Q}'_i + \tilde{Q}''_i, L^2, \tilde{F}_i \rangle_{i=1,2,3} \oplus H^2(R_L, \mathbb{Z})(-1). \tag{3.3}$$

By the description of the map $E_i \dashrightarrow Q_i$, the two fiber classes on $Q_i \simeq \mathbb{P}^1 \times \mathbb{P}^1$ pull back to hyperplanes in E_i containing either Q'_i or Q''_i , which correspond to the classes $-E_i'^2 - \tilde{Q}'_i$ or $-E_i''^2 - \tilde{Q}''_i$ on P' , respectively. The map $P' \rightarrow X$ is the blow-up along Q_i , $i = 1, 2, 3$, so

$$H^4(P', \mathbb{Z}) \simeq \langle E_i'^2 + \tilde{Q}'_i, E_i''^2 + \tilde{Q}''_i \rangle_{i=1,2,3} \oplus H^4(X, \mathbb{Z}). \tag{3.4}$$

Combining (3.3) and (3.4), we get

$$\begin{aligned} H^4(X, \mathbb{Z}) &\simeq \langle E_i'^2 + \tilde{Q}'_i, E_i''^2 + \tilde{Q}''_i \rangle_{i=1,2,3}^\perp \\ &\simeq \langle E_i'^2 + \tilde{Q}'_i + \tilde{Q}''_i, L^2, \tilde{F}_i \rangle_{i=1,2,3} \oplus H^2(R_L, \mathbb{Z})(-1). \end{aligned}$$

Both Q_i and $E_i'^2 + \tilde{Q}'_i + \tilde{Q}''_i$ are orthogonal to L^2, \tilde{F}_i and $H^2(R_L, \mathbb{Z})$, and $Q_i^2 = (E_i'^2 + \tilde{Q}'_i + \tilde{Q}''_i)^2 = 1$, so $Q_i = \pm(E_i'^2 + \tilde{Q}'_i + \tilde{Q}''_i)$. Therefore

$$\begin{aligned} H^4(X, \mathbb{Z}) &\simeq \langle Q_i, L^2, \tilde{F}_i \rangle_{i=1,2,3} \oplus H^2(R_L, \mathbb{Z})(-1) \\ &\simeq \langle Q_i, L^2, \tilde{F}_i \rangle_{i=1,2,3} \oplus NS(R_L)(-1) \oplus_\perp T(R_L)(-1) \\ &\simeq H^4(X, \mathbb{Z})_{\text{alg}} \oplus_\perp T(R_L)(-1), \end{aligned}$$

where $NS(R_L)$ is the Néron–Severi lattice of R_L .

When R_L has Picard number one, we have $NS(R_L)(-1) \simeq \langle \tilde{H}_L \rangle$. In this case

$$H^4(X, \mathbb{Z})_{\text{alg}} \simeq \langle Q_i, L^2, \tilde{F}_i, \tilde{H}_L \rangle_{i=1,2,3} = A_L(X). \quad \square$$

Lemma 3.3 also proves the decomposition

$$H^4(X, \mathbb{Z}) \simeq H^4(X, \mathbb{Z})_{\text{alg}} \oplus_\perp T(R_M)(-1)$$

from the side of f^{-1} . So there is an isometry

$$T(R_L) \simeq H^4(X, \mathbb{Z})_{\text{alg}}^\perp(-1) \simeq T(R_M)$$

which allows us to conclude the following proposition.

PROPOSITION 3.4. R_L and R_M are derived equivalent.

3.3 The discriminant groups

For an arbitrary lattice Λ with dual lattice $\Lambda^* := \text{Hom}(\Lambda, \mathbb{Z})$, we denote by $d\Lambda := \Lambda^*/\Lambda$ its discriminant group.

Let $A_M(X)$ be the lattice constructed in the same way as $A_L(X)$ from the side of f^{-1} . Assume R_L and R_M have Picard number one. Then Lemma 3.3 implies that there is an isometry

$$\varphi : A_M(X) \oplus_{\perp} T(R_M)(-1) \xrightarrow{\sim} A_L(X) \oplus_{\perp} T(R_L)(-1)$$

such that $\varphi = \varphi_A \oplus \varphi_T$ with respect to the decompositions. It induces the commutative diagram

$$\begin{array}{ccc} dA_M(X) & \xrightarrow[\sim]{\varphi_{A^*}} & dA_L(X) \\ \sim \downarrow & & \sim \downarrow \\ dT(R_M) & \xrightarrow[\sim]{\varphi_{T^*}} & dT(R_L). \end{array}$$

These groups are all isomorphic to $\mathbb{Z}/12\mathbb{Z}$. From the intersection matrix (3.1) we know that $dA_L(X)$ is generated by $-\tilde{H}_L/12$. Similarly, $dA_M(X)$ is generated by $-\tilde{H}_M/12$, where H_M is the polarization of R_M and \tilde{H}_M is the strict transform on X .

LEMMA 3.5. *We have the following equations in $H^4(X, \mathbb{Z})$:*

- (1) $M^2 = 7L^2 - 3\tilde{H}_L + 4(\tilde{F}_1 + \tilde{F}_2 + \tilde{F}_3) + 2(Q_1 + Q_2 + Q_3)$;
- (2) $\tilde{H}_M = 36L^2 - 17\tilde{H}_L + 24(\tilde{F}_1 + \tilde{F}_2 + \tilde{F}_3) + 12(Q_1 + Q_2 + Q_3)$.

Proof. The following computation is based on Corollary 2.2. Assume that

$$M^2 = aL^2 + b\tilde{H}_L + f_1\tilde{F}_1 + f_2\tilde{F}_2 + f_3\tilde{F}_3 + g_1Q_1 + g_2Q_2 + g_3Q_3.$$

Then $a = L^2M^2 = 7$. For $i = 1, 2, 3$, we have

$$\begin{aligned} g_i &= M^2Q_i = -M^2E_i'^2 \\ &= -(4L - E - 2\sum_j E_j')^2 E_i'^2 = -(-E - 2\sum_j E_j')^2 E_i'^2 \\ &= -E^2 E_i'^2 - 4EE_i'^3 - 4E_i'^4 = -2 - 0 + 4 = 2. \end{aligned}$$

Let \tilde{C} be the strict transform of the sectional curve C on P' . Note that

$$LM = 4L^2 - LE = 4L^2 - \tilde{C} = 4L^2 - \tilde{H}_L + \sum_i \tilde{F}_i,$$

so we find

$$\begin{aligned} 4 &= LM^3 = (LM)M^2 \\ &= (4L^2 - \tilde{H}_L + \sum_i \tilde{F}_i)(7L^2 + b\tilde{H}_L + \sum_i f_j \tilde{F}_j + 2\sum_k Q_k) \\ &= 28 + 12b + f_1 + f_2 + f_3, \end{aligned}$$

and thus

$$f_1 + f_2 + f_3 = -12b - 24. \tag{3.5}$$

We also have

$$\begin{aligned} 1 &= M^4 = (7L^2 + b\tilde{H}_L + \sum_i f_j \tilde{F}_j + 2\sum_k Q_k)^2 \\ &= 49 - 12b^2 + f_1^2 + f_2^2 + f_3^2 + 12, \end{aligned}$$

which is equivalent to

$$f_1^2 + f_2^2 + f_3^2 = 12b^2 - 60. \tag{3.6}$$

By the Cauchy–Schwarz inequality,

$$(f_1 + f_2 + f_3)^2 = ((1, 1, 1) \cdot (f_1, f_2, f_3))^2 \leq (1, 1, 1)^2 (f_1, f_2, f_3)^2 = 3(f_1^2 + f_2^2 + f_3^2). \tag{3.7}$$

Applying (3.5) and (3.6), we get $(-12b - 24)^2 \leq 3(12b^2 - 60)$, that is,

$$3b^2 + 16b + 21 = (3b + 7)(b + 3) \leq 0.$$

The only integer solution is $b = -3$. Because (3.7) becomes an equality in this case, we have $(f_1, f_2, f_3) = f(1, 1, 1)$ for some integer f . We obtain $f = 4$ by setting $b = -3$ in (3.5). As a result, we find

$$M^2 = 7L^2 - 3\tilde{H}_L + 4(\tilde{F}_1 + \tilde{F}_2 + \tilde{F}_3) + 2(Q_1 + Q_2 + Q_3).$$

Next, assume that

$$\tilde{H}_M = aL^2 + b\tilde{H}_L + f_1\tilde{F}_1 + f_2\tilde{F}_2 + f_3\tilde{F}_3 + g_1Q_1 + g_2Q_2 + g_3Q_3.$$

By symmetry, $\tilde{H}_L L^n M^{2-n} = \tilde{H}_M M^n L^{2-n}$ for $n = 0, 1, 2$. In particular,

$$a = \tilde{H}_M L^2 = \tilde{H}_L M^2 = -3\tilde{H}_L^2 = 36.$$

We have

$$\begin{aligned} 12 &= \tilde{H}_L(4L^2 - \tilde{H}_L + \Sigma_i \tilde{F}_i) = \tilde{H}_L(LM) = \tilde{H}_M(ML) \\ &= (36L^2 + b\tilde{H}_L + \Sigma_i f_i \tilde{F}_i + \Sigma_j g_j Q_j)(4L^2 - \tilde{H}_L + \Sigma_i \tilde{F}_i) \\ &= 144 + 12b + f_1 + f_2 + f_3. \end{aligned}$$

Rearrange to obtain

$$f_1 + f_2 + f_3 = -12b - 132. \tag{3.8}$$

Applying the symmetry again, we get

$$\begin{aligned} 0 &= \tilde{H}_L L^2 = \tilde{H}_M M^2 \\ &= (36L^2 + b\tilde{H}_L + \Sigma_i f_i \tilde{F}_i + \Sigma_j g_j Q_j)(7L^2 - 3\tilde{H}_L + 4\Sigma_i \tilde{F}_i + 2\Sigma_j Q_j) \\ &= 252 + 36b + 4(f_1 + f_2 + f_3) + 2(g_1 + g_2 + g_3), \end{aligned}$$

whence

$$2(f_1 + f_2 + f_3) + (g_1 + g_2 + g_3) = -18b - 126,$$

and combining with (3.8) gives

$$g_1 + g_2 + g_3 = 6b + 138. \tag{3.9}$$

We also have

$$\begin{aligned} -12 &= \tilde{H}_L^2 = \tilde{H}_M^2 = (36L^2 + b\tilde{H}_L + \Sigma_i f_i \tilde{F}_i + \Sigma_j g_j Q_j)^2 \\ &= 1296 - 12b^2 + f_1^2 + f_2^2 + f_3^2 + g_1^2 + g_2^2 + g_3^2, \end{aligned}$$

from which we obtain

$$f_1^2 + f_2^2 + f_3^2 + g_1^2 + g_2^2 + g_3^2 = 12b^2 - 1308. \tag{3.10}$$

By the Cauchy–Schwarz inequality,

$$\begin{aligned} (-12b - 132)^2 &= (f_1 + f_2 + f_3)^2 = ((1, 1, 1) \cdot (f_1, f_2, f_3))^2 \\ &\leq (1, 1, 1)^2 (f_1, f_2, f_3)^2 = 3(f_1^2 + f_2^2 + f_3^2) \end{aligned} \tag{3.11}$$

and

$$\begin{aligned} (6b + 138)^2 &= (g_1 + g_2 + g_3)^2 = ((1, 1, 1) \cdot (g_1, g_2, g_3))^2 \\ &\leq (1, 1, 1)^2 (g_1, g_2, g_3)^2 = 3(g_1^2 + g_2^2 + g_3^2). \end{aligned} \tag{3.12}$$

Add the two inequalities and then apply (3.10) to get

$$(-12b - 132)^2 + (6b + 138)^2 \leq 3(f_1^2 + f_2^2 + f_3^2 + g_1^2 + g_2^2 + g_3^2) = 3(12b^2 - 1308), \tag{3.13}$$

which can be arranged as

$$2b^2 + 67b + 561 = (2b + 33)(b + 17) \leq 0.$$

The only integer solution is $b = -17$ which makes (3.13) an equality. This forces (3.11) and (3.12) to be equalities also. Therefore $(f_1, f_2, f_3) = f(1, 1, 1)$ and $(g_1, g_2, g_3) = g(1, 1, 1)$ for some integers f and g . We get $f = 24$ from (3.8) and $g = 12$ from (3.9). As a consequence,

$$\tilde{H}_M = 36L^2 - 17\tilde{H}_L + 24(\tilde{F}_1 + \tilde{F}_2 + \tilde{F}_3) + 12(Q_1 + Q_2 + Q_3). \quad \square$$

PROPOSITION 3.6. *The isomorphism $\varphi_{A_*} : dA_M(X) \xrightarrow{\sim} dA_L(X)$ equals multiplication by 7 on $\mathbb{Z}/12\mathbb{Z}$.*

Proof. Recall that φ_A acts as the identity map on $H^4(X, \mathbb{Z})_{\text{alg}}$, thus $\varphi_A(\tilde{H}_M) = \tilde{H}_M$. By Lemma 3.5 we have

$$\varphi_A(\tilde{H}_M) = 36L^2 - 17\tilde{H}_L + 24(\tilde{F}_1 + \tilde{F}_2 + \tilde{F}_3) + 12(Q_1 + Q_2 + Q_3)$$

as a map from $A_M(X)$ to $A_L(X)$. Therefore

$$\begin{aligned} \varphi_{A_*}(-\frac{1}{12}\tilde{H}_M) &= -3L^2 + \frac{17}{12}\tilde{H}_L - 2(\tilde{F}_1 + \tilde{F}_2 + \tilde{F}_3) - (Q_1 + Q_2 + Q_3) \\ &= -17 \cdot (-\frac{1}{12}\tilde{H}_L) \pmod{A_L(X)} \\ &= 7 \cdot (-\frac{1}{12}\tilde{H}_L) \pmod{A_L(X)}. \end{aligned} \quad \square$$

Remark. By the symmetry of the Cremona transformation f , the rank-eight lattice $H^4(X, \mathbb{Z})_{\text{alg}}$ is also spanned by the classes

$$\{M^2, \tilde{H}_M, \tilde{G}_1, \tilde{G}_2, \tilde{G}_3, K_1, K_2, K_3\}$$

constructed in a similar way from the right-hand side. Here $\tilde{G}_1, \tilde{G}_2, \tilde{G}_3$ are from the exceptional curves and K_1, K_2, K_3 are from the transverse double points. The full transformation between the two set of bases is

$$\begin{pmatrix} M^2 \\ \tilde{H}_M \\ \tilde{G}_1 \\ \tilde{G}_2 \\ \tilde{G}_3 \\ K_1 \\ K_2 \\ K_3 \end{pmatrix} = \begin{pmatrix} 7 & -3 & 4 & 4 & 4 & 2 & 2 & 2 \\ 36 & -17 & 24 & 24 & 24 & 12 & 12 & 12 \\ 4 & -2 & 3 & 3 & 3 & 2 & 1 & 1 \\ 4 & -2 & 3 & 3 & 3 & 1 & 2 & 1 \\ 4 & -2 & 3 & 3 & 3 & 1 & 1 & 2 \\ 2 & -1 & 2 & 1 & 1 & 1 & 1 & 1 \\ 2 & -1 & 1 & 2 & 1 & 1 & 1 & 1 \\ 2 & -1 & 1 & 1 & 2 & 1 & 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} L^2 \\ \tilde{H}_L \\ \tilde{F}_1 \\ \tilde{F}_2 \\ \tilde{F}_3 \\ Q_1 \\ Q_2 \\ Q_3 \end{pmatrix}.$$

This expression is unique up to the ordering of the exceptional curves and the transverse double points on each side. The top two rows are computed by Lemma 3.5. The other rows can be computed in a similar way.

3.4 Proofs of Theorem 3.1 and its Corollary

We first prove the theorem.

The derived equivalence follows from Proposition 3.4. Note that this implies that the Picard numbers of R_L and R_M are the same.

Assume R_L and R_M have Picard number one. Suppose they are isomorphic. Then there is an isometry

$$\theta : T(R_L) \xrightarrow{\sim} T(R_M)$$

which induces the isomorphism

$$\begin{aligned} \theta_* : dT(R_L) &\xrightarrow{\sim} dT(R_M) \\ -\frac{\tilde{H}_L}{12} &\mapsto -\frac{\tilde{H}_M}{12} \end{aligned}$$

under the identifications $dT(R_L) \simeq dA_L(X)$ and $dT(R_M) \simeq dA_M(X)$.

By Proposition 3.6, the composition $\varphi_A \circ \theta$ is an automorphism on $T(R_L)$ acting as multiplication by 7 on $dT(R_L)$. This contradicts the fact that the only automorphism on $T(R_L)$ is the identity [Ogu02]. Hence R_L and R_M cannot be isomorphic to each other.

Next we prove the corollary.

The corollary is trivial if R_L and R_M are isomorphic, so we assume that they are non-isomorphic.

Given a generic triple of points $\Pi_L \in R_L^{[3]}$, we determine a degree-12 K3 surface R_M and a triple of points $\Pi_M \in R_M^{[3]}$ through the following steps.

- (1) Project R_L from Π_L to obtain $S_L \subset \mathbb{P}^4$, whose ideal defines a Cremona transformation $f : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$.
- (2) The base locus of f^{-1} is a surface S_M singular along three transverse double points. Normalize S_M to get Σ_M .
- (3) Σ_M is the blow-up of a degree-12 K3 surface R_M along three points. The three exceptional curves on Σ_M are contracted to $\Pi_M \in R_M^{[3]}$.

Recall that two derived equivalent K3 surfaces of degree 12 uniquely determine each other up to isomorphism. So R_M is independent of the choice of $\Pi_L \in R_L^{[3]}$ by Theorem 3.1. Hence there is a rational map

$$\begin{aligned} \sigma : R_L^{[3]} &\dashrightarrow R_M^{[3]} \\ \Pi_L &\mapsto \Pi_M. \end{aligned}$$

It is birational because Π_L is uniquely determined by Π_M through the same process as above.

3.5 Connections between our construction and other approaches

The derived equivalence and geometric connections between the degree-12 K3 surfaces (R_L, H_L) and (R_M, H_M) admit several interpretations.

3.5.1 *Mukai lattices.* For a K3 surface R , the Mukai lattice

$$\tilde{H}(R, \mathbb{Z}) := H^0(R, \mathbb{Z}) \oplus H^2(R, \mathbb{Z}) \oplus H^4(R, \mathbb{Z}),$$

equipped with a weight-two Hodge structure, that is, the standard Hodge structure on the middle summand and the outer summands taken as $(1, 1)$ classes. This is polarized by

$$(r_1, D_1, s_1) \cdot (r_2, D_2, s_2) = D_1 \cdot D_2 - r_1 \cdot s_2 - r_2 \cdot s_1.$$

Each coherent sheaf E yields a Mukai vector

$$v(E) = (r(E), c_1(E), s(E)),$$

where $r(E)$ is the rank and $r(E) + s(E) = \chi(E)$. Mukai [Muk87] has shown that the second cohomology of a moduli space $M_v(R)$ may be expressed as

$$H^2(M_v(R), \mathbb{Z}) = \begin{cases} v^\perp & \text{if } v \cdot v \geq 2, \\ v^\perp / \mathbb{Z}v & \text{if } v \cdot v = 0, \end{cases}$$

provided v is primitive and satisfies certain technical conditions. A derived equivalence between R and \hat{R} induces an isomorphism of Hodge structures

$$\Phi : \tilde{H}(R, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\hat{R}, \mathbb{Z})$$

which may be chosen so that $\hat{R} = M_v(R)$ with $\Phi(v) = (0, 0, 1)$.

We return to our degree-12 K3 surfaces R_M and R_L . We may interpret R_M as a moduli space of vector bundles on R_L and vice versa [Muk99]. Let $M_{(2, H_L, 3)}(R_L)$ denote the moduli space of rank-two stable bundles E with $c_1(E) = H_L$ and $\chi(R_L, E) = 5$, which is isomorphic to R_M . The universal bundle $\mathcal{E} \rightarrow R_L \times R_M$ induces a Hodge isometry

$$\Phi : \tilde{H}(R_L, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(R_M, \mathbb{Z})$$

described above. We have

$$\Phi(2, H_L, 3) = (0, 0, 1), \quad \Phi(0, 0, 1) = (2, H_M, 3),$$

and Φ restricts to the isogeny on transcendental cohomology mentioned in §3.1. It follows formally that

$$\Phi(1, 0, -2) = (-1, 0, 2),$$

thus after a shift the Mukai vector of ideal sheaves of length-three subschemes of R_L goes to the Mukai vector of length-three subschemes of R_M . We obtain an isomorphism

$$H^2(R_L^{[3]}, \mathbb{Z}) \simeq H^2(R_M^{[3]}, \mathbb{Z})$$

of Hodge structures arising from Mukai lattices. Thus the Torelli theorem [Mar11, Corollary 9.9] yields a birational equivalence

$$R_L^{[3]} \dashrightarrow R_M^{[3]}.$$

Corollary 3.2 is quite natural from this perspective.

Remark. We also have $\Phi(1, 0, -1) = (1, H_M, 5)$. Elements of $M_{(1, H_M, 5)}(R_M)$ may be interpreted as $I_Z(H_M)$ where $Z \subset R_M$ has length two. Similar reasoning gives

$$R_L^{[2]} \dashrightarrow R_M^{[2]}.$$

3.5.2 *Homological projective duality.* Mukai [Muk99, Example 1.3] proposed an interpretation of the derived equivalence via linear algebra, which was explored in detail by Iliev and Markushevich [IM04]. The key observation is that the components \mathcal{S}^+ and \mathcal{S}^- described in § 2.1 are naturally embedded in dual projective spaces

$$\mathcal{S}^+ \subset \mathbb{P}^{15}, \quad \mathcal{S}^- \subset \check{\mathbb{P}}^{15}.$$

Codimension- r subspaces $\mathbb{P} \subset \mathbb{P}^{15}$ correspond to codimension- $(16-r)$ subspaces $\mathbb{P}^\perp \subset \check{\mathbb{P}}^{15}$. When $r = 8$, the K3 surfaces $R = \mathcal{S}^+ \cap \mathbb{P}$ and $\check{R} = \mathcal{S}^- \cap \mathbb{P}^\perp$ are dual. These are derived equivalent and generally non-isomorphic. Kuznetsov [Kuz06, § 6.2] has interpreted this derived equivalence via homological projective duality.

It would be interesting to construct the Cremona transformation through this mechanism, by introducing the data of the three points on the K3 surface into the duality construction.

4. Zero-divisors in the Grothendieck ring

Let $K_0(\text{Var}/\mathbb{C})$ denote the Grothendieck ring of complex algebraic varieties. It is the abelian group generated by isomorphism classes of complex algebraic varieties subject to the relation

$$[Z] = [U] + [Z - U],$$

where U is an open subvariety of Z . The multiplication is induced by the Cartesian product

$$[X][Y] = [X \times Y],$$

which is associative and commutative with unit $1 = [\text{Spec } \mathbb{C}]$. More generally, if $Z \rightarrow X$ is a Zariski locally trivial bundle with fibers isomorphic to Y , by stratifying the base it is easy to prove that

$$[X][Y] = [Z].$$

Let $\mathbb{L} = [\mathbb{A}^1]$ be the class of the affine line in $K_0(\text{Var}/\mathbb{C})$. Consider a pair of non-isomorphic smooth projective varieties X and Y which are derived equivalent. It is interesting to know if there exists $k \geq 0$ satisfying

$$([X] - [Y])\mathbb{L}^k = 0 \tag{4.1}$$

and what the minimal k is if it exists [KS17].

When X is a generic K3 surface of degree 12, Ito *et al.* [IMOU16b] prove that there exists Y non-trivially derived equivalent to X such that (4.1) holds for $k = 3$. Actually, this can be improved to $k = 1$ straightforwardly from the point of view of the Cremona transformation.

THEOREM 4.1. *Let R_L and R_M be a generic pair of K3 surfaces associated with our Cremona transformation. Then we have*

$$([R_L] - [R_M])\mathbb{L} = 0.$$

in $K_0(\text{Var}/\mathbb{C})$. The relation is minimal in the sense that $[R_L] - [R_M] \neq 0$.

Proof. Recall that Σ_L is the normalization of S_L as well as the blow-up of R_L at three points. Hence we have

$$[S_L] = [\Sigma_L] - 3 = [R_L] + 3\mathbb{L} - 3.$$

From the blow-up $\pi_1 : X \rightarrow \mathbb{P}^4$ we obtain

$$\begin{aligned}
 [X] &= ([\mathbb{P}^4] - [S_L]) + [\pi_1^{-1}(S_L)] \\
 &= ([\mathbb{P}^4] - [S_L]) + (([S_L] - 3)[\mathbb{P}^1] + [Q_1] + [Q_2] + [Q_3]) \\
 &= ([\mathbb{P}^4] - [S_L]) + (([S_L] - 3)[\mathbb{P}^1] + 3[\mathbb{P}^1]^2) \\
 &= [\mathbb{P}^4] + 3[\mathbb{P}^1]([\mathbb{P}^1] - 1) + [S_L]([\mathbb{P}^1] - 1) \\
 &= [\mathbb{P}^4] + 3[\mathbb{P}^1]\mathbb{L} + [S_L]\mathbb{L} \\
 &= [\mathbb{P}^4] + 3[\mathbb{P}^1]\mathbb{L} + [R_L]\mathbb{L} + 3\mathbb{L}^2 - 3\mathbb{L}.
 \end{aligned}
 \tag{4.2}$$

By symmetry, we also have

$$[X] = [\mathbb{P}^4] + 3[\mathbb{P}^1]\mathbb{L} + [R_M]\mathbb{L} + 3\mathbb{L}^2 - 3\mathbb{L}.
 \tag{4.3}$$

Subtracting (4.3) from (4.2), we get

$$([R_L] - [R_M])\mathbb{L} = 0.$$

Next we show that $[R_L] \neq [R_M]$, and it is sufficient to show that $[R_L] \neq [R_M]$ modulo \mathbb{L} . According to [LL03], $[R_L] = [R_M] \bmod \mathbb{L}$ if and only if R_L and R_M are stably birational. Because a K3 surface is not rationally connected, this implies that R_L and R_M are birational and thus isomorphic, contradicting Theorem 3.1. \square

5. Exclusion of alternative constructions

This section shows that there exists just one class of Cremona transformations of \mathbb{P}^4 that can be resolved by blowing up an irreducible surface S with transverse double points: the class constructed in §2. Recall that [CK89] classified the case where S is smooth.

THEOREM 5.1. *Let $S \subset \mathbb{P}^4$ be an irreducible surface of degree d with $\delta > 0$ transverse double points. Assume there exists a Cremona transformation*

$$f : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$$

resolved by blowing up S . Let n and ξ denote the degrees of the homogeneous forms inducing f and f^{-1} respectively, and m the multiplicity of S in the base locus. Then we have

$$n = \xi = 4, \quad m = 1, \quad \delta = 3,$$

and S is obtained by projecting a degree-12 K3 surface from three points.

The remainder of this section is devoted to the proof of Theorem 5.1.

5.1 Extracting Diophantine equations

By Lemmas 1.1 and 1.2, equation (1.3) can be expressed as

$$\xi = n^3 - 3nm^2d + m^3(K_\Sigma C + 5d).
 \tag{5.1}$$

Similarly, equation (1.1) can be expressed as

$$1 = n^4 - 6n^2m^2d + 4nm^3(K_\Sigma C + 5d) - m^4(15d + 5K_\Sigma C + c_2(\Sigma) - 6\delta)
 \tag{5.2}$$

and equivalently as

$$1 = n^4 - 6n^2m^2d + 4nm^3(K_\Sigma C + 5d) + m^4(d^2 - 25d - 10K_\Sigma C - K_\Sigma^2 + 4\delta).
 \tag{5.2'}$$

The two formulas follow from the two expressions Lemma 1.2(3) and (3') for E^4 , respectively. The right-hand sides of these equations are arranged as polynomials in n and m . Note that only the coefficients of m^4 reflect the appearance of transverse double points.

5.2 Enumeration of combinatorial cases

LEMMA 5.2. *Only the following (n, m, ξ) can occur:*

| | n | m | ξ |
|-----|-----|-----|-------|
| (a) | 3 | 1 | 2 |
| (b) | 4 | 1 | 4 |
| (c) | 7 | 2 | 3 |
| (d) | 9 | 2 | 9 |
| (e) | 43 | 10 | 7 |
| (f) | 24 | 5 | 24 |
| (g) | 49 | 10 | 49 |

Proof. In the smooth case, the same list [CK89, Theorem 1.6] is obtained by using [CK89, Lemma 0.2] and [CK89, Formulae 0.3]. The proof of the former proceeds unchanged even with the transverse double points. The latter can be derived from (5.1) and (5.2) and only the terms with power of m up to two matter, so transverse double points do not change the result. Therefore the same elimination process works and we obtain the same list. □

5.3 Exclusion of cases

Here we show that only case (b) can occur.

LEMMA 5.3. *Cases (c) and (e) do not occur.*

Proof. The proof is similar to the smooth case [CK89, Lemma 3.2].

Assume case (c) holds. Then (5.1) reduces to

$$2K_{\Sigma}C = 11d - 85,$$

and (5.2) reduces to

$$465 = 62d - 2c_2(\Sigma) + 12\delta.$$

This is odd on the left and even on the right, a contradiction.

Assume case (e) holds. Now (5.1) reduces to

$$79d = 795 + 10K_{\Sigma}C,$$

so d is divisible by 5. On the other hand, (5.2) becomes

$$-34188 = -11094d + 1720(K_{\Sigma}C + 5d) - 100(15d + 5K_{\Sigma}C + c_2(\Sigma) - 6\delta).$$

Note that 5 divides the right but not the left, a contradiction. □

LEMMA 5.4. *Cases (d), (f) and (g) do not occur.*

Proof. Let I_S be the ideal sheaf of $S \subset \mathbb{P}^4$. Generally, the global sections of $I_S^m(n)$ and $\mathcal{O}_{P'}(M)$ are bijective canonically. So we have

$$h^0(\mathbb{P}^4, I_S^m(n)) = h^0(P', M) = 5 \tag{5.3}$$

by equation (1.4).

We prove the lemma case by case. In each case, we prove by contradiction in the following situations

$$h^0(\mathbb{P}^4, I_S(4)) = 0, = 1 \text{ and } \geq 2.$$

Assume case (d) holds. Suppose $h^0(\mathbb{P}^4, I_S(4)) = 0$. Consider the surjective map

$$\bigoplus_{k_1+k_2=9} H^0(\mathbb{P}^4, I_S(k_1)) \otimes H^0(\mathbb{P}^4, I_S(k_2)) \rightarrow H^0(\mathbb{P}^4, I_S^2(9)). \tag{5.4}$$

By hypothesis, $h^0(\mathbb{P}^4, I_S(k)) = 0$ for all $k \leq 4$. Since $k_1 + k_2 = 9$ implies $k_1 \leq 4$ or $k_2 \leq 4$, the left-hand side of (5.4) vanishes. Thus $h^0(\mathbb{P}^4, I_S^2(9)) = 0$, contradicting (5.3).

Let X_0, \dots, X_4 be a basis of degree-one forms on \mathbb{P}^4 in what follows.

Suppose $h^0(\mathbb{P}^4, I_S(4)) = 1$. Let $A \in H^0(\mathbb{P}^4, I_S(4))$ be a generator. This forces $h^0(\mathbb{P}^4, I_S(k)) = 0$ for all $k \leq 3$. It follows that $H^0(\mathbb{P}^4, I_S^2(8))$ is generated by A^2 . Then (5.3) indicates that A^2X_0, \dots, A^2X_4 form a basis for $H^0(\mathbb{P}^4, I_S^2(9))$. As a result, the linear system $|I_S^2(9)|$ defines an automorphism of \mathbb{P}^4 instead of a Cremona transformation.

Suppose $h^0(\mathbb{P}^4, I_S(4)) \geq 2$. Let $A, B \in H^0(\mathbb{P}^4, I_S(4))$ be independent. Then A^2 and AB are independent in $H^0(\mathbb{P}^4, I_S^2(8))$. We claim that there exists an i such that A^2X_i is not a linear combination of $ABX_j, j = 0, \dots, 4$. Suppose not, that is, $A^2X_i = ABL_i$ for some linear form $L_i, i = 0, \dots, 4$. Then we have $A/B = L_0/X_0 = L_1/X_1$, which implies that $L_0 = X_0L_1/X_1$, so X_1 divides L_0 . Therefore $A/B = L_1/X_1$ is a scalar, thus A and B are dependent, a contradiction. As a result, there exists an i such that A^2X_i and ABX_0, \dots, ABX_4 form an independent subset of $H^0(\mathbb{P}^4, I_S^2(9))$. Thus $h^0(\mathbb{P}^4, I_S^2(9)) \geq 6 > 5$, a contradiction.

Assume case (f) holds. Suppose $h^0(\mathbb{P}^4, I_S(4)) = 0$. Then $h^0(\mathbb{P}^4, I_S(k)) = 0$ for all $k \leq 4$. Now we consider the map

$$\bigoplus_{k_1+\dots+k_5=24} H^0(\mathbb{P}^4, I_S(k_1)) \otimes \dots \otimes H^0(\mathbb{P}^4, I_S(k_5)) \rightarrow H^0(\mathbb{P}^4, I_S^5(24)). \tag{5.5}$$

At least one $k_i \leq 4, i = 1, \dots, 5$, if their sum equals 24. Hence the left-hand side of (5.5) vanishes. Thus $h^0(\mathbb{P}^4, I_S^5(24)) = 0 \neq 5$.

Suppose $h^0(\mathbb{P}^4, I_S(4)) \geq 1$. Let $A \in H^0(\mathbb{P}^4, I_S(4))$ be a non-zero element. Then $A^5 \in H^0(\mathbb{P}^4, I_S^5(20))$. Multiplication by A^5 defines an injection

$$\cdot A^5 : H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(4)) \hookrightarrow H^0(\mathbb{P}^4, I_S^5(24)).$$

Thus $h^0(\mathbb{P}^4, I_S^5(24)) \geq \binom{8}{4} = 70 > 5$, a contradiction.

The elimination of case (g) is similar to that of case (f). In case (g), we use the surjection

$$\bigoplus_{k_1+\dots+k_{10}=49} H^0(\mathbb{P}^4, I_S(k_1)) \otimes \dots \otimes H^0(\mathbb{P}^4, I_S(k_{10})) \rightarrow H^0(\mathbb{P}^4, I_S^{10}(49))$$

to rule out the situation $h^0(\mathbb{P}^4, I_S(4)) = 0$. If $H^0(\mathbb{P}^4, I_S(4))$ contains $A \neq 0$, then multiplication of A^{10} with 9-forms produces $\binom{13}{4} = 715$ independent elements in $H^0(\mathbb{P}^4, I_S^{10}(49))$, which is not allowed. □

LEMMA 5.5. *In cases (a) and (b) we have:*

| | (a)(3, 1, 2) | (b)(4, 1, 4) |
|--------------------------------|----------------------------|------------------------------|
| d | ≤ 8 | ≤ 15 |
| $K_{\Sigma}C$ | $4d - 25$ | $7d - 60$ |
| K_{Σ}^2 | $d^2 - 11d + 4\delta + 30$ | $d^2 + d + 4\delta - 105$ |
| $c_2(\Sigma)$ | $19d - 95 + 6\delta$ | $46d - 405 + 6\delta$ |
| $12\chi(\mathcal{O}_{\Sigma})$ | $d^2 + 8d - 65 + 10\delta$ | $d^2 + 47d - 510 + 10\delta$ |
| $g(C)$ | $\frac{5d - 23}{2}$ | $4d - 29$ |

The invariants d and δ satisfy $(d - 5)^2 = 2\delta$ in case (a) and $(d - 10)(d - 15) = 2\delta$ in case (b).

Proof. In order to compute the invariants in the list, we first use (5.1) to express $K_\Sigma C$ in d with given $n, m,$ and ξ . Then (5.2) (respectively, (5.2')) allows us to express $c_2(\Sigma)$ (respectively, K_Σ^2) in d and δ . We compute $12\chi(\mathcal{O}_\Sigma)$ and $g(C)$ by Noether's formula and the genus formula, respectively. The upper bound for d comes from the inequality $d < (n/m)^2$ which holds generally [CK89, Formulae 0.3(v)].

We have $h^0(\mathbb{P}^4, I_S(n)) = h^0(P', M) = 5$ by (1.4). On the other hand, $h^1(\mathbb{P}^4, I_S(n)) = 0$ by [Dol12, Proposition 7.1.4]. Hence

$$h^0(\mathbb{P}^4, I_S(n)) = \chi(\mathbb{P}^4, I_S(n)) = \chi(P, I_{S'}(n)),$$

where the second equality follows from the functoriality of the Euler characteristic. The short exact sequence

$$0 \rightarrow I_{S'}(nL - 2\Sigma_i E_i) \rightarrow \mathcal{O}_P(nL - 2\Sigma_i E_i) \rightarrow \mathcal{O}_{S'}(nC - 2\Sigma_i(Q'_i + Q''_i)) \rightarrow 0$$

implies that

$$\chi(P, I_{S'}(n)) = \chi(P, nL - 2\Sigma_i E_i) - \chi(S', nC - 2\Sigma_i(Q'_i + Q''_i)).$$

$\chi(P, nL - 2\Sigma_i E_i)$ counts the dimension of the space of degree- n polynomials singular along Δ , so

$$\chi(P, nL - 2\Sigma_i E_i) = \binom{n+4}{4} - 5\delta.$$

By the previous computations and the Riemann–Roch formula, we have

$$\chi(S', nC - 2\Sigma_i(Q'_i + Q''_i)) = \begin{cases} \frac{1}{12}(d^2 - 10d + 385 - 62\delta) & \text{for (a),} \\ \frac{1}{12}(d^2 - 25d + 930 - 62\delta) & \text{for (b),} \end{cases}$$

whence

$$\chi(P, I_{S'}(n)) = \begin{cases} -\frac{1}{12}(d^2 - 10d - 35 - 2\delta) & \text{for (a),} \\ -\frac{1}{12}(d^2 - 25d + 90 - 2\delta) & \text{for (b).} \end{cases}$$

Then the two equations are obtained by setting $\chi(P, I_{S'}(n)) = 5$. □

LEMMA 5.6. *Case (a) does not occur.*

Proof. Assume (a) is satisfied. Then the same argument as in [CK89, Theorem 3.3] implies that $d = 5$. By Lemma 5.5 we have $\delta = 0$. □

5.4 Geometric analysis of the remaining case

To complete the proof of Theorem 5.1, it remains to analyze the last possible case.

LEMMA 5.7. *We have $(d, \delta) = (8, 7)$ or $(9, 3)$. The invariants in these cases are:*

| (d, δ) | $K_\Sigma C$ | K_Σ^2 | $c_2(\Sigma)$ | $\chi(\mathcal{O}_\Sigma)$ | $g(C)$ |
|---------------|--------------|--------------|---------------|----------------------------|--------|
| $(8, 7)$ | −4 | −5 | 5 | 0 | 3 |
| $(9, 3)$ | 3 | −3 | 27 | 2 | 7 |

Proof. By the previous part only case (b) is allowed.

By Lemma 5.5, we have $d \leq 15$ and $g(C) = 4d - 29 \geq 0$. Hence $8 \leq d \leq 15$. Then $(d - 10)(d - 15) = 2\delta$ and our hypothesis $\delta > 0$ force $d = 8$ or 9 , which implies that $\delta = 7$ or 3 , respectively.

The invariants are computed directly by using Lemma 5.5. □

Consider the linear system $|K_\Sigma + C|$ for both cases of Lemma 5.7. We have $h^1(K_\Sigma + C) = 0$ by Kodaira vanishing and $h^2(K_\Sigma + C) = h^0(-C) = 0$ by Serre duality.

LEMMA 5.8. *The case $(d, \delta) = (8, 7)$ is not allowed.*

Proof. By the Riemann–Roch formula,

$$\begin{aligned} h^0(K_\Sigma + C) &= \chi(K_\Sigma + C) \\ &= \chi(\mathcal{O}_\Sigma) + \frac{1}{2}(K_\Sigma + C)C \\ &= 0 + \frac{1}{2}(-4 + 8) = 2. \end{aligned}$$

Because $c_2(\Sigma) = 5$, Σ cannot be \mathbb{P}^2 , $\mathbb{P}^1 \times \mathbb{P}^1$, or a minimal ruled surface. This implies that $\mathcal{O}_\Sigma(K_\Sigma + C)$ is generated by global sections [Som81, Proposition 2.2]. Hence the system $|K_\Sigma + C|$ defines a morphism

$$\phi : \Sigma \rightarrow \mathbb{P}^1,$$

the *adjunction mapping*.

Consider the Stein factorization

$$\begin{array}{ccc} \Sigma & \xrightarrow{\phi} & \mathbb{P}^1 \\ & \searrow r & \nearrow s \\ & \Sigma' & \end{array}$$

where r is a proper morphism with connected fibers and s is a finite morphism. By [Som81, (2.3)], this leads to two possible situations.

- (1) $\dim \phi(\Sigma) = 0$. Here we have $g(C) = 1$, a contradiction.
- (2) $\dim \phi(\Sigma) = 1$. Then there exists a \mathbb{P}^1 -bundle $\pi : R \rightarrow \Sigma'$ such that r factors as

$$\begin{array}{ccc} \Sigma & \xrightarrow{\epsilon} & R \\ & \searrow r & \downarrow \\ & & \Sigma', \end{array}$$

where Σ is the blow-up of R in at most one point of each fiber blown up, and C meets the generic fiber with degree two. Furthermore, the map s is an isomorphism except possibly if $g(C) = 3$ and $h^{1,0}(\Sigma) = 1$.

Let us analyze situation (2): the map s cannot be an isomorphism. Otherwise, R is a Hirzebruch surface and $\chi(\mathcal{O}_\Sigma) = \chi(\mathcal{O}_R) = 1$, a contradiction. Hence we obtain

$$g(\Sigma') = h^{1,0}(R) = h^{1,0}(\Sigma) = 1.$$

Then $\chi(\mathcal{O}_\Sigma) = 0$ and $c_2(\Sigma) = 5$ implies that Σ has Hodge diamond

$$\begin{array}{ccccc} & & 1 & & \\ & & & & \\ & 1 & & 1 & \\ & & & & \\ 0 & & 7 & & 0. \end{array}$$

Since the Néron–Severi group of R has rank two, we conclude that Σ is the blow-up of R along five points on distinct fibers, and R is ruled over the elliptic curve Σ' .

Let h be the class of a section on R and f be the class of a fiber so that

$$h^2 = m, \quad hf = 1, \quad \text{and} \quad f^2 = 0$$

for some integer m . According to the description of (2), the image of C in R gives a class $H = 2h + bf$ for some integer b and $C = \epsilon^*H - \sum_{i=1}^5 F_i$ where F_1, \dots, F_5 are the exceptional curves on Σ . Note that $K_\Sigma = \epsilon^*K_R + \sum_{i=1}^5 F_i$. Thus we have

$$\begin{aligned} 8 = C^2 = H^2 - 5 &\Rightarrow H^2 = 13, \\ -4 = K_\Sigma C = K_R H + 5 &\Rightarrow K_R H = -9, \end{aligned}$$

and consequently,

$$\begin{aligned} \chi(H) &= \chi(\mathcal{O}_R) + \frac{1}{2}H(H - K_R) \\ &= 0 + \frac{1}{2}(13 + 9) = 11. \end{aligned}$$

On the other hand, we can use the exact sequence

$$0 \rightarrow \mathcal{O}_R \rightarrow \mathcal{O}_R(h) \rightarrow \mathcal{O}_h(m) \rightarrow 0$$

to get $\chi(h) = m$, and then use

$$0 \rightarrow \mathcal{O}_R(h) \rightarrow \mathcal{O}_R(2h) \rightarrow \mathcal{O}_h(2m) \rightarrow 0$$

to obtain $\chi(2h) = 3m$. Then an induction on n with the sequence

$$0 \rightarrow \mathcal{O}_R(2h + (n - 1)f) \rightarrow \mathcal{O}_R(2h + nf) \rightarrow \mathcal{O}_f(2) \rightarrow 0$$


implies that $\chi(H) = \chi(2h + bf) = 3m + 3b$. But this implies $11 = \chi(H)$ is divisible by 3, a contradiction. \square

PROPOSITION 5.9. *If S has a transverse double point, then it can only be the image of a $K3$ surface $R \subset \mathbb{P}^7$ of degree 12 projected from three points on R , and the number of transverse double points must be $\delta = 3$.*

Proof. By the Riemann–Roch formula,

$$\begin{aligned} h^0(K_\Sigma + C) &= \chi(K_\Sigma + C) \\ &= \chi(\mathcal{O}_\Sigma) + \frac{1}{2}(K_\Sigma + C)C \\ &= 2 + \frac{1}{2}(3 + 9) = 8. \end{aligned}$$

Because $c_2(\Sigma) = 27$, Σ cannot be \mathbb{P}^2 , $\mathbb{P}^1 \times \mathbb{P}^1$, or a minimal ruled surface. It follows that $\mathcal{O}_\Sigma(K_\Sigma + C)$ is generated by global sections [Som81, Proposition 2.2]. Hence $|K_\Sigma + C|$ defines an adjunction morphism with Stein factorization



There are three possible situations [Som81, (2.3)].

- (1) $\dim \phi(\Sigma) = 0$. We have $g(C) = 1$, a contradiction.
- (2) $\dim \phi(\Sigma) = 1$. Then $r : \Sigma \rightarrow \Sigma'$ is again obtained by blowing up a \mathbb{P}^1 -bundle, with no more than one point in a fiber blown up. In particular, $1 \geq \chi(\mathcal{O}_\Sigma) = 2$, a contradiction.
- (3) $\dim \phi(\Sigma) = 2$. Then $r : \Sigma \rightarrow \Sigma'$ expresses Σ as the blow-up of a smooth surface Σ' along a finite set with $F \cdot C = 1$ for any positive-dimensional fiber F of r . Moreover, $s : \Sigma' \rightarrow \mathbb{P}^7$ is an embedding.

Now we are in situation (3). Let F_1, \dots, F_k be the exceptional curves on Σ relative to r and let H be the very ample divisor on Σ' which defines s . Then

$$C = r^*H - \sum_{i=1}^k F_i \quad \text{and} \quad K_\Sigma = r^*K_{\Sigma'} + \sum_{i=1}^k F_i,$$

and it follows that

$$\begin{aligned} 9 &= C^2 = H^2 - k, \\ 3 &= K_\Sigma C = K_{\Sigma'} H + k, \\ -3 &= K_\Sigma^2 = K_{\Sigma'}^2 - k. \end{aligned}$$

By the Riemann–Roch formula,

$$\begin{aligned} 8 &= \chi(H) = \chi(\mathcal{O}_{\Sigma'}) + \frac{1}{2}H(H - K_{\Sigma'}) \\ &= 2 + \frac{1}{2}((9 + k) - (3 - k)) \\ &= 5 + k, \end{aligned}$$

which implies that $k = 3$. Hence Σ is obtained by blowing up Σ' along three distinct points, and $\Sigma' \subset \mathbb{P}^7$ has

$$\deg(\Sigma') = H^2 = 12, \quad K_{\Sigma'} H = K_{\Sigma'}^2 = 0, \quad c_2(\Sigma') = 24, \quad \chi(\mathcal{O}_{\Sigma'}) = 2.$$

We claim that Σ' is a K3 surface. Indeed, its Kodaira dimension $\kappa \neq 2$ since $K_{\Sigma'} H = 0$. If $\kappa = 1$, then Σ' has minimal model R an elliptic surface, such that nK_R is numerically equivalent to a positive linear combination of some fiber classes if n is large enough [Bea96, Proposition IX.3]. This implies that $K_{\Sigma'}$ is numerically effective, which contradicts the fact that $K_{\Sigma'} H = 0$. If $\kappa = -\infty$, then $h^{1,0}(\Sigma') = 0$ and thus $1 \geq \chi(\mathcal{O}_{\Sigma'}) = 2$, a contradiction. As a result, Σ' has $\kappa = 0$ and thus is a K3 surface.

Besides, the birational map $R \dashrightarrow \Sigma \rightarrow S$ can be realized as the projection from three points on R . Furthermore, the fact that $\delta = 3$ can also be verified directly by the double-point formula as in § 2.3. □

ACKNOWLEDGEMENTS

We are grateful for conversations with Lev Borisov that inspired this work. The manuscript benefited from correspondence with Alexander Kuznetsov. The authors are grateful for the support of the National Science Foundation through grant DMS-1551514.

REFERENCES

- Bea96 A. Beauville, *Complex algebraic surfaces*, London Mathematical Society Student Texts, vol. 34, second edition (Cambridge University Press, Cambridge, 1996); translated from the 1978 French original by R. Barlow, with assistance from N. I. Shepherd-Barron and M. Reid.
- Bor15 L. Borisov, *The class of the affine line is a zero divisor in the Grothendieck ring*, Preprint (2015), [arXiv:1412.6194v3](https://arxiv.org/abs/1412.6194v3).
- CG72 C. H. Clemens and P. A. Griffiths, *The intermediate Jacobian of the cubic threefold*, Ann. of Math. (2) **95** (1972), 281–356.
- CK89 B. Crauder and S. Katz, *Cremona transformations with smooth irreducible fundamental locus*, Amer. J. Math. **111** (1989), 289–307.
- DGPS15 W. Decker, G.-M. Greuel, G. Pfister and H. Schönemann, *Singular 4-0-2 — A computer algebra system for polynomial computations*, <http://www.singular.uni-kl.de>, 2015.
- Dol12 I. V. Dolgachev, *Classical algebraic geometry: A modern view* (Cambridge University Press, Cambridge, 2012).
- Ful98 W. Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, vol. 2, second edition (Springer, Berlin, 1998).
- GS14 S. Galkin and E. Shinder, *The Fano variety of lines and rationality problem for a cubic hypersurface*, Preprint (2014), [arXiv:1405.5154v2](https://arxiv.org/abs/1405.5154v2).
- Has16 B. Hassett, *Cubic fourfolds, K3 surfaces, and rationality questions*, in *Rationality problems in algebraic geometry*, Lecture Notes in Mathematics, vol. 2172 (Springer, Cham, 2016).
- HLOY03 S. Hosono, B. H. Lian, K. Oguiso and S.-T. Yau, *Fourier–Mukai partners of a K3 surface of Picard number one*, in *Vector bundles and representation theory (Columbia, MO, 2002)*, Contemporary Mathematics, vol. 322 (American Mathematical Society, Providence, RI, 2003), 43–55.
- IM04 A. Iliev and D. Markushevich, *Elliptic curves and rank-2 vector bundles on the prime Fano threefold of genus 7*, Adv. Geom. **4** (2004), 287–318.
- IMOU16a A. Ito, M. Miura, S. Okawa and K. Ueda, *The class of the affine line is a zero divisor in the Grothendieck ring: via G_2 -Grassmannians*, Preprint (2016), [arXiv:1606.04210v2](https://arxiv.org/abs/1606.04210v2).
- IMOU16b A. Ito, M. Miura, S. Okawa and K. Ueda, *Derived equivalence and Grothendieck ring of varieties: via K3 surfaces of degree 12*, Preprint (2016), [arXiv:1612.08497v1](https://arxiv.org/abs/1612.08497v1).
- KS17 A. Kuznetsov and E. Shinder, *Grothendieck ring of varieties, D- and L-equivalence, and families of quadrics*, Selecta Math. (N.S.) (2017), doi:[10.1007/s00029-017-0344-4](https://doi.org/10.1007/s00029-017-0344-4).
- Kuz06 A. G. Kuznetsov, *Hyperplane sections and derived categories*, Izv. Math. **70** (2006), 447–547.
- Kuz16 A. Kuznetsov, *Derived equivalence of Ito–Miura–Okawa–Ueda Calabi–Yau 3-folds*, J. Math. Soc. Japan, to appear. Preprint (2016), [arXiv:1611.08386v1](https://arxiv.org/abs/1611.08386v1).
- LL03 M. Larsen and V. A. Lunts, *Motivic measures and stable birational geometry*, Mosc. Math. J. **3** (2003), 85–95; 259.
- Mar11 E. Markman, *A survey of Torelli and monodromy results for holomorphic-symplectic varieties*, in *Complex and differential geometry*, Springer Proceedings in Mathematics, vol. 8 (Springer, Heidelberg, 2011), 257–322.
- Mar16 N. Martin, *The class of the affine line is a zero divisor in the Grothendieck ring: an improvement*, C. R. Math. Acad. Sci. Paris **354** (2016), 936–939.
- Muk87 S. Mukai, *On the moduli space of bundles on K3 surfaces. I*, in *Vector bundles on algebraic varieties (Bombay, 1984)*, Tata Institute of Fundamental Research Studies in Mathematics, vol. 11 (Tata Institute of Fundamental Research, Bombay, 1987), 341–413.
- Muk88 S. Mukai, *Curves, K3 surfaces and Fano 3-folds of genus ≤ 10* , in *Algebraic geometry and commutative algebra, Vol. I* (Kinokuniya, Tokyo, 1988), 357–377.

- Muk99 S. Mukai, *Duality of polarized $K3$ surfaces*, in *New trends in algebraic geometry (Warwick, 1996)*, London Mathematical Society Lecture Note Series, vol. 264 (Cambridge University Press, Cambridge, 1999), 311–326.
- Ogu02 K. Oguiso, *$K3$ surfaces via almost-primes*, *Math. Res. Lett.* **9** (2002), 47–63.
- Orl97 D. O. Orlov., *Equivalences of derived categories and $K3$ surfaces*, *J. Math. Sci. (N.Y.)* **84** (1997), 1361–1381.
- Som81 A. J. Sommese, *Hyperplane sections*, in *Algebraic geometry (Chicago, Ill., 1980)*, Lecture Notes in Mathematics, vol. 862 (Springer, Berlin, 1981), 232–271.

Brendan Hassett bhassett@math.brown.edu
 Department of Mathematics, Brown University,
 Box 1917, 151 Thayer Street, Providence, RI 02912, USA

Kuan-Wen Lai kwlai@math.brown.edu
 Department of Mathematics, Brown University,
 Box 1917, 151 Thayer Street, Providence, RI 02912, USA