

THE JOURNAL OF AGRICULTURAL SCIENCE

EDITED BY

G. D. H. BELL, C.B.E., PH.D., F.R.S., Plant Breeding Institute, Cambridge

K. L. BLAXTER, PH.D., N.D.A. (HONS.), D.SC., F.R.S.E., F.R.S.,
Rowett Research Institute, Bucksburn, Aberdeen

G. W. COOKE, PH.D., F.R.I.C., Rothamsted Experimental Station, Harpenden

JOHN HAMMOND, Jr., M.A., School of Agriculture, Cambridge

PROF. SIR J. B. HUTCHINSON, C.M.G., SC.D., F.R.S., School of Agriculture,
Cambridge

H. L. PENMAN, O.B.E., PH.D., F.R.S., Rothamsted Experimental Station, Harpenden

W. J. RIDGMAN, M.A., School of Agriculture, Cambridge

H. H. ROGERS, B.SC., DIP.AG.SCI., Plant Breeding Institute, Cambridge

PROF. E. W. RUSSELL, C.M.G., PH.D., F.INST.P., Department of Soil Science,
University of Reading

F. YATES, C.B.E., SC.D., F.R.S., Rothamsted Experimental Station, Harpenden



CAMBRIDGE UNIVERSITY PRESS

BENTLEY HOUSE, 200 EUSTON ROAD, LONDON, N.W. 1

AMERICAN BRANCH: 32 EAST 57TH STREET, NEW YORK, N.Y. 10022

Published bi-monthly

Annual subscription £9 net (U.S.A. \$29.00)

Single parts £2 net (U.S.A. \$6.00)

Second class postage paid at New York, N.Y.

INSTRUCTIONS TO AUTHORS

Failure to comply with the Instructions to Authors may delay publication

Papers intended for publication should be submitted to Dr G. D. H. Bell, C.B.E., F.R.S., Plant Breeding Institute, Cambridge, or to one of the Associate Editors. It must be understood that, if accepted by the Editorial Board, the paper will not be published elsewhere in the same form.

MANUSCRIPTS. Papers, written in English, are accepted from any country and should be typed in double-spacing on one side of the paper with a margin at least 4 cm wide on the left-hand side. Authors should instruct typists on the style required. A top copy and one carbon copy should be submitted.

SHORT NOTES may be accepted provided they are based on adequate experimental evidence; special provision is made for their publication with the least possible delay. MSS should not exceed 1500 words in length or their equivalent. For tabulated matter allow 25 words per line of the table (including headings). For line illustrations allow 225 words per quarter of a page.

TABLES must be self-explanatory. They should be typed on separate sheets, numbered consecutively and carry an appropriate title. When possible, tables should be arranged so that they can be printed in the normal orientation of the text and without rules.

LINE DRAWINGS (with photocopies) should be 25 cm wide and drawn in black waterproof ink on Bristol board, graph paper with blue lines or tracing paper. Legends should be typed on a separate copy and numbering inserted lightly and clearly in soft pencil on the drawing. Tables and figures should not reproduce the same data.

The approximate position of tables and figures should be noted in the text.

PLATES should make a definite contribution to the value of the paper and the number submitted should be kept to a minimum. They should be good quality, unmounted, glossy prints and be lightly numbered in pencil on the reverse side. If several, or coloured, plates are submitted the author may be asked to contribute to the cost of reproducing them.

TITLE. The title must be specific and suitable for indexing by the mechanical methods now being employed. The full name and address of the institution in which the research has been carried out should be stated. Change of address may be given as a footnote. A short title, not exceeding 50 characters, should be provided for the running headlines.

STYLE. Experimental details and results should be recorded in the past tense and there should be no unnecessary repetition or loose phrases. Manuscripts are likely to be returned for modification if the presentation is not clear and precise.

LAYOUT. Authors are recommended to study 'General Notes on the Preparation of Scientific Papers' (Royal Society, London, 2nd edn, 1965). The Editorial Board do not insist on a rigid format but it is usually convenient to divide the paper into sections, e.g. Introduction, Materials and Methods, Results, and Discussion. An excess of headings and sub-headings should be avoided.

Authors are advised to note the following points: a detailed review of literature is not necessary; relevant details should be given of the plant or animal material, the experimental design and chemical or other techniques employed; mean results with their relevant standard errors should be presented rather than detailed data; the statistical methods used should be clearly stated; the discussion should relate the author's experiments to other work on the subject and give the author's conclusions. Footnotes should be avoided.

SUMMARIES of papers are placed at the beginning of the text and authors should submit MSS with the summaries so placed. The summary should be factual and suitable for use in abstracting journals; paragraphs should not be numbered.

REFERENCES. The bibliography must be given in the form—Surname of authors, initials, year of publication (in parentheses), title of paper, name of journal (abbreviated according to the *World List of Scientific Periodicals*, 4th edn, Butterworths, London), volume and pages of reference. References should be in alphabetical order. In the text a reference should be quoted by the author's name and date (in parentheses). Where there are more than two authors, the initial reference in the text should include the names of all authors, but subsequent citations should be in the form—first author followed by *et al.* Authors should check that all references in the text appear at the end of the paper and vice versa, and that names and dates correspond in the two places.

PROOF CORRECTION. Standard proof correction marks (British Standard 1219) should be made as legibly as possible in ink, not pencil. Directions to the printer which are not to be set up in type should be encircled. Captions to illustrations and all references should be checked. Queries marked by the printer should be answered. Proofs are provided in order that authors can check the correctness of the type-setting—excessive alterations may be charged to the author.

OFFPRINTS. Contributors will receive 25 copies of their papers free and can order others when they receive the proofs.

Dieldrin v The Desert Locust

Check-mate before the first move

Dieldrin is widely recognised as the best insecticide for preventing desert locusts ever leaving their breeding ground.

Small amounts of dieldrin solution safeguard vast food supplies. One litre sprayed over 2 hectares can control 40,000 hopper locusts. These would consume as much food in one day as 20 people, if allowed to reach the swarm stage.


Shell companies are situated in key areas to ensure that supplies are immediately available where they are needed.

Locust control is just one of dieldrin's many uses. It is being used all over the world to protect many different crops against a wide range of pests. It is also much used in industry and public health schemes. Dieldrin is available in a

range of formulations, and when used as directed is harmless to both applicators and consumers of crops.

For full details of approved uses and advice on application, please consult your Shell company or Shell Chemicals distributor.

Shell Chemicals



JOURNAL OF REPRODUCTION AND FERTILITY

The Journal publishes original papers, abstracts of Proceedings, and occasionally reviews and bibliographies, dealing with the morphology, physiology, biochemistry and pathology of reproduction in man and animals, and with the biological, medical and veterinary problems of fertility and lactation. Nine numbers are issued to give three volumes a year.

Papers appearing during 1969 include:

- D. H. HALE and R. B. SYMINGTON. Control of sexual activity in ranch cows by intramuscular and intravaginal administration of progestagens.
- P. J. QUINN, I. G. WHITE and K. W. CLELAND. Chemical and ultrastructural changes in ram spermatozoa after washing, cold shock and freezing.
- E. PARR. The role of inflammation in the uterine weight increase caused by an IUD.
- L. E. DONALDSON. Effect of continued daily injections of oxytocin on oestrous cycle length and reproductive tract morphology in the cow.
- P. E. MATTNER. Differential leucocytic responses to spermatozoa in the cervix and the uterus in ewes.
- H. BOYD, A. JAMESON and J. O. HALL. Fertilization rate and embryonic survival in dairy cows in relation to transferrin and J antigen groupings of sires and dams.
- C. E. ADAMS. Intraperitoneal insemination in the rabbit.

Brief Communications

- F. W. STRATMAN, N. L. FIRST and R. K. MEYER. Failure of androgen to affect ovulation and puberty in swine.
- J. C. HINDSON and B. M. SCHOFIELD. Notes on parturition in the sheep.
- D. R. LAMOND. Ovulation block with stilboestrol in the cow.
- P. J. MCCOSKER. Abnormal spermatozoan chromatin in infertile bulls.
- S. C. MILLS and T. W. SCOTT. Metabolism of fatty acids by testicular and ejaculated ram spermatozoa.
- P. J. QUINN and I. G. WHITE. Osmotic shock of ram spermatozoa.

Editorial correspondence should be sent to the Executive Editor, Journal of Reproduction and Fertility, 8 Jesus Lane, Cambridge, England, and business correspondence to the publishers.

The subscription rate for 1968 is £7 per volume or £21 per year (three volumes). U.S.A. and Canada \$23.50 per volume or \$70.00 per year.

BLACKWELL SCIENTIFIC PUBLICATIONS
5 ALFRED STREET OXFORD

Experimental Agriculture

Volume 5, Number 1
January 1969

- H. DOGGETT Yields of hybrid sorghums
- A. PINKERTON AND K. R. BOCK Parallel streak of sisal (*Agave sisalana*) in Kenya
- D. A. RIJKS AND J. F. HARROP Irrigation and fertilizer experiments on cotton at Mubuku, Uganda
- J. J. HARDON, C. N. WILLIAMS AND I. WATSON Leaf area and yield in the oil palm in Malaya
- P. D. TURNER Evaluation of fungicides for the control of *Helminthosporium heveae* on hevea rubber in Malaysia. I. Laboratory assessment
- R. L. WASTIE Evaluation of fungicides for the control of *Helminthosporium heveae*, on hevea rubber in Malaysia. II. Field tests
- R. J. HAGGAR Use of companion crops in grassland establishment in Nigeria
- B. WALKER Effects of feeding hay on the early wet season weight loss of cattle in Western Tanzania
- J. J. HARDON AND M. D. DAVIES Effects of vacuum-drying on the viability of oil palm pollen
- S. C. PEARCE Multivariate techniques of use in biological research

REVIEWS

35s. net (\$5.00) Annual subscription
£5 net (\$16.50) for four issues

**CAMBRIDGE
UNIVERSITY PRESS**

Just Published

M. G. Bekker

Introduction to Terrain-Vehicle Systems

In this authoritative and abundantly illustrated volume, Dr Bekker deals with the various applications of systems analysis to problems of off-the-road locomotion. Concentrating on methods of approach, Dr Bekker critically reviews existing knowledge of terrain-vehicle systems and integrates this knowledge from the viewpoint of practical engineering.

\$27.50

To be Published in May

Gilbert F. White

Strategies of American Water Management

'In a country of parching drought, stinking streams, and muddy floods'—to use the author's words—the management of America's water resources is a vital national issue. In his informed study, Dr White outlines a framework in which society's decisions as to water management may be examined and then discusses six of the major strategies which have actually been practiced in the United States.

\$5.95

AVAILABLE THROUGH YOUR
BOOKSELLER OR ORDER DIRECT



The UNIVERSITY OF
MICHIGAN PRESS
ANN ARBOR

Azodrin

does its job inside out—outside in

... and that means double trouble for cotton pests, because Shell's powerful new organophosphorus insecticide Azodrin works *two* ways.

By contact, immediately destroying insects on the plant surface. And systemically, by penetrating foliage rapidly where it kills sucking and chewing pests as they try to eat.

This sophisticated dual action makes Azodrin by far the most wide-ranging insecticide yet devised giving *continuing* control over *more* pests *more* effectively.


Azodrin has already been employed with outstanding success in the United States and

Central America, and is now proving world-wide to be one of the cotton grower's most powerful allies.

Suddenly pests that took a bite out of profits are losing their appetite.

For further information or supplies of Azodrin contact your Shell company or Shell chemicals distributor. Shell works today for your more profitable tomorrow.

Shell Chemicals



CORRIGENDUM

**Heterosis and combining ability in hybrids of
Avena sativa L. and *A. byzantina* C. Koch.**

BY G. JENKINS

J. agric. Sci. 72

p. 87 line 5:

The author regrets the following error:

'...with the s.c.a. residual mean square.'

Should read

'...with the error mean square from the lattice square analyses, after adjustment.'