

## CONVERGENCE OF SOLUTIONS TO SOME ELLIPTIC EQUATIONS IN BOUNDED NEUMANN THIN DOMAINS

CÉSAR R. DE OLIVEIRA<sup>✉</sup> and ALESSANDRA A. VERRI

(Received 19 June 2014; accepted 24 August 2015; first published online 20 November 2015)

Communicated by A. Hassell

### Abstract

Consider the (elliptic) stationary nonlinear reaction–diffusion equation in a sequence of bounded Neumann tubes in a space that is squeezed to a reference curve. It is supposed that the forcing term is square integrable and that the nonlinear one satisfies some growth and dissipative conditions. A norm convergence of the resolvents of the operators associated with the linear terms of such equations is proven, and this fact is used to provide new and simpler proofs of the asymptotic behaviour of the solutions to the full nonlinear equations (previously known in similar singular problems).

*2010 Mathematics subject classification:* primary 35B25; secondary 35K57, 81Q15.

*Keywords and phrases:* thin tubes, resolvent convergence, reduction of dimension.

### 1. Introduction

The possible effects of domain squeezing and dimensional reduction have been of great interest, particularly questions about effective operators, spectral behaviour and the dynamics of reaction–diffusion equations. Roughly, as some directions are squeezed, the first task is to find possible remaining subspaces of the Hilbert space on which the operator acts; the next is to investigate the behaviour of the systems restricted to such subspaces and also the possible contribution from the process of discarding dimensions. Usually it is not an easy task to carry out all the estimates needed, since they involve extremely singular limits, and effective operators and dynamics are sensitive to the boundary conditions on the border of the regions of interest [2, 3, 5–9, 12, 13, 15–17].

Here we concentrate on solutions to (elliptic, linear and nonlinear) stationary reaction–diffusion equations in some bounded domains (of plane and space) with Neumann boundary conditions. Our main goals are twofold: by considering certain domains, we will rigorously reinforce the choice of the remaining subspace after dimensional reduction, as is commonly done in the literature, and then we will present

simpler proofs of the limit of the sequence of solutions to such equations in the squeezing process. Our techniques are mainly based on [2, 8] and previous works by the present authors [11, 13].

Let  $s \in I$  denote the arc length parameter of the curve  $r(s)$  in  $R^3$  of class  $C^3$ , with  $I \subset R$  being an open and bounded interval, and let  $k(s)$  and  $\tau(s)$  be its curvature and torsion at the point  $r(s)$ , respectively. Pick  $S \neq \emptyset$ : an open, bounded, smooth and connected subset of  $R^2$ . Build a tube (waveguide) in  $R^3$  by properly moving the region  $S$  along  $r(s)$ ; at each point  $r(s)$  the cross-section region  $S$  may present a (continuously differentiable) rotation angle  $\alpha(s)$  (see details in Section 2). For each small enough  $\varepsilon > 0$ , one can realize this same construction with the region  $\varepsilon S$  and so obtain a sequence of tubes  $\Omega_\varepsilon$  in  $R^3$  that is squeezed to the reference curve  $r$  as  $\varepsilon \rightarrow 0$ .

Let  $-\Delta$  be the (negative) Laplacian in  $\Omega_\varepsilon$  with Neumann condition at the boundary  $\partial\Omega_\varepsilon$ . Pick  $\tilde{g}_\varepsilon \in L^2(\Omega_\varepsilon)$ , and consider the stationary reaction–diffusion equation

$$\begin{aligned}
 -\Delta u + u + f(u) &= \tilde{g}_\varepsilon \quad \text{in } \Omega_\varepsilon, \\
 \frac{\partial u}{\partial \eta_\varepsilon} &= 0 \quad \text{on } \partial\Omega_\varepsilon,
 \end{aligned}
 \tag{1.1}$$

where  $\eta_\varepsilon$  denotes the outward unit normal vector field to  $\partial\Omega_\varepsilon$ , and  $f : R \rightarrow R$  is either the null function or a  $C^2$ -function satisfying some growth and dissipative conditions (see Section 5). We are interested in the limiting behaviour of one sequence of solutions to (1.1) as  $\varepsilon \rightarrow 0$ . This subject has received the attention of several authors (see [1, 4, 9, 16, 18]).

We note that Equation (1.1) is elliptic and actually involves no diffusion, since there is no time dependence. Before we present the main focus of this paper, we will briefly recall some results that motivated our questions and studies.

Let  $\Lambda$  be an arbitrary smooth bounded domain in  $R^m \times R^n$ . Write  $(x, y)$  for a generic point of  $R^m \times R^n$  and, for each  $\varepsilon > 0$ , put  $\Lambda_\varepsilon = \{(x, \varepsilon y) : (x, y) \in \Lambda\}$ . Prizzi and Rybakowski [16] considered the reaction–diffusion equation

$$\begin{aligned}
 u_t - \Delta u + f(u) &= 0 \quad \text{in } \Lambda_\varepsilon, \\
 \frac{\partial u}{\partial \eta_\varepsilon} &= 0 \quad \text{on } \partial\Lambda_\varepsilon,
 \end{aligned}
 \tag{1.2}$$

where, as before,  $\eta_\varepsilon$  denotes the outward unit vector field to  $\partial\Lambda_\varepsilon$  and  $f : R \rightarrow R$  is a nonlinearity satisfying some growth and dissipative conditions ensuring that (1.2) generates a semiflow on the Sobolev space  $H^1(\Lambda_\varepsilon)$ .

By considering the stationary linear equation associated with (1.2) (that is,  $f \equiv 0$ ) and performing a dilation of the domain  $\Lambda_\varepsilon$  by a factor  $\varepsilon$  in the  $y$ -direction, one gets

$$\begin{aligned}
 -\Delta_x u - \frac{1}{\varepsilon^2} \Delta_y u &= 0 \quad \text{in } \Lambda, \\
 \langle \nabla_x u, \eta_x \rangle + \frac{1}{\varepsilon^2} \langle \nabla_y u, \eta_y \rangle &= 0 \quad \text{on } \partial\Lambda,
 \end{aligned}
 \tag{1.3}$$

with  $(\eta_x, \eta_y)$  denoting the components of the outward unitary normal vector field to  $\partial\Lambda$ . Consider the associated quadratic form

$$b_\varepsilon(u) = \int_\Lambda \left( |\nabla_x u|^2 + \frac{|\nabla_y u|^2}{\varepsilon^2} \right) dx dy, \quad u \in H^1(\Lambda),$$

and the subspace

$$\mathcal{K} := \{u \in H^1(\Lambda) : \nabla_y u = 0 \text{ almost everywhere in } \Lambda\}. \tag{1.4}$$

By direct inspection, as  $\varepsilon \rightarrow 0$ , one gets

$$b_\varepsilon(u) \rightarrow \begin{cases} \int_\Lambda |\nabla_x u|^2 dx dy & \text{if } u \in \mathcal{K}, \\ \infty & \text{if } u \in H^1(\Lambda) \setminus \mathcal{K}. \end{cases}$$

Thus, in order to establish a limit for  $b_\varepsilon(u)$  [16, 18] one just restricts study to the subspace  $\mathcal{K}$ . Note that  $\mathcal{K}$  is directly related to the fact that the first eigenvalue of the Neumann Laplacian in a bounded region is zero (so there is no need for a renormalization: see ahead), and the constant functions are the corresponding eigenfunctions.

By studying Equation (1.3), Prizzi and Rybakowski have proven that, in some strong sense, (1.2) has a limiting equation

$$u_t + Au + f(u) = 0,$$

as  $\varepsilon \rightarrow 0$ , where  $A$  is the self-adjoint operator associated with the quadratic form

$$\int_\Lambda |\nabla_x u|^2 dx, \quad u \in \mathcal{K}.$$

By using the same notation as above, given a bounded sequence  $(h_\varepsilon)_\varepsilon$  in  $L^2(\Lambda)$ , Silva [18] performed a similar analysis of the stationary linear problem

$$\begin{aligned} -\Delta_x u - \frac{1}{\varepsilon^2} \Delta_y u &= h_\varepsilon \quad \text{in } \Lambda, \\ \langle \nabla_x u, \eta_x \rangle + \frac{1}{\varepsilon^2} \langle \nabla_y u, \eta_y \rangle &= 0 \quad \text{on } \partial\Lambda, \end{aligned} \tag{1.5}$$

and studied the limiting behaviour of its solutions as  $\varepsilon \rightarrow 0$ . In comparison to [16], Silva presented a simpler and more direct proof of the operator convergence of resolvents.

We draw attention to the fact that this strategy to select the correct action of the limiting quadratic form may not be convenient in some settings. A simple example of dimensional reduction involving the Laplacian with Dirichlet boundary condition illustrates our point. Consider the sequence of tubes  $\Omega_\varepsilon$ , as defined at the beginning of this Introduction, and the particular case in which  $r(s)$  is a piece of a circumference with radius  $a > 0$  (and  $\alpha \equiv 0$ ); the quadratic form associated with the problem is  $\int_{\Omega_\varepsilon} |\nabla u|^2 ds dy$  with domain  $H_0^1(\Omega_\varepsilon)$ . In this case it is necessary to perform a renormalization. More exactly, since the first eigenvalue  $\lambda_0$  of the Dirichlet Laplacian

in  $H_0^1(S)$  is greater than zero, one considers

$$\int_{\Omega_\varepsilon} \left( |\nabla u|^2 - \frac{\lambda_0}{\varepsilon^2} |u|^2 \right) ds dy,$$

and after a natural change of variables it is transformed into

$$\int_{I \times S} \left[ \frac{1}{\gamma_\varepsilon} |u'|^2 + \frac{\gamma_\varepsilon}{\varepsilon^2} (|\nabla_y u|^2 - \lambda_0 |u|^2) \right] ds dy, \quad u \in H_0^1(I \times S), \tag{1.6}$$

where  $u'$  denotes the derivative with respect to variable  $s$ ,  $\gamma_\varepsilon(s, y) = 1 - \varepsilon y_1/a$  (here  $(s, y_1, y_2) = (s, y)$  denotes points of  $I \times S$ ).

If  $u_0 = u_0(y)$  is the positive and normalized eigenfunction associated with  $\lambda_0$ , the subspace corresponding to  $\mathcal{K}$  (see Equation (1.4)) in this setting is

$$\mathcal{J} := \{wu_0 : w \in H_0^1(I)\} \subset L^2(I \times S). \tag{1.7}$$

Restricted to  $\mathcal{J}$ , in the limit as  $\varepsilon \rightarrow 0$ , the quadratic form (1.6) becomes (see the Appendix for details of this convergence)

$$\int_I |w'|^2 ds, \quad w \in H_0^1(I), \tag{1.8}$$

and one could naively guess that (1.8) would be the correct limiting form. However, by some careful considerations, in [2, 11, 14] it was proven that, although (1.7) is really the correct remaining subspace, the curvature  $k = 1/a$  explicitly appears in the action of the limiting quadratic form, which is actually given by

$$\int_I \left( |w'|^2 - \frac{|w|^2}{4a^2} \right) ds, \quad w \in H_0^1(I). \tag{1.9}$$

Further, this limit is also obtained in the sense of norm resolvent convergence of the associated self-adjoint operators [11]. Such an additional (curvature) term with respect to (1.8) may be interpreted as a contribution from the discarded dimensions in the process of squeezing, so that special caution must be exercised in the selection of the action of the limiting quadratic forms.

In view of the example just mentioned, one purpose of this work is to present a more complete justification—through an important norm resolvent convergence—such that, for some regions (that is, particular cases of (1.2) and (1.5)), under dimensional reduction with Neumann boundary conditions, one may actually restrict the analysis to suitable ‘remaining’ subspaces (for example, (1.4)) and with *no* net contribution from the discarded dimensions in the process of squeezing (in contrast to the Dirichlet case (1.9)). This is the main content of Theorem 1.1 below.

Now we return to Equation (1.1) in tubes. To study the limiting behaviour of a sequence of its solutions as  $\varepsilon \rightarrow 0$ , we perform a change of variables which takes  $\Omega_\varepsilon$  onto the fixed domain  $\Omega := I \times S$  (we keep the notation  $(s, y_1, y_2)$  for a point of  $\Omega$ ). Thus, we obtain the equivalent equations

$$\begin{aligned} A_\varepsilon u + f(u) &= g_\varepsilon \quad \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} &= \langle B_\varepsilon u, \nu \rangle \quad \text{on } \partial\Omega, \end{aligned} \tag{1.10}$$

where  $g_\varepsilon(s, y) = \tilde{g}_\varepsilon(F_\varepsilon(s, y))$  (for the definitions of  $B_\varepsilon$  and  $F_\varepsilon$  and details of this change of variables, see Section 2),  $\nu$  is the unit outward normal to  $\partial\Omega$  and  $A_\varepsilon$  is the self-adjoint operator associated with the quadratic form

$$a_\varepsilon(u) = \int_\Omega \left( |u'| + \langle \nabla_y u, Ry \rangle (\tau + \alpha') \right)^2 + \frac{\beta_\varepsilon}{\varepsilon^2} |\nabla_y u|^2 + |u|^2 \Big) ds dy, \quad u \in H^1(\Omega).$$

The definition of  $\beta_\varepsilon(s, y)$  also appears in Section 2. We denote by  $\mathcal{E}_\varepsilon$  the set of solutions to Equation (1.10).

If  $1 = 1(y)$  denotes the constant function on the cross-section  $S$ , let

$$\mathcal{L} := \{w1 : w \in L^2(I)\},$$

with the decomposition  $L^2(\Omega) = \mathcal{L} \oplus \mathcal{L}^\perp$  and denote by  $P$  the orthogonal projection onto the subspace  $\mathcal{L}$ . In the Hilbert space  $L^2(I)$ , let  $A_0$  be the self-adjoint operator associated with the quadratic form

$$a_0(w) := \int_I (|w'|^2 + |w|^2) ds, \quad w \in H^1(I). \tag{1.11}$$

**THEOREM 1.1.** *There are two positive numbers  $C$  and  $\varepsilon_0$ , so that, for  $0 < \varepsilon < \varepsilon_0$ ,*

$$\|A_\varepsilon^{-1} - A_0^{-1}P\|_{L^2(\Omega)} \leq C\varepsilon.$$

To compare this theorem with the results of [2, 11, 14] mentioned above, we note that, in the Neumann case, some geometric characteristics of the tube (such as curvature, torsion and rotation angle) have no influence on effective operators. Further, at least for the regions we consider here, this norm resolvent convergence of Theorem 1.1 supports the idea that one may just restrict the initial problem to the subspace  $\mathcal{L}$  for small  $\varepsilon$ , since it is the subspace that ‘remains’ in the process, and the quadratic form (1.11) is the restriction of the form  $a_\varepsilon$  to  $\mathcal{L}$ ; no contribution from the discarded dimensions shows up.

Now we move on to convergence of solutions to the stationary problem (1.10). We begin with the null function  $f$ . Theorem 1.1 allows us to weaken the condition that  $(g_\varepsilon)_\varepsilon$  is a bounded sequence in  $L^2(\Omega)$ , as done in [18], and still obtain a similar result, that is, the following theorem.

**THEOREM 1.2.** *In the problem (1.10) with null function  $f$ , let  $(g_\varepsilon)_\varepsilon$  be a sequence in  $L^2(\Omega)$  such that  $\|Pg_\varepsilon\|_{L^2(\Omega)} < c$ , for all small enough  $\varepsilon > 0$  and some constant  $c > 0$ . Consider a sequence  $(u_\varepsilon)_\varepsilon$ ,  $u_\varepsilon \in \mathcal{E}_\varepsilon$ . Then there exist  $g_0 \in \mathcal{L}$ ,  $u_0 := A_0^{-1}g_0 \in \text{dom } A_0$ , and a subsequence of  $(u_\varepsilon)_\varepsilon$ , denoted by the same symbol  $(u_\varepsilon)_\varepsilon$ , such that*

$$\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u_0\|_{L^2(\Omega)} = 0. \tag{1.12}$$

However, if  $\|g_\varepsilon\|_{L^2(\Omega)} \leq c$ , for all small enough  $\varepsilon > 0$ , then more can be said, that is,

$$\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u_0\|_{H^1(\Omega)} = 0. \tag{1.13}$$

In the general case, that is, with nonzero  $f$ , we have two possibilities, which are discussed in Theorems 1.3 and 1.4.

**THEOREM 1.3.** *Let  $f : R \rightarrow R$  be a nonlinear  $C^2$ -function satisfying*

$$f(0) = 0, \quad -f'(x) \leq K_1, \quad \forall x \in R, \tag{1.14}$$

$$\limsup_{|x| \rightarrow \infty} \frac{-f(x)}{x} \leq 0, \tag{1.15}$$

$$|f'(x)| \leq K_2(1 + |x|^\gamma), \quad \forall x \in R, \tag{1.16}$$

with some  $0 \leq \gamma \leq 2$ ; assume also that it is bounded and Lipschitz. Let  $(g_\varepsilon)_\varepsilon$  be a sequence in  $L^2(\Omega)$  such that  $\|Pg_\varepsilon\|_{L^2(\Omega)} < c$ , for all  $\varepsilon > 0$  small enough and for some constant  $c$ . Consider a sequence  $(u_\varepsilon)_\varepsilon$ ,  $u_\varepsilon \in \mathcal{E}_\varepsilon$ . Then there exist  $g_0 \in L^2(I)$ ,  $u_0 \in \text{dom } A_0 \subset L^2(I)$  so that  $u_0 = A_0^{-1}P(-f(u_0) + g_0)$ , and a subsequence of  $(u_\varepsilon)_\varepsilon$ , denoted by the same symbol  $(u_\varepsilon)_\varepsilon$ , such that

$$\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u_0\|_{L^2(\Omega)} = 0.$$

Note that, since  $f$  is a nonlinear function, the control on the term  $Pf(u)$ ,  $u \in H^1(\Omega)$ , can be far from trivial. Thus, if  $f$  is neither bounded nor Lipschitz, we will impose the condition  $\|g_\varepsilon\|_{L^2(\Omega)} < c$  on the forcing terms. In this case, we have the following theorem.

**THEOREM 1.4.** *Let  $f : R \rightarrow R$  be a nonlinear, bounded and Lipschitz  $C^2$ -function satisfying conditions (1.14)–(1.16), and  $(g_\varepsilon)_\varepsilon$  a sequence in  $L^2(\Omega)$  such that  $\|g_\varepsilon\|_{L^2(\Omega)} < c$ , for all small enough  $\varepsilon > 0$ . Consider a sequence  $(u_\varepsilon)_\varepsilon$ ,  $u_\varepsilon \in \mathcal{E}_\varepsilon$ . Then, there exist  $g_0 \in L^2(\Omega)$ ,  $u_0 \in \text{dom } A_0 \subset L^2(I)$  so that  $u_0 = A_0^{-1}P(-f(u_0) + g_0)$ , and a subsequence of  $(u_\varepsilon)_\varepsilon$ , denoted by the same symbol  $(u_\varepsilon)_\varepsilon$ , such that*

$$\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u_0\|_{H^1(\Omega)} = 0. \tag{1.17}$$

Although the settings in Theorems 1.3 and 1.4 have been previously considered, here, due to Theorem 1.1, we have simpler proofs on top of additional justification of the choice of remaining subspaces and the action of limiting forms. Further, in the case of Theorem 1.3, we were able to weaken the condition on  $g_\varepsilon$  with respect to [16, 18], by only requiring that the projection  $\|Pg_\varepsilon\|$  is uniformly bounded.

The above results have similar versions to strips in the plane, again built along differentiable curves  $r$  but with cross sections given by bounded intervals. The proofs are even simpler than in the case of tubes, and we do not explicitly discuss them here.

In addition, we are able to say something about the asymptotic behaviour of some semilinear equations in thin planar domains as in particular cases studied by Hale and Raugel [9]. Let  $g : [a, b] \rightarrow R$ ,  $-\infty < a < b < +\infty$ , be a  $C^3$ -function delimiting a planar region, that is, for each  $\varepsilon > 0$ , let

$$Q_\varepsilon := \{(s, y) \in J \times R : 0 < y < \varepsilon g(s)\}, \quad J := (a, b).$$

Consider the problem

$$\begin{aligned} u_t - \Delta u + f(u) &= 0 \quad \text{in } Q_\varepsilon, \\ \frac{\partial u}{\partial \nu_\varepsilon} &= 0 \quad \text{on } \partial Q_\varepsilon, \end{aligned} \tag{1.18}$$

where  $\nu_\varepsilon$  is the exterior unit normal vector field to  $\partial Q_\varepsilon$ , and  $f$  is a nonlinearity such that this equation generates a semiflow on  $H^1(Q_\varepsilon)$ . In [9] it was proven that the limiting semiflow, as  $\varepsilon \rightarrow 0$ , is the one generated by the one-dimensional boundary value problem

$$\begin{aligned} u_t - (1/g)(gu')' - f(u) &= 0 \quad \text{in } J, \\ u'(a) = u'(b) &= 0. \end{aligned}$$

In what follows, we shall present a simpler proof, with respect to [9], of the convergence of the solutions to the stationary equation associated with (1.18), namely,

$$\begin{aligned} -\Delta u + f(u) &= 0 \quad \text{in } Q_\varepsilon, \\ \frac{\partial u}{\partial \nu_\varepsilon} &= 0 \quad \text{on } \partial Q_\varepsilon. \end{aligned} \tag{1.19}$$

The first step in the study of Equation (1.19) is an analysis of its linear term, which is reduced, after an appropriate change of variables (see Section 6 for details), to

$$m_\varepsilon(u) = \int_Q \left[ \left( u' - \frac{g'}{2g}u \right)^2 + \frac{1}{\varepsilon^2 g^2} u_y^2 \right] ds dy,$$

where  $\text{dom } m_\varepsilon = H^1(Q)$  and  $Q := J \times (0, 1)$ . We denote by  $M_\varepsilon$  the self-adjoint operator associated with  $m_\varepsilon(u)$ . Thus, we pass to the equation

$$\begin{aligned} M_\varepsilon u + f(u) &= 0 \quad \text{in } Q, \\ \frac{\partial u}{\partial \nu} &= 0 \quad \text{on } \partial Q. \end{aligned} \tag{1.20}$$

Denote by  $\mathcal{E}_\varepsilon^g$  the set of solutions to (1.20). As before, we consider the subspace  $\tilde{\mathcal{J}} = \{w1 : w \in L^2(J)\}$  and the decomposition  $L^2(Q) = \tilde{\mathcal{J}} \oplus \tilde{\mathcal{J}}^\perp$ . Denote by  $\tilde{P}$  the orthogonal projection onto the subspace  $\tilde{\mathcal{J}}$ . Define the quadratic form

$$m(w) := \int_J \left( w' - \frac{g'}{2g}w \right)^2 ds, \quad \text{dom } m = H^1(J), \tag{1.21}$$

and denote by  $M$  the associated self-adjoint operator. In this case, we have a similar result to Theorem 1.1 (see Theorem 6.1) and, as a consequence, we shall prove the following theorem.

**THEOREM 1.5.** *Let  $f : R \rightarrow R$ , as in Theorem 1.4, and take a sequence  $(u_\varepsilon)_\varepsilon$ ,  $u_\varepsilon \in \mathcal{E}_\varepsilon^g$ ,  $\varepsilon \rightarrow 0$ . Then there exist  $u_0 \in \text{dom } M$ , so that  $u_0 = M^{-1}\tilde{P}(-f(u_0))$ , and a subsequence of  $(u_\varepsilon)_\varepsilon$ , denoted by the same symbol  $(u_\varepsilon)_\varepsilon$ , such that*

$$\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u_0\|_{H^1(Q)} = 0.$$

The rest of this paper is organized as follows. In Section 2, we present details of the construction of the tubular region  $\Omega_\varepsilon$  and we study the quadratic form associated with the Neumann Laplacian in this tube. Section 3 is dedicated to the proof of Theorem 1.1. In Section 4, we prove Theorem 1.2, whereas the proofs of Theorems 1.3 and 1.4 appear in Section 5 and, finally, Section 6 is reserved for the proof of Theorem 1.5.

### 2. Quadratic forms

In this section we are going to construct the region in which the Neumann Laplacian is considered, along with its associated quadratic form.

Let  $I$  be an open and bounded interval. We suppose that  $r : I \rightarrow R^3$  is a simple  $C^3$  curve in  $R^3$  parametrized by its arc length parameter  $s$ . The curvature of  $r$  at the position  $s$  is  $k(s) := \|r''(s)\|$ . We choose the usual orthonormal triad of vector fields  $\{T(s), N(s), B(s)\}$ , the so-called Frenet frame, given the tangent, normal and binormal vectors, respectively, moving along the curve and defined by

$$T = r'; \quad N = k^{-1}T'; \quad B = T \times N. \tag{2.1}$$

To justify the construction (2.1), it is assumed that  $k > 0$ , but if  $r$  has a piece of a straight line (that is,  $k = 0$  identically in this piece), usually one can choose a constant Frenet frame instead. It is possible to combine constant Frenet frames with the Frenet frame (2.1) to include other types of curves, for instance, curves with  $k(s) > 0$  only on a compact interval of values of  $s$  (and so obtaining a global  $C^2$  Frenet frame (see [10, Theorem 1.3.6]). In each situation we assume that a global Frenet frame exists and that the Frenet equations are satisfied, that is,

$$\begin{pmatrix} T' \\ N' \\ B' \end{pmatrix} = \begin{pmatrix} 0 & k & 0 \\ -k & 0 & \tau \\ 0 & -\tau & 0 \end{pmatrix} \begin{pmatrix} T \\ N \\ B \end{pmatrix}, \tag{2.2}$$

where  $\tau(s)$  is the torsion of  $r(s)$ , actually defined by (2.2). Let  $\alpha : I \rightarrow R$  be a bounded  $C^1$  function so that  $\alpha(s_0) = 0$  ( $s_0$  is a fixed point of  $I$ ), and  $S$  an open, bounded, connected and smooth (nonempty) subset of  $R^2$ . For small enough  $\varepsilon > 0$  and  $y = (y_1, y_2) \in S$ , write

$$\vec{x}(s, y) = r(s) + \varepsilon y_1 N(s) + \varepsilon y_2 B(s)$$

and consider the domain

$$\Omega_\varepsilon = \{\vec{x}(s, y) \in R^3 : s \in I, y = (y_1, y_2) \in S\},$$

where

$$\begin{aligned} N_\alpha(s) &:= \cos \alpha(s)N(s) + \sin \alpha(s)B(s), \\ B_\alpha(s) &:= -\sin \alpha(s)N(s) + \cos \alpha(s)B(s). \end{aligned}$$



Hence, this tube  $\Omega_\varepsilon$  is obtained by putting the region  $\varepsilon S$  along the curve  $r(s)$ , which is simultaneously rotated by an angle  $\alpha(s)$  with respect to the cross section at the position  $s = s_0$ . We suppose that  $k, \tau + \alpha' \in L^\infty(I)$ .

In order to study the Neumann Laplacian  $-\Delta$  in  $\Omega_\varepsilon$  we initially consider the corresponding family of quadratic forms

$$\hat{a}_\varepsilon(\psi) := \int_{\Omega_\varepsilon} (|\nabla\psi|^2 + |\psi|^2) d\vec{x}, \quad \psi \in \text{dom } \hat{a}_\varepsilon = H^1(\Omega_\varepsilon). \tag{2.3}$$

Consider the mapping

$$F_\varepsilon : I \times S \rightarrow \Omega_\varepsilon$$

$$(s, y) \mapsto r(s) + \varepsilon y_1 N_\alpha(s) + \varepsilon y_2 B_\alpha(s),$$

that will be used to perform an important change of variables. The condition  $k \in L^\infty(I)$ , is to guarantee that  $F_\varepsilon$  will be a (global) diffeomorphism for small  $\varepsilon$ . With this change of variables we work with a fixed region  $I \times S$  for all  $\varepsilon > 0$ : more precisely, in the new variables the domain of the quadratic form (2.3) turns out to be  $H^1(I \times S)$ . On the other hand, the price to be paid is a nontrivial Riemannian metric  $G = G_\varepsilon$  which is induced by  $F_\varepsilon$ , that is,

$$G = (G_{ij}), \quad G_{ij} = \langle e_i, e_j \rangle = G_{ji}, \quad 1 \leq i, j \leq 3,$$

where

$$e_1 = \frac{\partial F_\varepsilon}{\partial s}, \quad e_2 = \frac{\partial F_\varepsilon}{\partial y_1}, \quad e_3 = \frac{\partial F_\varepsilon}{\partial y_2}.$$

Some calculations show that, in the Frenet frame,

$$W = \begin{pmatrix} e_1 \\ e_2 \\ e_3 \end{pmatrix} = \begin{pmatrix} \beta_\varepsilon & -\varepsilon(\tau + \alpha')\langle z_\alpha^\perp, y \rangle & \varepsilon(\tau + \alpha')\langle z_\alpha, y \rangle \\ 0 & \varepsilon \cos \alpha & \varepsilon \sin \alpha \\ 0 & -\varepsilon \sin \alpha & \varepsilon \cos \alpha \end{pmatrix},$$

where

$$\beta_\varepsilon(s, y) = 1 - \varepsilon k(s)\langle z_\alpha, y \rangle, \quad z_\alpha = (\cos \alpha, -\sin \alpha) \quad \text{and} \quad z_\alpha^\perp = (\sin \alpha, \cos \alpha).$$

The inverse matrix of  $W$  is given by

$$W^{-1} = \begin{pmatrix} 1/\beta_\varepsilon & (\tau + \alpha')y_2/\beta_\varepsilon & -(\tau + \alpha')y_1/\beta_\varepsilon \\ 0 & (1/\varepsilon) \cos \alpha & -(1/\varepsilon) \sin \alpha \\ 0 & (1/\varepsilon) \sin \alpha & (1/\varepsilon) \cos \alpha \end{pmatrix}.$$

Note that  $WW^t = G$  and  $\det W = |\det G|^{1/2} = \varepsilon^2 \beta_\varepsilon$ . Since  $k$  is a bounded function, for small enough  $\varepsilon$ ,  $F_\varepsilon$  does not vanish in  $I \times S$ . Thus  $\beta_\varepsilon > 0$ , and  $F_\varepsilon$  is a local diffeomorphism. In the case where  $F_\varepsilon$  is injective (again by requiring that  $\varepsilon > 0$  is small), a global diffeomorphism is obtained.

Introducing the notation

$$\|u\|_G^2 = \int_{I \times S} |u|^2 \beta_\varepsilon ds dy,$$

and the unitary transformation

$$U_\varepsilon : L^2(\Omega_\varepsilon) \rightarrow L^2(I \times S, \beta_\varepsilon)$$

$$u \mapsto \varepsilon u \circ F_\varepsilon,$$

we obtain, from (2.3), the sequence of quadratic forms

$$\tilde{a}_\varepsilon(U_\varepsilon u) := \|W^{-1}\nabla(U_\varepsilon u)\|_G^2 + \|U_\varepsilon u\|_G^2.$$

Again, since  $k \in L^\infty(I)$ ,  $\beta_\varepsilon \rightarrow 1$  uniformly in  $I \times S$ , as  $\varepsilon \rightarrow 0$ . Therefore, the spaces  $L^2(I \times S, \beta_\varepsilon)$  and  $L^2(I \times S)$  are topologically equivalent and the strong convergence in  $L^2(I \times S, \beta_\varepsilon)$  is equivalent to the convergence in the fixed space  $L^2(I \times S)$ . Thus, we move on to work in  $L^2(I \times S)$ .

After the norms are written out, and using the same notation  $u$  for  $U_\varepsilon u$ , we obtain

$$\tilde{a}_\varepsilon(u) = \int_{I \times S} \left( \frac{1}{\beta_\varepsilon} |u'| + \langle \nabla_y u, Ry \rangle (\tau + \alpha') \right)^2 + \frac{\beta_\varepsilon}{\varepsilon^2} |\nabla_y u|^2 + \beta_\varepsilon |u|^2 \Big) ds dy,$$

$\text{dom } \tilde{a}_\varepsilon = H^1(I \times S)$ .

Now, for technical reasons, we define the quadratic form

$$a_\varepsilon(u) := \int_{I \times S} \left( |u'| + \langle \nabla_y u, Ry \rangle (\tau + \alpha') \right)^2 + \frac{\beta_\varepsilon}{\varepsilon^2} |\nabla_y u|^2 + |u|^2 \Big) ds dy,$$

$\text{dom } a_\varepsilon = H^1(I \times S)$ , which is very similar to the quadratic form  $\tilde{a}_\varepsilon(u)$ . We denote by  $\tilde{A}_\varepsilon$  and  $A_\varepsilon$  the self-adjoint operators associated with  $\tilde{a}_\varepsilon$  and  $a_\varepsilon$ , respectively. Since  $\beta_\varepsilon \rightarrow 1$  uniformly in  $I \times S$ , as  $\varepsilon \rightarrow 0$ , we have the following theorem, the proof of which is very similar to the proof of [13, Theorem 3.1] and for this reason will be omitted here.

**THEOREM 2.1.** *There exist numbers  $D > 0$  and  $\varepsilon_1 > 0$  so that*

$$\|\tilde{A}_\varepsilon^{-1} - A_\varepsilon^{-1}\|_{L^2(\Omega)} \leq D\varepsilon, \quad 0 < \varepsilon < \varepsilon_1.$$

Recall that  $\Omega = I \times S$ . Due to the changes of variables presented above and Theorem 2.1, instead of studying Equation (1.5) we move on to work with

$$A_\varepsilon u_\varepsilon + f(u_\varepsilon) = g_\varepsilon \quad \text{in } \Omega,$$

$$\frac{\partial u}{\partial \nu} = \langle B_\varepsilon u, \nu \rangle \quad \text{on } \partial\Omega,$$

where  $\nu$  is the unit outward normal to  $\partial\Omega$  and  $B_\varepsilon u = \beta_\varepsilon G^{-1} \nabla u$ .

Again, for technical reasons, we define the space  $H_\varepsilon^1(\Omega) := \{u \in L^2(\Omega) : \|u\|_{H_\varepsilon^1(\Omega)} < \infty\}$ , where

$$\|u\|_{H_\varepsilon^1(\Omega)} := \left( \int_\Omega (|u'| + \langle \nabla_y u, Ry \rangle (\tau + \alpha'))^2 + \beta_\varepsilon |\nabla_y u|^2 + |u|^2 \Big) ds dy \right)^{1/2}.$$

Since  $k, \tau + \alpha' \in L^\infty(I)$ , we can observe that  $\|u\|_{H^1(\Omega)}$  and  $\|u\|_{H_\varepsilon^1(\Omega)}$  are equivalent norms in  $H^1(\Omega)$ .

Another property that we are going to use throughout this text is that, since  $k \in L^\infty(I)$ , there exist  $E_1, E_2 > 0$  and  $\varepsilon_2 > 0$ , so that

$$E_1 < \beta_\varepsilon(s, y) < E_2, \quad \forall (s, y) \in I \times S, \quad 0 < \varepsilon < \varepsilon_2. \tag{2.4}$$

### 3. Reduction of dimension

As already mentioned in the Introduction, we consider the subspace  $\mathcal{L} = \{w1 : w \in L^2(I)\}$  and the orthogonal decomposition

$$L^2(\Omega) = \mathcal{L} \oplus \mathcal{L}^\perp.$$

For  $u \in L^2(\Omega)$ , we can write

$$u(s, y) = w(s) + v(s, y),$$

with  $w \in L^2(I)$  and  $v \in \mathcal{L}^\perp$ . Observe that  $v \in \mathcal{L}^\perp$  implies

$$\int_S v(s, y) dy = 0 \quad \text{a.e. } [s]. \quad (3.1)$$

Note that  $w \in H^1(\Omega)$  if  $w \in H^1(I)$ . For  $u \in H^1(\Omega)$ , write  $u = w + v$  with  $w \in H^1(I)$  and  $v \in H^1(\Omega) \cap \mathcal{L}^\perp$ .

For  $w \in H^1(I)$ , we may identify  $a_\varepsilon(w)$  with the ‘one-dimensional’ quadratic form (1.11), that is,

$$a_0(w) = \int_I (|w'|^2 + |w|^2) ds,$$

and recall that  $A_0$  is the associated self-adjoint operator.

The method of the proof of Theorem 1.1 is as follows. First one identifies suitable subspaces of the Hilbert space on which the operator acts and, second, one applies the powerful functional analytic technique of Friedlander and Solomyak [8]. The essence of the proof is an application of such a technique; however, showing that it can be applied, by demonstrating the required estimates, is in itself not trivial.

**PROOF OF THEOREM 1.1.** For  $u \in H^1(\Omega)$  write, as above,

$$u(s, y) = w(s) + v(s, y),$$

with  $w \in H^1(I)$  and  $v \in H^1(\Omega) \cap \mathcal{L}^\perp$ . Thus, the quadratic form  $a_\varepsilon(u)$  can be rewritten as

$$a_\varepsilon(w + v) = a_0(w) + 2a_\varepsilon(w, v) + a_\varepsilon(v).$$

Here,  $a_\varepsilon(u_1, u_2)$  denotes the bilinear form associated with the quadratic form  $a_\varepsilon(u)$ .

We are going to show that  $a_0(w)$ ,  $a_\varepsilon(v)$  and  $a_\varepsilon(w, v)$  satisfy conditions (3.2), (3.3), (3.4) and (3.5) in [8, Section 3], and so the theorem will follow.

Firstly, we observe that

$$a_0(w) \geq \|w\|_{L^2(I)}^2, \quad \forall w \in H^1(I),$$

and then condition (3.2) holds true.

Now, let  $\lambda_1 > 0$  be the second eigenvalue of the Neumann Laplacian in  $H^1(I)$ . For  $v \in H^1(\Omega) \cap \mathcal{L}^\perp$ ,

$$a_\varepsilon(v) \geq \frac{\lambda_1}{\varepsilon^2} \|v\|_{L^2(\Omega)}^2,$$

and (3.3) and (3.4) are satisfied.

Finally, we assert that there exist  $C_1 > 0$  and  $\varepsilon_0 > 0$ , so that, for  $0 < \varepsilon < \varepsilon_0$ ,

$$a_\varepsilon(w, v) \leq C_1 \varepsilon (a_0(w))^{1/2} (a_\varepsilon(v))^{1/2}.$$

In fact, due to property (3.1),

$$a_\varepsilon(w, v) = \int_\Omega w' \langle \nabla_y v, Ry \rangle (\tau + \alpha') ds dy.$$

By (2.4), we also note that

$$\frac{E_1}{\varepsilon^2} \int_\Omega |\nabla_y v|^2 ds dy \leq a_\varepsilon(v).$$

Let  $C_2 := \sup_{(s,y) \in \Omega} \{ \|Ry\| (\tau + \alpha')(s) \}$ . Thus,

$$\begin{aligned} \left| \int_\Omega w' \langle \nabla_y v, Ry \rangle (\tau + \alpha') ds dy \right| &\leq C_2 \left( \int_I |w'|^2 ds \right)^{1/2} \left( \int_\Omega |\nabla_y v|^2 ds dy \right)^{1/2} \\ &\leq \varepsilon (C_2 / \sqrt{E_1}) (a_\varepsilon(w))^{1/2} (a_\varepsilon(v))^{1/2}, \end{aligned}$$

and the condition (3.5) in [8] is also satisfied. □

#### 4. The linear problem

In this section we study the problem (1.10) and prove Theorem 1.2 presented in the Introduction. Lemma 5.2 in the next section will be used; note that it also holds in case of the (nonlinear) null function  $f$ .

**PROOF OF THEOREM 1.2.** First we prove (1.12). Let  $\varepsilon_n \rightarrow 0$ ,  $n \in N$  ( $n \rightarrow \infty$ ),  $A_n := A_{\varepsilon_n}$ ,  $g_n := g_{\varepsilon_n}$  and  $u_n := u_{\varepsilon_n}$  a sequence so that  $u_n \in \mathcal{E}_{\varepsilon_n}$ , that is,  $A_n u_n = g_n$ . Theorem 1.1 ensures that

$$\|A_n^{-1} g_n - A_0^{-1} P g_n\|_{L^2(\Omega)} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Since  $(P g_n)_n$  is a bounded sequence and  $A_0^{-1}$  is a compact operator, the sequence  $(A_0^{-1} P g_n)_n$  has a convergent subsequence. Thus, there exist  $u_0 \in L^2(I)$  and an infinite subset  $N_1$  of  $N$  so that

$$A_0^{-1} P g_n \rightarrow u_0 \quad \text{in } L^2(I), \quad n \in N_1.$$

On the other hand, there exist  $g_0 \in L^2(I)$  and an infinite subset  $N_2$  of  $N_1$  so that  $P g_n \rightarrow g_0$  in  $L^2(I)$ ,  $n \in N_2$ . Thus,  $A_0^{-1} P g_n \rightarrow A_0^{-1} g_0$ ,  $n \in N_2$ , and we conclude that  $A_0^{-1} P g_0 = u_0$  and  $u_n \rightarrow u_0$  in  $L^2(I \times S)$ ,  $n \in N_2$ .

To prove (1.13), we are going to show that, up to a subsequence,

$$\|u_n - u_0\|_{H^1_{\varepsilon_n}(\Omega)} \rightarrow 0. \tag{4.1}$$

Thus, since the norms  $\|u\|_{H^1(\Omega)}$  and  $\|u\|_{H^1_{\varepsilon_n}(\Omega)}$  are equivalent, this implies Theorem 1.2. □

Suppose that  $\|g_\varepsilon\| \leq c$ , for all small enough  $\varepsilon > 0$ . There exist  $\tilde{g}_0 \in L^2(\Omega)$  and an infinite subset  $N_3$  of  $N_2$  so that  $g_{\varepsilon_n} \rightarrow \tilde{g}_0$ ,  $n \in N_3$ . Write  $\tilde{g}_0 = P\tilde{g}_0 + P^\perp\tilde{g}_0$ . In this case,  $P\tilde{g}_0 = g_0$ . Now, by Lemma 5.2 and since the immersion  $H^1(\Omega) \rightarrow L^2(\Omega)$  is compact, we have  $u_n \rightarrow u_0$  in  $H^1(\Omega)$ ,  $n \in N_3$  (this convergence also holds in  $H^1_{\varepsilon_n}(\Omega)$ ). To conclude (4.1), it is enough to show that  $\|u_\varepsilon\|_{H^1_\varepsilon(\Omega)} \rightarrow \|u_0\|_{H^1(\Omega)}$ : for this, just observe that

$$\begin{aligned} \|u_0\|_{H^1(\Omega)}^2 &\leq \liminf_{n \in N_3} \int_{\Omega} (|u'_n + \langle \nabla_y u_n, Ry \rangle (\tau + \alpha')|^2 + \beta_{\varepsilon_n} |\nabla_y u_n|^2 + |u_n|^2) \, ds \, dy \\ &\leq \limsup_{n \in N_3} \int_{\Omega} (|u'_n + \langle \nabla_y u_n, Ry \rangle (\tau + \alpha')|^2 + \beta_{\varepsilon_n} |\nabla_y u_n|^2 + |u_n|^2) \, ds \, dy \\ &\leq \limsup_{n \in N_3} \int_{\Omega} (|u'_n + \langle \nabla_y u_n, Ry \rangle (\tau + \alpha')|^2 + \frac{\beta_{\varepsilon_n}}{\varepsilon_n^2} |\nabla_y u_n|^2 + |u_n|^2) \, ds \, dy \\ &= \limsup_{n \in N_3} \int_{\Omega} (A_n u_n) u_n \, ds \, dy \\ &= \limsup_{n \in N_3} \int_{\Omega} g_n u_n \, ds \, dy \\ &= \int_{\Omega} (P\tilde{g}_0 + P^\perp\tilde{g}_0) u_0 \, ds \, dy \\ &= \int_{\Omega} g_0 u_0 \, ds \, dy \\ &= \int_{\Omega} (A_0 u_0) u_0 \, ds \, dy \\ &= \|u_0\|_{H^1(\Omega)}^2. \end{aligned}$$

### 5. The nonlinear problem

In this section we present the required conditions for the function  $f$  in Equation (1.10). We assume that  $f : R \rightarrow R$  is a nonlinear  $C^2$ -function satisfying conditions (1.14)–(1.16), as in the statement of Theorem 1.3. The following related result is well known and its proof can be found in [15].

**LEMMA 5.1.** *Under conditions (1.14)–(1.16), the assignment  $u \mapsto f \circ u$  defines a map  $f : H^1(\Omega) \rightarrow L^2(\Omega)$ , which is Lipschitz continuous on every bounded set in  $H^1(\Omega)$ . Moreover, wherever  $u, u_1, u_2 \in H^1(\Omega)$  and  $\|u_1\|_{H^1(\Omega)}, \|u_2\|_{H^1(\Omega)} \leq L_1$  the following estimates hold.*

$$\begin{aligned} \|f(u)\|_{L^2(\Omega)} &\leq L_2(\|u\|_{L^2(\Omega)} + \tilde{L}_2 \|u\|_{H^1(\Omega)}^{\gamma+1}), \\ \|f(u_1) - f(u_2)\|_{L^2(\Omega)} &\leq \tilde{L}_1(1 + 2L_1^{2\gamma}) \|u_1 - u_2\|_{H^1(\Omega)}. \end{aligned}$$

Here,  $L_2, \tilde{L}_2$  and  $\tilde{L}_1$  are positive constants.

Note that, due to condition (1.15), for any  $\eta > 0$  there exists a positive constant  $C_\eta$  such that,

$$-f(x)x \leq \eta x^2 + C_\eta, \quad \forall x \in R. \tag{5.1}$$

This inequality will be useful later on.

**PROOF OF THEOREM 1.3.** Let  $\varepsilon_n \rightarrow 0, n \in N (n \rightarrow \infty), A_n := A_{\varepsilon_n}, g_n := g_{\varepsilon_n}$  and  $u_n := u_{\varepsilon_n}$  be a sequence so that  $u_n \in \mathcal{E}_{\varepsilon_n}$ . Since  $f$  is a bounded function,  $(f(u_n))_n$  is a bounded sequence in  $L^2(\Omega)$ . Thus, there exist  $f_0 \in L^2(\Omega)$  and an infinite subset  $N_4 \subset N$  so that

$$f(u_n) \rightharpoonup f_0 \quad \text{in } L^2(\Omega), \quad n \in N_4.$$

On the other hand, since  $(Pg_n)_n$  is a bounded sequence, there exist  $g_0 \in L^2(\Omega)$  and an infinite subset  $N_5 \subset N_4$  so that

$$Pg_n \rightharpoonup g_0 \quad \text{in } L^2(\Omega), \quad n \in N_5.$$

By Theorem 1.1,

$$\|A_n^{-1}(-f(u_n) + g_n) - A_0^{-1}P(-f(u_n) + g_n)\|_{L^2(\Omega)} \rightarrow 0, \quad n \in N_5.$$

Also note that

$$A_0^{-1}P(-f(u_n) + g_n) \rightharpoonup A_0^{-1}P(-f_0 + g_0) \quad \text{in } L^2(\Omega), \quad n \in N_5.$$

Now, since  $A_0^{-1}$  is a compact operator,  $u_n^s := A_0^{-1}P(-f(u_n) + g_n)$  has a convergent subsequence. Thus, there exists  $u_0 \in L^2(I)$ , so that

$$u_n^s \rightarrow u_0 \quad \text{in } L^2(\Omega), \quad n \in N_6,$$

where  $N_6$  is an infinite subset of  $N_5$ . Consequently,

$$u_n \rightarrow u_0 \quad \text{in } L^2(\Omega), \quad n \in N_6,$$

and  $u_0 = A_0^{-1}P(-f_0 + g_0)$ . To finish the proof, since  $f$  is a Lipschitz function,  $f(u_n) \rightarrow f(u_0)$ , in  $L^2(\Omega), n \in N_6$ , and so we conclude that  $f_0 = f(u_0)$ .

Observe that Lemma 5.1 was not necessary to prove Theorem 1.3. To prove Theorem 1.4, besides Lemma 5.1, we will need the following result.

**LEMMA 5.2.** *Suppose that there exists  $c > 0$  so that  $\|g_\varepsilon\| < c$ , for all small enough  $\varepsilon > 0$ . Then, there are two positive constants  $\varepsilon_3$  and  $M$ , such that*

$$\|u_\varepsilon\|_{H^1(\Omega)} \leq M, \quad \forall u_\varepsilon \in \mathcal{E}_\varepsilon, \quad 0 < \varepsilon < \varepsilon_3.$$

**PROOF.** Due to (5.1), given  $\eta > 0$ , there exists  $C_\eta > 0$  so that

$$\begin{aligned} & \int_{\Omega} \left( |u'_\varepsilon + \langle \nabla_y u_\varepsilon, Ry \rangle (\tau + \alpha')|^2 + \frac{\beta_\varepsilon}{\varepsilon^2} |\nabla_y u_\varepsilon|^2 + |u_\varepsilon|^2 \right) ds dy \\ &= \int_{\Omega} (A_\varepsilon u_\varepsilon) u_\varepsilon ds dy \\ &= \int_{\Omega} (-f(u_\varepsilon) + g_\varepsilon) u_\varepsilon ds dy \\ &= \int_{\Omega} -f(u_\varepsilon) u_\varepsilon ds dy + \int_{\Omega} g_\varepsilon u_\varepsilon ds dy \\ &\leq \int_{\Omega} (\eta u_\varepsilon^2 + C_\eta) ds dy + \frac{1}{2\eta} \int_{\Omega} g_\varepsilon^2 ds dy + \frac{\eta}{2} \int_{\Omega} u_\varepsilon^2 ds dy \\ &\leq \frac{3}{2} \eta \|u_\varepsilon\|_{H^1(\Omega)}^2 + \tilde{S}, \end{aligned}$$

where  $\tilde{S} := C_\eta \text{mes}(\Omega) + c^2/(2\eta)$ . Recalling (2.4), we also have

$$E_1 \int_{\Omega} |\nabla_y u_\varepsilon|^2 ds dy \leq \int_{\Omega} \frac{\beta_\varepsilon}{\varepsilon^2} |\nabla_y u_\varepsilon|^2 ds dy \leq \frac{3}{2} \eta \|u_\varepsilon\|_{H^1(\Omega)}^2 + \tilde{S}.$$

With the above inequalities, we obtain

$$\begin{aligned} E_1 \|u_\varepsilon\|_{H^1(\Omega)}^2 &= E_1 \int_{\Omega} (|u'_\varepsilon|^2 + |\nabla_y u_\varepsilon|^2 + |u_\varepsilon|^2) ds dy \\ &\leq E_1 \left[ \left( \int_{\Omega} |u'_\varepsilon + \langle \nabla_y u_\varepsilon, Ry \rangle (\tau + \alpha')|^2 ds dy \right)^{1/2} \right. \\ &\quad \left. + \left( \int_{\Omega} |\langle \nabla_y u_\varepsilon, Ry \rangle (\tau + \alpha')|^2 ds dy \right)^{1/2} \right]^2 \\ &\quad + E_1 \int_{\Omega} (|\nabla_y u_\varepsilon|^2 + |u_\varepsilon|^2) ds dy \\ &\leq E_1 \left[ \left( \int_{\Omega} |u'_\varepsilon + \langle \nabla_y u_\varepsilon, Ry \rangle (\tau + \alpha')|^2 ds dy \right)^{1/2} \right. \\ &\quad \left. + C_2 \left( \int_{\Omega} |\nabla_y u_\varepsilon|^2 ds dy \right)^{1/2} \right]^2 \\ &\quad + E_1 \int_{\Omega} (|\nabla_y u_\varepsilon|^2 + |u_\varepsilon|^2) ds dy \\ &\leq (3E_1 + 2E_1 C_2 + E_1 C_2^2) \left( \frac{3}{2} \eta \|u_\varepsilon\|_{H^1(\Omega)}^2 + \tilde{S} \right). \end{aligned}$$

Recall that  $C_2 = \sup_{(s,y) \in \Omega} \{ \|Ry\| (\tau + \alpha')(s) \}$ . Now it is sufficient to take  $\eta > 0$  so that

$$\frac{3}{2} \eta (3E_1 + 2E_1 C_2 + E_1 C_2^2) < E_1,$$

and the proof is complete. □

**PROOF OF THEOREM 1.4.** Let  $\varepsilon_n \rightarrow 0, n \in N (n \rightarrow \infty), g_n := g_{\varepsilon_n}, A_n := A_{\varepsilon_n}$  and  $u_n := u_{\varepsilon_n}$  be a sequence so that  $u_n \in \mathcal{E}_{\varepsilon_n}$ . Lemmas 5.1 and 5.2 ensure that  $(f(u_n))_n$  is a bounded sequence in  $L^2(\Omega)$ . Thus, there exist  $f_0 \in L^2(\Omega)$  and an infinite subset  $N_7 \subset N$  with

$$f(u_n) \rightharpoonup f_0 \text{ in } L^2(\Omega), n \in N_7.$$

Further, since  $(g_n)_n$  is a bounded sequence, there exist  $g_0 \in L^2(\Omega)$  and an infinite subset  $N_8 \subset N_7$  so that

$$g_n \rightarrow g_0 \text{ in } L^2(\Omega), n \in N_8.$$

By Theorem 1.1,

$$\|A_n^{-1}(-f(u_n) + g_n) - A_0^{-1}P(-f(u_n) + g_n)\|_{L^2(\Omega)} \rightarrow 0, n \in N_8.$$

Also note that

$$A_0^{-1}(-f(u_n) + g_n) \rightharpoonup A_0^{-1}P(-f_0 + g_0) \text{ in } L^2(\Omega), n \in N_8.$$

Since  $A_0^{-1}$  is a compact operator,  $u_n^s := A_0^{-1}P(-f(u_n) + g_n)$  has a convergent subsequence. Thus, there exists  $u_0 \in L^2(I)$  so that

$$u_n^s \rightarrow u_0 \text{ in } L^2(\Omega), n \in N_9,$$

where  $N_9$  is an infinite subset of  $N_8$ . Consequently,

$$u_n \rightarrow u_0 \text{ in } L^2(\Omega), n \in N_9,$$

and  $u_0 = A_0^{-1}P(-f_0 + g_0)$ . Lemma 5.2 also ensures that

$$u'_n \rightharpoonup u'_0 \text{ and } \nabla_y u_n \rightarrow 0 \text{ in } L^2(\Omega), n \in N_9.$$

Now, since the norms  $\|u\|_{H^1(\Omega)}$  and  $\|u\|_{H^1_\varepsilon(\Omega)}$  are equivalent, to prove the convergence in (1.17) it is enough to check that  $\|u_\varepsilon\|_{H^1_\varepsilon(\Omega)} \rightarrow \|u_0\|_{H^1(\Omega)}$ , which follows from the following estimates.

$$\begin{aligned} \|u_0\|_{H^1(\Omega)}^2 &\leq \liminf_{n \in N_9} \int_{\Omega} (|u'_n + \langle \nabla_y u_n, Ry \rangle (\tau + \alpha')|^2 + \beta_{\varepsilon_n} |\nabla_y u_n|^2 + |u_n|^2) ds dy \\ &\leq \limsup_{n \in N_9} \int_{\Omega} (|u'_n + \langle \nabla_y u_n, Ry \rangle (\tau + \alpha')|^2 + \beta_{\varepsilon_n} |\nabla_y u_n|^2 + |u_n|^2) ds dy \\ &\leq \limsup_{n \in N_9} \int_{\Omega} \left( |u'_n + \langle \nabla_y u_n, Ry \rangle (\tau + \alpha')|^2 + \frac{\beta_{\varepsilon_n}}{\varepsilon_n^2} |\nabla_y u_n|^2 + |u_n|^2 \right) ds dy \\ &= \limsup_{n \in N_9} \int_{\Omega} (A_n u_n) u_n ds dy \\ &= \limsup_{n \in N_9} \int_{\Omega} (-f(u_n) + g_n) u_n ds dy \\ &= \int_{\Omega} (-f_0 + g_0) u_0 ds dy \\ &= \int_{\Omega} (A_0 u_0) u_0 ds \\ &= \|u_0\|_{H^1(\Omega)}^2. \end{aligned}$$



Thus,  $\|u_n - u_0\|_{H^1(\Omega)} \rightarrow 0$ . This limit, combined with Lemma 5.1, implies that  $f(u_n) \rightarrow f(u_0)$ ,  $n \in N_9$ . Therefore,  $u_0 = A_0^{-1}P(-f(u_0) + g_0)$ . □

### 6. Regions bounded by a smooth function

Let  $g : [a, b] \rightarrow \mathbb{R}$ ,  $-\infty < a < b < +\infty$ , be a function of class  $C^3$  satisfying,

$$\left| \frac{g'(s)}{g(s)} \right| \leq X_1, \quad X_2 \leq g(s) \leq X_1, \quad \frac{1}{g^2(s)} - \left( \frac{g'(s)}{g(s)} \right)^2 \geq \frac{1}{2X_1^2}, \quad \forall s \in [a, b],$$

where  $X_1, X_2$  are positive numbers.

Recall that we use the notation  $J = (a, b)$ . As mentioned in the Introduction, we consider the domain

$$Q_\varepsilon = \{(s, y) \in J \times \mathbb{R} : 0 < y < \varepsilon g(s)\}$$

and the Equation (1.19), that is,

$$\begin{aligned} -\Delta u + f(u) &= 0 \quad \text{in } Q_\varepsilon, \\ \frac{\partial u}{\partial \nu_\varepsilon} &= 0 \quad \text{on } \partial Q_\varepsilon, \end{aligned}$$

where  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a nonlinear  $C^2$ -function satisfying the conditions (1.14)–(1.16).

The linear term in Equation (1.19) is described by the quadratic form

$$\hat{m}_\varepsilon(u) = \int_{Q_\varepsilon} |\nabla u|^2 dx, \quad u \in H^1(Q_\varepsilon). \tag{6.1}$$

By performing the change of variables

$$\begin{aligned} \tilde{F}_\varepsilon : J \times (0, 1) &\rightarrow Q_\varepsilon \\ (s, y) &\mapsto (s, \varepsilon y g(s)), \end{aligned}$$

(6.1) becomes

$$\tilde{m}_\varepsilon(u) = \int_Q \left[ \frac{1}{g} (gu' - yg'u_y)^2 + \frac{1}{\varepsilon^2 g} u_y^2 \right] ds dy, \tag{6.2}$$

where  $\text{dom } \tilde{m}_\varepsilon = H^1(Q)$  and  $Q = J \times (0, 1)$ . Further,  $\text{dom } \tilde{m}_\varepsilon$  is a subspace of the Hilbert space  $L^2(Q, g)$ . The details of this change of variables can be found in [9].

Now, we propose an additional change of variables in order to work in the Hilbert space  $L^2(Q)$ : that is, consider the unitary operator

$$\begin{aligned} V_\varepsilon : L^2(Q, g) &\rightarrow L^2(Q) \\ v &\mapsto g^{1/2}v, \end{aligned}$$

so that the quadratic form (6.2) becomes

$$m_\varepsilon(u) = \int_Q \left[ \left( u' - \frac{g'}{2g}u \right)^2 + \frac{1}{\varepsilon^2 g^2} u_y^2 \right] ds dy, \tag{6.3}$$

and now  $\text{dom } m_\varepsilon = H^1(Q)$  as a subspace of  $L^2(Q)$ .

Recall that  $M_\varepsilon$  is the self-adjoint operator associated with the quadratic form (6.3) and consider Equation (1.20), whose set of solutions is  $\mathcal{E}_\varepsilon^g$ . Recall that  $\tilde{\mathcal{J}} = \{w1 : w \in L^2(J)\}$  and that  $M$  is the self-adjoint operator associated with the quadratic form  $m$  given by (1.21).

The proof of the next theorem follows the same strategy employed in the proof of Theorem 1.1: that is, one must identify the correct subspaces and demonstrate the required estimates for the application of the technique of [8].

**THEOREM 6.1.** *There are two positive numbers  $N$  and  $\varepsilon_4$ , so that, for  $0 < \varepsilon < \varepsilon_4$ ,*

$$\|M_\varepsilon^{-1} - M^{-1}\tilde{P}\|_{L^2(Q)} \leq N\varepsilon,$$

where  $\tilde{P}$  denotes the orthogonal projection onto the subspace  $\tilde{\mathcal{J}}$ .

**PROOF.** If  $u \in \text{dom } m_\varepsilon$ , write  $u(s, y) = w(s) + v(s, y)$ , where  $w \in H^1(J)$  and  $v \in \tilde{\mathcal{J}}^\perp \cap H^1(Q)$ . Thus,

$$m_\varepsilon(u) = m_\varepsilon(w) + 2m_\varepsilon(w, v) + m_\varepsilon(v),$$

where  $m_\varepsilon(u_1, u_2)$  is the bilinear form associated with quadratic form  $m_\varepsilon(u)$ .

Again, we are going to show that the conditions (3.2), (3.3), (3.4) and (3.5) of [8, Section 3] are satisfied, and so the theorem follows. For this, just note that

$$\begin{aligned} m_\varepsilon(w) &= m(w), \quad \forall w \in H^1(J), \\ m_\varepsilon(w, v) &= 0, \quad \forall w \in H^1(J) \quad \text{and} \quad v \in \tilde{\mathcal{J}}^\perp \cap H^1(Q), \\ m_\varepsilon(v) &\geq \frac{\pi^2}{\varepsilon^2 X_1^2} \int_Q |v|^2 \, ds \, dy, \quad \forall v \in \tilde{\mathcal{J}}^\perp \cap H^1(Q), \end{aligned}$$

and the proof is complete. □

The following lemma follows from [9, Theorem 2.4].

**LEMMA 6.2.** *There are two positive numbers  $T$  and  $\varepsilon_5 > 0$ , so that, for  $0 < \varepsilon < \varepsilon_5$ ,*

$$\|u_\varepsilon\|_{H^2(Q)} \leq T \quad \forall u_\varepsilon \in \mathcal{E}_\varepsilon^g.$$

**PROOF OF THEOREM 1.5.** Let  $\varepsilon_n \rightarrow 0$ ,  $n \in N$  ( $n \rightarrow \infty$ ),  $M_n := M_{\varepsilon_n}$  and  $u_n := u_{\varepsilon_n}$  be a sequence so that  $u_n \in \mathcal{E}_{\varepsilon_n}^g$ . Lemmas 6.2 and 5.1 (see Section 5 for Lemma 5.1) ensure that  $(f(u_n))_n$  is a bounded sequence in  $L^2(Q)$ . Thus, there exist  $f_0 \in L^2(Q)$  and an infinite subset  $N_{10} \subset N$  so that

$$f(u_n) \rightharpoonup f_0 \quad \text{in } L^2(Q), \quad n \in N_{10}.$$

By Theorem 6.1,

$$\|M_n^{-1}(-f(u_n)) - M^{-1}\tilde{P}(-f(u_n))\|_{L^2(Q)} \rightarrow 0, \quad n \in N_{10},$$

and also note that

$$M_0^{-1}(-f(u_n)) \rightharpoonup M^{-1}\tilde{P}(-f_0) \quad \text{in } L^2(Q), \quad n \in N_{10}.$$

Since  $M^{-1}$  is a compact operator (in fact,  $M^{-1}$  is a norm limit of compact operators),  $u_n^s := M^{-1}\tilde{P}(-f(u_n))$  has a convergent subsequence. Thus, there exists  $u_0 \in L^2(J)$  so that

$$\|u_n^s - u_0\|_{L^2(Q)} \rightarrow 0, \quad n \in N_{11},$$

where  $N_{11}$  is an infinite subset of  $N_{10}$ . Consequently,  $u_0 = M^{-1}\tilde{P}(-f_0)$ . Since the immersion  $H^2(Q) \rightarrow L^2(Q)$  is compact, Lemma 6.2 ensures that

$$\|u_n - u_0\|_{H^1(Q)} \rightarrow 0, \quad n \in N_{12},$$

where  $N_{12}$  is an infinite subset of  $N_{11}$ . This limit, combined with Lemma 5.1, implies that  $f(u_n) \rightarrow f(u_0)$ ,  $n \in N_{12}$ . Thus,  $u_0 = M^{-1}\tilde{P}(-f(u_0))$ .

### Appendix

In this appendix we present some details of the limiting behaviour of the quadratic form (1.6), as  $\varepsilon \rightarrow 0$ . Recall that it is given by

$$\int_{I \times S} \left( \frac{1}{\gamma_\varepsilon} |u'|^2 + \frac{\gamma_\varepsilon}{\varepsilon^2} (|\nabla_y u|^2 - \lambda_0 |u|^2) \right) ds dy,$$

where  $\gamma_\varepsilon(s, y) = 1 - (\varepsilon y_1)/a$ ,  $\lambda_0$  is the first eigenvalue of the Dirichlet Laplacian in  $H_0^1(S)$ , and recall that  $u_0 = u_0(y)$  is the corresponding (positive) normalized eigenfunction.

Consider the subspace  $\mathcal{J} = \{wu_0 : w \in H_0^1(I)\}$  and the quadratic form

$$\int_S \left( \frac{\gamma_\varepsilon}{\varepsilon^2} (|\nabla_y u|^2 - \lambda_0 |u|^2) \right) ds dy$$

restricted to  $\mathcal{J}$ . For this, integration by parts shows that

$$\begin{aligned} & \int_S \frac{(1 - (\varepsilon y_1)/a)}{\varepsilon^2} (|\nabla_y u_0|^2 - \lambda_0 u_0^2) dy \\ &= \int_S \frac{-y_1/a}{\varepsilon} (|\nabla_y u_0|^2 - \lambda_0 u_0^2) dy \\ &= \int_S \left[ \frac{\partial}{\partial y_1} \left( \frac{y_1}{a\varepsilon} \frac{\partial u_0}{\partial y_1} \right) u_0 + \frac{\partial}{\partial y_2} \left( \frac{y_1}{a\varepsilon} \frac{\partial u_0}{\partial y_2} \right) u_0 + \frac{y_1 \lambda_0}{a\varepsilon} u_0^2 \right] dy \\ &= \int_S \left( \frac{1}{a\varepsilon} \frac{\partial u_0}{\partial y_1} u_0 + \frac{y_1}{a\varepsilon} \Delta_y u_0 + \frac{y_1}{a\varepsilon} \lambda_0 u_0^2 \right) dy \\ &= \int_S \left( \frac{1}{a\varepsilon} \frac{\partial u_0}{\partial y_1} u_0 \right) dy = 0. \end{aligned}$$

Thus, since  $\gamma_\varepsilon(s, y) \rightarrow 1$  uniformly, as  $\varepsilon \rightarrow 0$ , we have, for the full quadratic form,

$$\begin{aligned} & \int_{I \times S} \left( \frac{1}{\gamma_\varepsilon} |w'u_0|^2 + |w|^2 \frac{\gamma_\varepsilon}{\varepsilon^2} |\nabla_y u_0|^2 - \lambda_0 |w|^2 |u_0|^2 \right) ds dy \\ &= \int_{I \times S} \frac{1}{\gamma_\varepsilon} |w'u_0|^2 ds dy \longrightarrow \int_I |w'|^2 ds. \end{aligned}$$

### Acknowledgements

The authors would like to thank Dr. Ricardo P. Silva for useful discussions. We acknowledge the good work done by the referee who encountered additional difficulties due to a typographical error in an important reference.

### References

- [1] J. M. Arrieta and A. N. Carvalho, ‘Spectral convergence and nonlinear dynamics of reaction–diffusion equations under perturbations of the domain’, *J. Differential Equations* **199** (2004), 143–178.
- [2] G. Bouchitté, M. L. Mascarenhas and L. Trabucho, ‘On the curvatures and torsion effects in one-dimensional waveguides’, *ESAIM Control Optim. Calc. Var.* **13** (2007), 793–808.
- [3] G. Bouchitté, M. L. Mascarenhas and L. Trabucho, ‘Thin waveguides with Robin boundary conditions’, *J. Math. Phys.* **53** (2012), 123517, 24 pages.
- [4] A. N. Carvalho and S. Piskarev, ‘A general approximation scheme for attractors of abstract parabolic problems’, *Numer. Funct. Anal. Optim.* **27**(7-8) (2006), 785–829.
- [5] T. Elsken, ‘Limiting behavior of attractors for systems on thin domains’, *Hiroshima Math. J.* **32** (2002), 389–415.
- [6] T. Elsken, ‘A reaction–diffusion equation on a net-shaped thin domain’, *Studia Math.* **165** (2004), 159–199.
- [7] T. Elsken, ‘Continuity of attractors for net-shaped thin domain’, *Topol. Methods Nonlinear Anal.* **26** (2005), 315–354.
- [8] L. Friedlander and M. Solomyak, ‘On the spectrum of the Dirichlet Laplacian in a narrow infinite strip’, in: *Spectral Theory of Differential Operators*, American Mathematical Society Translation Series 2, 225 (American Mathematical Society, Providence, RI, 2008), 103–116.
- [9] K. Hale and G. Raugel, ‘Reaction-diffusion equation on thin domains’, *J. Math. Pures Appl.* **71** (1992), 33–95.
- [10] W. Klingenberg, *A Course in Differential Geometry* (Springer, New York, 1978).
- [11] C. R. de Oliveira, ‘Quantum singular operator limits of thin Dirichlet tubes via  $\Gamma$ -convergence’, *Rep. Math. Phys.* **67** (2011), 1–32.
- [12] C. R. de Oliveira and A. F. Rossini, ‘Effective operators for Robin Laplacian in thin two- and three-dimensional curved waveguides’, Preprint.
- [13] C. R. de Oliveira and A. A. Verri, ‘On the spectrum and weakly effective operator for Dirichlet Laplacian in thin deformed tubes’, *J. Math. Anal. Appl.* **381** (2011), 454–468.
- [14] C. R. de Oliveira and A. A. Verri, ‘Norm resolvent convergence of Dirichlet Laplacian in unbounded thin waveguides’, *Bull. Braz. Math. Soc. (N.S.)* **46** (2015), 139–158.
- [15] M. Prizzi, ‘A remark on reaction–diffusion equations in unbounded domains’, *Discrete Contin. Dyn. Syst.* **9** (2003), 281–286.
- [16] M. Prizzi and K. P. Rybakowski, ‘The effect of domain squeezing upon the dynamics of reaction diffusion equations’, *J. Differential Equations* **173** (2001), 271–320.
- [17] A. M. Rekaló, ‘Asymptotic behavior of solutions of nonlinear parabolic equations on two-layer thin domains’, *Nonlinear Anal.* **52** (2003), 1393–1410.
- [18] R. P. Silva, ‘A note on resolvent convergence on a thin domain’, *Bull. Aust. Math. Soc.* **89** (2014), 141–148.

CÉSAR R. DE OLIVEIRA, Departamento de Matemática, UFSCar,  
São Carlos, SP 13560-970, Brazil  
e-mail: [oliveira@dm.ufscar.br](mailto:oliveira@dm.ufscar.br)

ALESSANDRA A. VERRI, Departamento de Matemática, UFSCar,  
São Carlos, SP 13560-970, Brazil  
e-mail: [alessandraverri@dm.ufscar.br](mailto:alessandraverri@dm.ufscar.br)