

- resistant *Staphylococcus aureus* for the 1990s and effective control measures. *Am J Med.* 1991;91 (suppl 3B):221S-225S.
2. Schaberg DR, Culver DH, Gaynes RI? Major trends in the microbial etiology of nosocomial infection. *Am J Med.* 1991;91:72S-75s.
 3. Gelmi M, Forest I, Ravizzola G, et al. Antibiotic resistances and plasmids in *Staphylococcus aureus* from Italian hospitals. *J Med Microbiol.* 1987;23:111-118.
 4. Pavillard P, Harvey K, Douglas D, et al. Epidemic of hospital-acquired infection due to methicillin-resistant *Staphylococcus aureus* in major Victorian hospitals. *Med J Aust.* 1982;1:451-454.
 5. Bradley JM, Noone P, Townsend DE, Grubb WB. Methicillin-resistant *Staphylococcus aureus* in a London hospital. *Lancet.* 1985;i:1493-1495.
 6. Wey SB, Cardo DM, Walker E, Carratu FP, Saes AC. Distribution and analysis of 8,268 nosocomial infections at the Hospital Sao Paulo: 1985 to 1989. *Rev Hosp Sao Paulo.* 1990;1:169-174.
 7. Rodrigues JN, Amaral JLG, Pignatari A, et al. Molecular epidemiology and antimicrobial susceptibility testing of quinolone-resistant *Staphylococcus aureus* strains isolated in Brazil. *Diagn Microbiol Infect Dis.* 1993;16:9-16.
 8. Thompson RL, Cabezudo IC, Wenzel RI? Epidemiology of nosocomial infections caused by methicillin-resistant *Staphylococcus aureus*. *Ann Intern Med.* 1982;97:309-317.
 9. Sesso R, Draibe S, Castelo A, et al. *Staphylococcus aureus* skin carriage and development of peritonitis in patients on continuous ambulatory peritoneal dialysis. *Clin Nephrol.* 1989;31:264-268.
 10. National Committee for Clinical Laboratory Standards. *Approved Standard, M7-A2: Standard Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically.* 2nd ed. Villanova, PA NCCLS; 1990.
 11. Nahaie MR, Goodfellow M, Harwood CR. A rapid screening procedure for staphylococcal plasmids. *J Microbiol Methods.* 1984;2:73-81.
 12. Finch RG, Hawkey PM, Townner KJ, eds. RP-59500, a semisynthetic injectable streptogramin antibiotic. *J Antimicrob Chemother.* 1992;30(suppl A): 1-131.
 13. Haley RW, Hightower AW, Khabbaz RF, et al. The emergence of methicillin-resistant *Staphylococcus aureus* infections in United States hospitals: possible role of the house staff-patient transfer circuit. *Ann Intern Med.* 1982;97:297-308.
 14. Lyon BR, Iuorio JL, May JW, Skurray RA. Molecular epidemiology of multiresistant *Staphylococcus aureus* in Australian hospitals. *J Med Microbiol.* 1984;17:79-89.
 15. Townsend DE, Ashdown N, Bradley JM, Pearman JW, Grubb WB. Australian methicillin-resistant *Staphylococcus aureus* in a London hospital. *Med J Aust.* 1984;141:339-340.
 16. Zuccarelli AJ, Roy I, Harding GP, Couperus JJ. Diversity and stability of restriction enzyme profiles of plasmid DNA from methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol.* 1990;28:97-102.
 17. Doebbeling BN, Pfaller MA, Hollis RJ, et al. Restriction endonuclease analysis of *Staphylococcus aureus* plasmid DNA from three continents. *Eur J Clin Microbiol Infect Dis.* 1992;11:4-8.

Reintroduction of Discontinued TB Drugs to Combat Epidemic

by Gina Pugliese, RN, MS
Medical News Editor

Mary Pendergast, senior advisor to the commissioner of the Food and Drug Administration (FDA), reported at the recent World Congress on Tuberculosis in Bethesda, Maryland, that the FDA is concerned with the need for more effective drugs for treating tuberculosis because many of the existing antituberculosis drugs are either unavailable or in insufficient supply. She explained that the FDA is currently tracking down sources for existing antitu-

berculosis drugs throughout the world, as well as urging United States drug manufacturers to keep antituberculosis drugs on the market, bring back discontinued drugs, and seek approval for new ones. The FDA is also planning to expedite the review of antituberculosis drugs, biologics, and devices.

Emergency supplies of streptomycin sulfate are being distributed through the Centers for Disease Control and Prevention (CDC) as an investigational drug until drug production can resume in the United States. Two generic drugs were approved last year for

tuberculosis treatment, a form of amikacin and pyrazinamide. Additionally, the only injectable form of isoniazid, Nydrazid, has just been reintroduced at the request of the FDA and CDC. It will be useful for patients who are unable to take pills.

The FDA's willingness to accept foreign data for approval of drugs from outside the United States is evidenced by the drug Rifater. This agent is a combination of rifampin, isoniazid, and pyrazinamide and would make compliance with therapy much easier.