



COMPOSITIO MATHEMATICA

On two examples by Iyama and Yoshino

Bernhard Keller, Daniel Murfet and Michel Van den Bergh

Compositio Math. **147** (2011), 591–612.

[doi:10.1112/S0010437X10004902](https://doi.org/10.1112/S0010437X10004902)



FOUNDATION
COMPOSITIO
MATHEMATICA

*The London
Mathematical
Society*





On two examples by Iyama and Yoshino

Bernhard Keller, Daniel Murfet and Michel Van den Bergh

ABSTRACT

In a recent paper, Iyama and Yoshino considered two interesting examples of isolated singularities over which it is possible to classify the indecomposable maximal Cohen–Macaulay modules in terms of linear algebra data. In this paper, we present two new approaches to these examples. In the first approach we give a relation with cluster categories. In the second approach we use Orlov’s result on the graded singularity category.

Contents

1	Introduction	591
2	Notation and conventions	593
3	First approach to the second example	594
4	The singularity category of graded Gorenstein rings	599
5	The Iyama–Yoshino examples (again)	602
6	A remark on gradability of rigid modules	603
	Acknowledgements	604
	Appendix A Generators of singularity categories	605
	References	610

1. Introduction

Throughout, k is a field. In [OY08], Iyama and Yoshino considered the following two settings.

Example 1.1. Let $S = k[[x_1, x_2, x_3]]$ and let $C_3 = \langle \sigma \rangle$ be the cyclic group of three elements. Consider the action of C_3 on S via $\sigma x_i = \omega x_i$, where $\omega^3 = 1$, $\omega \neq 1$. Put $R = S^{C_3}$.

Example 1.2. Let $S = k[[x_1, x_2, x_3, x_4]]$ and let $C_2 = \langle \sigma \rangle$ be the cyclic group of two elements. Consider the action of C_2 on S via $\sigma x_i = -x_i$. Put $R = S^{C_2}$.

In both examples, Iyama and Yoshino reduced the classification of maximal Cohen–Macaulay modules over R to the representation theory of certain generalized Kronecker quivers. They used this to classify the rigid Cohen–Macaulay modules over R . As predicted by deformation theory, the latter are described by discrete data.

The explicit description of the stable category of maximal Cohen–Macaulay modules over a commutative Gorenstein ring (also known as the singularity category [Buc87, BEH87, Orl93]) is a problem that has received much attention over the years. This appears to be in general a difficult problem and perhaps the best one can hope for is a reduction to linear algebra or, in other words, the representation theory of quivers. This is precisely what Iyama and Yoshino have accomplished.

Received 9 April 2008, accepted in final form 16 February 2010, published online 9 February 2011.

2000 Mathematics Subject Classification 13C14, 16E65 (primary).

Keywords: singularity category, maximal Cohen–Macaulay modules.

This journal is © Foundation Compositio Mathematica 2011.

The proofs of Iyama and Yoshino are based on the machinery of mutation in triangulated categories, a general theory developed by them. In the current paper we present two alternative approaches to the examples. Hopefully the additional insight obtained in this way may be useful elsewhere.

Our first approach applies to Example 1.2 and is inspired by the treatment in [KR08] of Example 1.1, where the authors used the fact that in this case the stable category $\underline{\text{MCM}}(R)$ of maximal Cohen–Macaulay R -modules is a 2-Calabi–Yau category which has a cluster tilting object whose endomorphism ring is the path algebra kQ_3 of the Kronecker quiver with three arrows. Then they invoked their acyclicity result (slightly specialized).

THEOREM 1.3 [KR08, §1, Theorem]. *Assume that \mathcal{T} is a k -linear algebraic Krull–Schmidt 2-Calabi–Yau category with a cluster tilting object T such that $A = \text{End}(T)$ is hereditary. Then there is an exact equivalence between \mathcal{T} and the orbit category $D^b(\text{mod}(A))/(\tau[-1])$.*

From this result, they obtained immediately that $\underline{\text{MCM}}(R)$ is the orbit category $D^b(\text{mod}(kQ_3))/(\tau[-1])$. This gives a very satisfactory description of $\underline{\text{MCM}}(R)$ and implies in particular the results by Iyama and Yoshino.

In the first part of this paper we show that Example 1.2 is amenable to a similar approach. Iyama and Yoshino proved that $\underline{\text{MCM}}(R)$ is a 3-Calabi–Yau category with a 3-cluster tilting object T such that $\text{End}(T) = k$ [OY08, Theorem 9.3]. We show that under these circumstances there is an analogue of the acyclicity result of the first author and Reiten.

THEOREM 1.4 (See §3.4). *Assume that \mathcal{T} is a k -linear algebraic Krull–Schmidt 3-Calabi–Yau category with a 3-cluster tilting object T such that $\text{End}(T) = k$. Then there is an exact equivalence of \mathcal{T} with the orbit category $D^b(\text{mod}(kQ_n))/(\tau^{1/2}[-1])$, $n = \dim \text{Ext}_{\mathcal{T}}^{-1}(T, T)$, where Q_n is the generalized Kronecker quiver with n arrows and $\tau^{1/2}$ is a natural square root of the Auslander–Reiten translate of $D^b(\text{mod}(kQ_n))$, which on the pre-projective/pre-injective component corresponds to ‘moving one place to the left’.*

In the second part of this paper, which is logically independent of the first, we give yet another approach to Examples 1.1 and 1.2 based on the following observation, which might have independent interest.

PROPOSITION 1.5 (See Proposition A.8). *Let $A = k + A_1 + A_2 + \dots$ be a finitely generated commutative graded Gorenstein k -algebra with an isolated singularity. Let \widehat{A} be the completion of A at $A_{\geq 1}$. Let $\underline{\text{MCM}}_{\text{gr}}(A)$ be the stable category of graded maximal Cohen–Macaulay A -modules. Then the obvious functor $\underline{\text{MCM}}_{\text{gr}}(A) \rightarrow \underline{\text{MCM}}(\widehat{A})$ induces an equivalence*

$$\underline{\text{MCM}}_{\text{gr}}(A)/(1) \cong \underline{\text{MCM}}(\widehat{A}), \tag{1.1}$$

where $M \mapsto M(1)$ is the shift functor for the grading.

In this proposition the quotient $\underline{\text{MCM}}_{\text{gr}}(A)/(1)$ has to be understood as the triangulated/Karoubian hull (as explained in [Kel05]) of the naive quotient of $\underline{\text{MCM}}_{\text{gr}}(A)$ by the shift functor $?(1)$. This result is similar in spirit to [AR90], which treats the finite representation type case. Note however that one of the main results in [AR90] is that in the finite representation type case every indecomposable maximal Cohen–Macaulay \widehat{A} -module is gradable. This does not seem to be a formal consequence of Proposition 1.5. It would be interesting to investigate this further.

In §6, we show that at least rigid Cohen–Macaulay modules are always gradable so they are automatically in the image of $\text{MCM}_{\text{gr}}(A)$. We expect this to be well known in some form but we have been unable to locate a reference.

Hence, in order to understand $\text{MCM}(\widehat{A})$ it is sufficient to understand $\text{MCM}_{\text{gr}}(A)$. The latter is the graded singularity category [Orl09a] of A and, by [Orl09a, Theorem 2.5], it is related to $D^b(\text{coh}(X))$, where $X = \text{Proj } A$.

In Examples 1.1 and 1.2, R is the completion of a graded ring A which is the Veronese of a polynomial ring. Hence, $\text{Proj } A$ is simply a projective space. Using Orlov’s results and the existence of exceptional collections on projective space, we get very quickly in Example 1.1

$$\underline{\text{MCM}}_{\text{gr}}(A) \cong D^b(\text{mod}(kQ_3))$$

and in Example 1.2

$$\underline{\text{MCM}}_{\text{gr}}(A) \cong D^b(\text{mod}(kQ_6))$$

(where here and below \cong actually stands for a quasi-equivalence between the underlying DG-categories). Finally, it suffices to observe that in Example 1.1 we have $?(-1) = \tau[-1]$ and in Example 1.2 we have $?(-1) = \tau^{1/2}[-1]$ (see §5 below).

Finally, we mention the following interesting side result.

PROPOSITION 1.6. *Let (R, m) be a Gorenstein local ‘G-ring’ (for example R may be essentially of finite type over a field) with an isolated singularity. Then the natural functor*

$$\widehat{R} \otimes_R ? : \underline{\text{MCM}}(R) \rightarrow \underline{\text{MCM}}(\widehat{R}) \tag{1.2}$$

is an equivalence up to direct summands. In particular, every maximal Cohen–Macaulay module over \widehat{R} is a direct summand of the completion of a maximal Cohen–Macaulay module over R .

The original proof (by the first and third authors) of this result was unnecessarily complicated. After the paper was put on the arXiv, Daniel Murfet (who has become the second author) informed us about the existence of a much nicer proof in the context of singularity categories (see Proposition A.1). The same argument also applies to Proposition 1.5. So, we dropped our original proofs and put the new argument in an appendix, to which we refer.

Meanwhile, Orlov [Orl04] has proved (independently and using different methods) a very general result, which implies in particular Proposition 1.6.

2. Notation and conventions

We hope that the notation is self explanatory but nevertheless we give it here. If R is a ring, then $\text{Mod}(R)$ and $\text{mod}(R)$ denote respectively the category of all left R -modules and the full subcategory of finitely generated R -modules. The derived category of all R -modules is denoted by $D(R)$. If R is graded, then we use $\text{Gr}(R)$ and $\text{gr}(R)$ for the category of graded left modules and its subcategory of finitely generated modules. The shift functor on $\text{Gr}(R)$ is denoted by $?(1)$. Explicitly, $M(1)_i = M_{i+1}$. If we want to refer to right modules, then we use R° instead of R . If X is a scheme, then $\text{Qch}(X)$ is the category of quasi-coherent \mathcal{O}_X -modules. If X is noetherian, then $\text{coh}(X)$ is the category of coherent \mathcal{O}_X -modules. We are generally very explicit about which categories we use. For example, we write $D^b(\text{mod}(R))$ rather than something like $D_f^b(R)$. If R is graded and M, N are graded R -modules, then $\text{Ext}_R^i(M, N)$ is the ungraded Ext between M and N . If we need Ext in the category of graded R -modules, then we write $\text{Ext}_{\text{Gr}(R)}^i(M, N)$.

3. First approach to the second example

3.1 Some preliminaries on tilting complexes

Let C, E be rings. We denote the unbounded derived category of right C -modules by $D(C^\circ)$. We let $\text{Eq}(D(C^\circ), D(E^\circ))$ be the set of triangle equivalences of $D(C^\circ) \rightarrow D(E^\circ)$ modulo natural isomorphisms. Define $\text{Tilt}(C, E)$ as the set of pairs (ϕ, T) , where T is a perfect complex generating $D(E^\circ)$ and ϕ is an isomorphism $C \rightarrow \text{RHom}_E(T)$. Associated to $(\phi, T) \in \text{Tilt}(C, E)$ there is a canonical equivalence $\theta : D(C^\circ) \rightarrow D(E^\circ)$ such that $\theta(C) = T$. It may be constructed either directly [Ric89] or using DG-algebras [Kel98]. The induced map

$$\text{Tilt}(C, E) \rightarrow \text{Eq}(D(C^\circ), D(E^\circ))$$

is obviously injective (as it is canonically split), but not known to be surjective. Below we will informally refer to the elements of $\text{Tilt}(C, E)$ as tilting complexes.

3.2 A square root of τ for a generalized Kronecker quiver

Let W be a finite-dimensional k -vector space and let C be the path algebra of the quiver¹

$$\begin{array}{ccc} & W & \\ \bullet & \longleftarrow & \bullet \\ 1 & & 2 \end{array} \tag{3.1}$$

Let E be the path algebra of the quiver

$$\begin{array}{ccc} & W^* & \\ \bullet & \longleftarrow & \bullet \\ 1 & & 2 \end{array}$$

which we think of as being obtained from (3.1) by ‘inverting the arrows’ and renumbering the vertices $(1, 2) \mapsto (2, 1)$.

Let P_i, I_i, S_i be respectively the projective, injective, and simple right C -modules corresponding to vertex i . For E , we use P'_i, I'_i, S'_i . Let $r_i : \text{mod}(C^\circ) \rightarrow \text{mod}(E^\circ)$ be the reflection functor at vertex i . Recall that if (U, V) is a representation of C , then $r_1(U, V)$ is given by (V, U') , where $U' = \ker(V \otimes W \rightarrow U)$ (taking into account the renumbered vertices).

The right derived functor Rr_1 of r_1 defines an equivalence $D(C^\circ) \rightarrow D(E^\circ)$. It is obtained from the tilting complex $S'_2[-1] \oplus P'_1$ [APR79]. One has (see [Gab80])

$$Rr_1 \circ Rr_1 = \tau_C, \tag{3.2}$$

where τ_C is the Auslander–Reiten translate on $D(C^\circ)$. Assume now that W is equipped with an isomorphism $\pi : W \rightarrow W^*$. Then π yields an equivalence $D(E^\circ) \cong D(C^\circ)$, which we denote by the same symbol. We use the same convention for the transpose isomorphism $\pi^* : W \rightarrow W^*$.

LEMMA 3.2.1. *We have $r_1 \circ \pi^{-1} = \pi^* \circ r_1$ as functors $D(C^\circ) \rightarrow D(C^\circ)$.*

Proof. Let (U, V) be a representation of C determined by a linear map $c : V \otimes W \rightarrow U$ and put $(V, U'') = (r_1 \circ \pi^{-1})(U, V)$. Then one checks that U'' is given by the exact sequence

$$0 \rightarrow U'' \rightarrow V \otimes W^* \xrightarrow{c \circ (\pi^{-1} \otimes \text{id})} U \rightarrow 0,$$

¹ We use the convention that multiplication in the path algebra is concatenation. So, representations correspond to right modules.

where the first non-trivial map induces the action $U'' \otimes W \rightarrow V$. Similarly, if we put $(V, U') = (\pi^* \circ r_1)(U, V)$, then one gets the same sequence

$$0 \rightarrow U' \rightarrow V \otimes W^* \xrightarrow{c\circ(\pi^{-1}\otimes\text{id})} U,$$

where the first non-trivial map again yields the action $U' \otimes W \rightarrow V$. Thus, we have $(V, U') = (V, U'')$. □

Below we put $a = \pi \circ Rr_1$.

LEMMA 3.2.2. *One has $(\pi^* \circ \pi^{-1}) \circ a^2 = \tau$. In particular, $\tau \cong a^2$ if and only if π is self-adjoint or anti-self-adjoint.*

Proof. This is a straightforward verification using Lemma 3.2.1 and (3.2). □

For use below, we record

$$\begin{aligned} aP_2 &= P_1, \\ aP_1 &= I_2[-1], \\ aI_2 &= I_1. \end{aligned}$$

3.3 A 3-Calabi–Yau category with a 3-cluster tilting object

We let the notation be as in the previous section.

Put $\mathcal{H} = D^b(\text{mod}(C^\circ))$, $\mathcal{D} = \mathcal{H}/a[-1]$. As \mathcal{H} is hereditary, we have

$$\text{Ind}(\mathcal{D}) = \text{Ind}(\mathcal{H})/a[-1].$$

Inspection reveals that

$$\text{Ind}(\mathcal{D}) = \text{Ind}(\mathcal{H}) \cup \{I_2[-1]\}. \tag{3.3}$$

LEMMA 3.3.1. *\mathcal{D} is 3-Calabi–Yau if and only if π is self-adjoint or anti-self-adjoint.*

Proof. Let S be the Serre functor for \mathcal{H} . Being canonical, S commutes with the auto-equivalence $a[-1]$. Hence, S induces an auto-equivalence on \mathcal{D} , which is easily seen to be the Serre functor of \mathcal{D} .

In \mathcal{D} , we have $S = \tau[1] = (\pi^* \circ \pi^{-1}) \circ a^2[1] = (\pi^* \circ \pi^{-1})[3]$. Thus, \mathcal{D} is 3-Calabi–Yau if and only if $\pi^* \circ \pi^{-1}$ is isomorphic to the identity functor. It is easy to see that this is the case if and only if $\pi^* \circ \pi^{-1} = \pm 1$ in $\text{End}_k(W)$. □

LEMMA 3.3.2. *The object P_1 in \mathcal{D} satisfies*

$$\text{Ext}_{\mathcal{D}}^i(P_1, P_1) = 0 \text{ for } i = 1, 2, \quad \text{Hom}_{\mathcal{D}}(P_1, P_1) = k, \quad \text{and} \quad \text{Ext}_{\mathcal{D}}^{-1}(P_1, P_1) = W.$$

Proof. For $N \in \text{Ind}(\mathcal{H}) \cup \{I_2[-1]\}$, one computes

$$\text{Hom}_{\mathcal{D}}(P_1, N) = \text{Hom}_{\mathcal{H}}(P_1, N). \tag{3.4}$$

Thus, we find that

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(P_1, P_1[-1]) &= \text{Hom}_{\mathcal{D}}(P_1, a^{-1}P_1) \\ &= \text{Hom}_{\mathcal{D}}(P_1, P_2) \\ &= W, \end{aligned}$$

$$\text{Hom}_{\mathcal{D}}(P_1, P_1) = k,$$

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(P_1, P_1[1]) &= \text{Hom}_{\mathcal{D}}(P_1, aP_1) \\ &= \text{Hom}_{\mathcal{D}}(P_1, I_2[-1]) \\ &= 0, \end{aligned}$$

and

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(P_1, P_1[2]) &= \text{Hom}_{\mathcal{D}}(P_1, aP_1[1]) \\ &= \text{Hom}_{\mathcal{D}}(P_1, I_2) \\ &= 0. \end{aligned} \quad \square$$

The following lemma is not used explicitly.

LEMMA 3.3.3. *The object P_1 in \mathcal{D} has the properties of a 3-cluster tilting object, i.e. if $\text{Ext}_{\mathcal{D}}^i(P_1, N) = 0$ for $i = 1, 2$, then N is a sum of copies of P_1 .*

Proof. Assume that $N \in \text{Ind}(\mathcal{H}) \cup \{I_2[-1]\}$ is such that $\text{Hom}_{\mathcal{D}}(P_1, N[1]) = \text{Hom}_{\mathcal{D}}(P_1, N[2]) = 0$. We have to prove that $N = P_1$.

We may rewrite

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(P_1, N[2]) &= \text{Hom}_{\mathcal{D}}(P_1[-1], N[1]) \\ &= \text{Hom}_{\mathcal{D}}(a^{-1}P_1, N[1]) \\ &= \text{Hom}_{\mathcal{D}}(P_2, N[1]). \end{aligned}$$

Thus, we find that $\text{Hom}_{\mathcal{D}}(P_1, aN) = \text{Hom}_{\mathcal{D}}(P_2, aN) = 0$. Hence, $aN \notin \text{Ind}(\mathcal{H})$. We deduce that $N \in \{P_1, I_2[-1]\}$.

But, if $N = I_2[-1]$, then

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(P_1, N[2]) &= \text{Hom}_{\mathcal{D}}(P_1, I_2[1]) \\ &= \text{Hom}_{\mathcal{D}}(P_1, aI_2) \\ &= \text{Hom}_{\mathcal{D}}(P_1, I_1) \\ &\neq 0. \end{aligned}$$

So, we are left with the possibility $N = P_1$, which finishes the proof. □

3.4 Proof of Theorem 1.4

Let \mathcal{T} be an algebraic Ext-finite Krull–Schmidt 3-Calabi–Yau category containing a 3-cluster tilting object T such that $\text{End}_{\mathcal{T}}(T) = k$.

LEMMA 3.4.1. *Let $N \in \mathcal{T}$. Then there exists a distinguished triangle in \mathcal{T} :*

$$T^a \rightarrow T^b \oplus T[-1]^c \rightarrow N[1] \rightarrow . \tag{3.5}$$

Proof. Let Y be defined (up to isomorphism) by the following distinguished triangle:²

$$Y \rightarrow T^{\text{Ext}_{\mathcal{T}}^1(T, N)} \oplus T[-1]^{\text{Ext}_{\mathcal{T}}^2(T, N)} \rightarrow N[1] \rightarrow .$$

A quick check reveals that $\text{Ext}_{\mathcal{T}}^1(T, Y) = \text{Ext}_{\mathcal{T}}^2(T, Y) = 0$. Hence, $Y = T^a$ for some a . □

² It would be more logical to write e.g. $\text{Ext}_{\mathcal{T}}^1(T, N) \otimes_k T$ for $T^{\text{Ext}_{\mathcal{T}}^1(T, N)}$, but this would take a lot more space.

We need to consider the special case $N = T[1]$. Then the distinguished triangle (3.5) (constructed as in the proof) has the form

$$T^{\text{Ext}_{\mathcal{T}}^{-1}(T, T)} \xrightarrow{\phi} T[-1] \xrightarrow{\alpha} T[2] \xrightarrow{\beta}, \tag{3.6}$$

where ϕ is the universal map (this follows from applying $\text{Hom}_{\mathcal{T}}(T, -)$). Since $\text{End}_{\mathcal{C}}(T[2]) = k$, it follows that α, β are determined up to a (the same) scalar.

This has a surprising consequence. Applying $\text{Hom}_{\mathcal{T}}(-, T)$ to the triangle (3.6), we find that $\text{Hom}_{\mathcal{T}}(\beta[-1], T)^{-1}$ defines an isomorphism

$$\pi : \text{Ext}_{\mathcal{T}}^{-1}(T, T) \rightarrow \text{Ext}_{\mathcal{T}}^{-1}(T, T)^*.$$

Thus, $W \stackrel{\text{def}}{=} \text{Ext}_{\mathcal{C}}^{-1}(T, T)$ comes equipped with an isomorphism $\pi : W \rightarrow W^*$, which is canonical up to a scalar. In other words, we are in the setting of §3.2 and we now use the notation introduced in §§3.2 and 3.3.

As a is obtained from the reflection in vertex 1, one verifies (see §3.2) that a is associated to the element of $\text{Tilt}(C, C)$ given by $(\theta, I_2[-1] \oplus P_1)$, where $\theta : C \rightarrow \text{End}_C(I_2[-1] \oplus P_1)$ is the composition

$$C = \begin{pmatrix} k & 0 \\ W & k \end{pmatrix} \xrightarrow{\pi} \begin{pmatrix} k & 0 \\ W^* & k \end{pmatrix} = \text{End}_C(I_2[-1] \oplus P_1). \tag{3.7}$$

Since the auto-equivalence a is a derived functor that commutes with coproducts, it is isomorphic to a derived tensor functor $- \overset{L}{\otimes}_C X$ for some $X \in D(C^e)$, by [Kel94, 6.4]. As a right C -module, we have $X \cong I_2[-1] \oplus P_1$.

Now we use the assumption that \mathcal{H} is algebraic and we proceed more or less as in the appendix to [KR08]. By [Kel94, Theorem 4.3], we may assume that \mathcal{T} is a strict (= closed under isomorphism) triangulated subcategory of a derived category $D(\mathcal{A})$ for some DG-category \mathcal{A} . We denote by ${}_{\mathcal{C}}\mathcal{T}$ the full subcategory of $D(C \otimes \mathcal{A})$ whose objects are differential graded $C \otimes \mathcal{A}$ -modules which are in \mathcal{T} when considered as \mathcal{A} -modules. Clearly, ${}_{\mathcal{C}}\mathcal{T}$ is triangulated. By [KR08, Lemma A.2.1(a)], T may be lifted to an object in ${}_{\mathcal{C}}\mathcal{T}$, which we also denote by T . Put $S = T \oplus T[-1]$.

LEMMA 3.4.2. *One has an isomorphism in ${}_{\mathcal{C}}\mathcal{T}$:*

$$X \overset{L}{\otimes}_B S \cong S[1].$$

Proof. As objects in \mathcal{T} , we have

$$\begin{aligned} X \overset{L}{\otimes}_C S &= (I_2[-1] \oplus P_1) \overset{L}{\otimes}_C S \\ &= I_2 \overset{L}{\otimes}_C S[-1] \oplus P_1 \overset{L}{\otimes}_C S. \end{aligned}$$

Clearly, $P_1 \overset{L}{\otimes}_C S \cong T$. To compute $I_2 \overset{L}{\otimes}_C S$, we use the resolution

$$0 \rightarrow P_1^{\text{Ext}_{\mathcal{T}}^{-1}(T, T)} \rightarrow P_2 \rightarrow I_2 \rightarrow 0.$$

Tensoring with S , we get a distinguished triangle

$$T^{\text{Ext}_{\mathcal{T}}^{-1}(T, T)} \rightarrow T[-1] \rightarrow I_2 \overset{L}{\otimes}_C S \rightarrow .$$

Comparing with (3.6), we find that $I_2 \overset{L}{\otimes}_C S \cong T[2]$. Thus, we have indeed an isomorphism

$$\varphi : X \overset{L}{\otimes}_B S \rightarrow S[1]$$

in \mathcal{T} .

Now we check that φ is C -equivariant in \mathcal{T} . The left C -module structure on $X \overset{L}{\otimes}_B S$ is obtained from the (homotopy) C -action on $I_2[-1] \oplus P_1$ as given in (3.7).

Let μ be an element of $W = \text{Hom}_C(P_1, P_2) = \text{Ext}_{\mathcal{T}}^{-1}(T, T)$. We need to prove that the following diagram is commutative in \mathcal{T} .

$$\begin{array}{ccc} I_2[-1] \overset{L}{\otimes}_B S & \xrightarrow{\cong} & T[1] \\ \pi(\mu) \overset{L}{\otimes}_B \text{id}_S \downarrow & & \downarrow \mu \\ P_1 \overset{L}{\otimes}_B S & \xrightarrow{\cong} & T \end{array}$$

We write this out in triangles.

$$\begin{array}{ccccccc} T^{\text{Ext}^{-1}(T,T)} & \xrightarrow{\phi} & T[-1] & \xrightarrow{\alpha} & T[2] & \xrightarrow{\beta} & \\ \pi(\mu) \downarrow & & \downarrow & & \downarrow \mu & & \\ T & \longrightarrow & 0 & \longrightarrow & T[1] & \xrightarrow{\text{id}} & \end{array}$$

Rotating the triangles, we need to prove that the following square is commutative.

$$\begin{array}{ccc} T[1] & \xrightarrow{\beta[-1]} & T^{\text{Ext}^{-1}(T,T)} \\ \mu \downarrow & & \downarrow \pi(\mu) \\ T & \xlongequal{\quad} & T \end{array}$$

This commutativity holds precisely because of the definition of π . So, ϕ is indeed C -equivariant.

But, according to [KR08, Lemma A.2.2], any C -equivariant morphism in \mathcal{T} between objects in ${}^c\mathcal{T}$ may be lifted to a morphism in ${}^c\mathcal{T}$. This finishes the proof. \square

We now have a functor

$$? \overset{L}{\otimes}_C T : \mathcal{C} \rightarrow \mathcal{T}$$

and, by Lemma 3.4.2, one finds that $a[-1](?) \overset{L}{\otimes}_C T$ is isomorphic to $? \overset{L}{\otimes}_C T$. By the universal property of orbit categories [Kel05], we obtain a triangulated functor

$$Q : \mathcal{D} \rightarrow \mathcal{T}$$

that sends P_1 to T .

LEMMA 3.4.3. *Q is an equivalence.*

Proof. We observe that analogues of the distinguished triangles (3.5) exist in \mathcal{D} (with P_1 replacing T). Indeed, let $N \in \text{Ind}(\mathcal{D})$. By (3.3), we have $N \in \text{Ind}(\mathcal{H}) \cup \{I_2[-1]\}$. If $N \in \text{Ind}(\mathcal{H})$, then $N[1] \cong aN$ and the analogue of (3.5) is simply the image in \mathcal{D} of the projective resolution of aN in \mathcal{H} (taking into account that $P_2 = a^{-1}P_1 = P_1[-1]$).

If $N = I_2[-1]$, then $N[1] = I_2$ and the analogue of (3.5) is the image in \mathcal{D} of the projective resolution of I_2 in \mathcal{H} .

To prove that Q is fully faithful, we have to prove that Q induces an isomorphism $\text{Hom}_{\mathcal{D}}(M, N) \rightarrow \text{Hom}_{\mathcal{T}}(QM, QN)$. Using the analogues of (3.5), we reduce to $M = P_1[i]$. But, since $\text{Hom}_{\mathcal{D}}(P_1[i], N) = \text{Hom}_{\mathcal{D}}(P_1[-1], N[-i-1])$, we reduce in fact to $M = P_1[-1]$. It now suffices to apply $\text{Hom}_{\mathcal{D}}(P_1[-1], -)$ to

$$P_1^a \rightarrow P_1^b \oplus P_1[-1]^c \rightarrow N[1] \rightarrow,$$

taking into account that $\text{Hom}_{\mathcal{D}}(M, N) \rightarrow \text{Hom}_{\mathcal{T}}(QM, QN)$ is an isomorphism for $M = P_1, N = P_1, P_1[1], P_2[2]$ by Lemma 3.3.2.

As a last step, we need to prove that Q is essentially surjective. But this follows from the distinguished triangles (3.5) together with the fact that $QP_1 = T$. \square

To finish the proof of Theorem 1.4, we observe that since \mathcal{T} is 3-Calabi–Yau, so is \mathcal{D} . Hence, by Lemma 3.3.1, π is either self-adjoint or anti-self-adjoint. By Lemma 3.2.2, we deduce that $a^2 \cong \tau$ and hence we may write $a = \tau^{1/2}$.

Remark 3.4.4. It would be interesting to deduce the fact that π is (anti-)self-adjoint directly from the Calabi–Yau property of \mathcal{T} , without going through the construction of \mathcal{D} first. This would have made our arguments above more elegant.

Remark 3.4.5. Iyama and Yoshino also considered $2n + 1$ -Calabi–Yau categories \mathcal{T} equipped with a $2n + 1$ -cluster tilting object T such that $\text{End}(T) = k$ and $\text{Ext}^{-i}(T, T) = 0$ for $0 < i < n$. They related such \mathcal{T} to the representation theory of the generalized Kronecker quiver Q_m , where $m = \dim \text{Ext}^{-n}(T, T)$.

One may show that our techniques are applicable to this case as well and yield $\mathcal{T} \cong D^b(\text{mod}(kQ_m))/(\tau^{1/2}[-n])$. We thank Osamu Iyama for bringing this point to our attention.

4. The singularity category of graded Gorenstein rings

4.1 Orlov’s results

Let $A = k + A_1 + A_2 + \dots$ be a commutative finitely generated graded k -algebra. As in [AZ94], we write $\text{qgr}(A)$ for the quotient of $\text{gr}(A)$ by the Serre subcategory of graded finite length modules. We write $\pi : \text{gr}(A) \rightarrow \text{qgr}(A)$ for the quotient functor. If A is generated in degree one and $X = \text{Proj } A$, then by Serre’s theorem [Ser55] we have $\text{coh}(X) = \text{qgr}(A)$.

Now assume that A is Gorenstein. Then we have $\text{RHom}_A(k, A) \cong k(a)[-d]$, where d is the Krull dimension of R and $a \in \mathbb{Z}$. The number a is called the Gorenstein parameter of A (see [Orl09a, Definition 2.1]).

Example 4.1.1. If A is a polynomial ring in n variables (of degree one), then $d = n, a = n$.

For use below we record another incarnation of the Gorenstein parameter. Let A' be the graded k -dual of A . Then

$$R\Gamma_{A_{>0}}(A) \cong A'(a)[-d], \tag{4.1}$$

where $R\Gamma_{A_{>0}}$ denotes cohomology with support in the ideal $A_{>0}$.

The following is a particular case of [Orl09a, Theorem 2.5].

THEOREM 4.1.2. *If $a \geq 0$, then there are fully faithful functors*

$$\Phi_i : \text{MCM}_{\text{gr}}(A) \rightarrow D^b(\text{qgr}(A))$$

such that for $\mathcal{T}_i = \Phi_i \underline{\text{MCM}}_{\text{gr}}(A)$ there is a semi-orthogonal decomposition

$$D^b(\text{qgr}(A)) = \langle \pi A(-i - a + 1), \dots, \pi A(-i), \mathcal{T}_i \rangle.$$

Hence, under the hypotheses of the theorem, we obtain in particular that

$$\underline{\text{MCM}}_{\text{gr}}(A) \cong {}^\perp \langle \pi A(-i - a + 1), \dots, \pi A(-i) \rangle \subset D^b(\text{qgr}(A))$$

for arbitrary i .

4.2 The action of the shift functor on the singularity category

Unfortunately, the functors Φ_i introduced in the previous section are not compatible with $?(1)$. Our aim in this section is to understand how $?(1)$ acts on the image of Φ_i . This requires us to dig deeper into Orlov’s construction, which has the unusual feature of depending on the category $D^b(\text{gr}_{\geq i} A)$, where $\text{gr}_{\geq i} A$ are the finitely generated graded A -modules with non-zero components concentrated in degrees $\geq i$. The quotient functor

$$D^b(\text{gr}_{\geq i} A) \hookrightarrow D^b(\text{gr} A) \xrightarrow{\pi} D^b(\text{qgr} A)$$

has a right adjoint $R\omega_i A$. Its image is denoted by \mathcal{D}_i .

We let P_i be the graded projective A -module of rank one generated in degree i (i.e. $P_i = A(-i)$). Likewise, S_i is the simple A -module concentrated in degree i . As in [Orl09a], we put $\mathcal{P}_{\geq i} = \langle (P_j)_{j \geq i} \rangle$, $\mathcal{S}_{\geq i} = \langle (S_j)_{j \geq i} \rangle$, and obvious variants with other types of inequality signs. In [Orl09a], it is proved that the image \mathcal{T}_i of Φ_i is the left orthogonal to $\mathcal{P}_{\geq i}$ inside $D^b(\text{gr}_{\geq i} A)$. The identification of \mathcal{T}_i with the graded singularity category is through the composition

$$\mathcal{T}_i \cong D^b(\text{gr}_{\geq i} A) / \mathcal{P}_{\geq i} \cong D^b(\text{gr} A) / \text{perf}(A) \cong \underline{\text{MCM}}_{\text{gr}}(A). \tag{4.2}$$

Assume that $a \geq 0$. Then the relation between \mathcal{T}_i , \mathcal{D}_i is given by the following semi-orthogonal decompositions:

$$D^b(\text{gr} A) = \left\langle \mathcal{S}_{< i}, \overbrace{\mathcal{P}_{\geq i+a}, P_{i+a-1}, \dots, P_i, \mathcal{T}_i}^{D^b(\text{gr}_{\geq i} A)} \right\rangle_{\mathcal{D}_i \cong D^b(\text{qgr}(A))}.$$

This is a refinement of Theorem 4.1.2.

The category $\underline{\text{MCM}}_{\text{gr}}(A)$ comes equipped with the shift functor $?(1)$. We denote the induced endofunctor on \mathcal{T}_i by σ_i . We will now compute it.

LEMMA 4.2.1. *For $M \in \mathcal{T}_i \subset D^b(\text{qgr}(A))$, we have*

$$\sigma_i M = \text{cone}(\text{RHom}_{\text{qgr}(A)}(\pi A(-i), M) \otimes_k \pi A(-i + 1) \rightarrow M(1)), \tag{4.3}$$

where the symbol ‘cone’ is to be understood in a functorial sense, for example by computing it on the level of complexes after first replacing M by an injective resolution.

Proof. Let $N \in \mathcal{T}_i \subset D^b(\text{gr}(A))$. To compute $\sigma_i N$, we see by (4.2) that we have to find $\sigma_i N \in \mathcal{T}_i$ such that $\sigma_i N \cong N(1)$ up to projectives. It is clear that we should take

$$\begin{aligned} \sigma_i N &= \text{cone}(\text{RHom}_{\text{gr}(A)}(P_{i-1}, N(1)) \otimes_k P_{i-1} \rightarrow N(1)) \\ &= \text{cone}(\text{RHom}_{\text{gr}(A)}(P_i, N) \otimes_k P_{i-1} \rightarrow N(1)). \end{aligned}$$

Now we note that RHom can be computed in $\mathcal{D}_i \cong D^b(\text{qgr}(A))$. Furthermore, since the result lies in $\mathcal{T}_i \subset \mathcal{D}_i$, we can characterize it uniquely by applying π to it. Since π commutes with $?(1)$, we obtain (4.3) with $M = \pi N$. □

4.3 The Serre functor for a graded Gorenstein ring

Let A, a, d be as above but now assume that A has an isolated singularity and let $M, N \in \underline{\text{MCM}}_{\text{gr}}(A)$. Then by a variant of [OY08, Theorem 8.3] we have a canonical graded isomorphism

$$\text{Ext}_A^d(\underline{\text{Hom}}_A(M, N), A) \cong \underline{\text{Hom}}_A(N, M[d - 1])$$

and furthermore an appropriate version of local duality yields

$$\text{Ext}_A^d(\underline{\text{Hom}}_A(M, N), A) = \underline{\text{Hom}}_A(M, N)^*(a).$$

In other words, we find that

$$\underline{\text{Hom}}_A(M, N)^* = \underline{\text{Hom}}_A(N, M[d - 1](-a))$$

and hence the Serre functor S on $\underline{\text{MCM}}(A)$ is given by $?[d - 1](-a)$.

It is customary to write $S = \tau[1]$, so that we have the usual formula

$$\underline{\text{Hom}}_A(M, N)^* = \text{Ext}^1(N, \tau M).$$

In this setting, we find that

$$\tau = ?[d - 2](-a). \tag{4.4}$$

4.4 The Gorenstein parameter of a Veronese subring

We remind the reader of the following well-known result.

PROPOSITION 4.4.1. *Let B be a polynomial ring in n variables of degree one. Assume that $m|n$ and let $B^{(m)}$ be the corresponding Veronese subring of B , i.e. $B_i^{(m)} = B_{mi}$. Then $B^{(m)}$ is Gorenstein with Gorenstein parameter n/m .*

Proof. The Gorenstein property is standard. To compute the Gorenstein invariant, we first let A be the ‘blown-up’ Veronese, i.e.

$$A_i = \begin{cases} B_i & \text{if } m|i, \\ 0 & \text{otherwise.} \end{cases}$$

Let $a, b = n$ be respectively the Gorenstein parameters of A and B . If M is a B -module, write M^+ for $\bigoplus_i M_{mi}$, considered as a graded A -module. We have

$$\begin{aligned} A'(a)[-n] &= R\Gamma_{A>0}(A) \quad (\text{see (4.1)}) \\ &= R\Gamma_{A>0}(B)^+ \\ &= R\Gamma_{B>0}(B)^+ \\ &= (B'(b)[-n])^+ \\ &= A'(b)[-n]. \end{aligned}$$

In the third equality, we have used that local homology is insensitive to finite extensions. We deduce that $a = b = n$. Since $B^{(m)}$ is obtained from A by dividing the grading by m , we obtain n/m as the Gorenstein parameter for $B^{(m)}$. □

Remark 4.4.2. In characteristic zero we could have formulated the result for invariant rings of finite subgroups of $\text{Sl}_n(k)$ (with the same proof). However, in finite characteristic Veronese subrings are not always invariant rings (consider the case where the characteristic divides m).

5. The Iyama–Yoshino examples (again)

5.1 Example 1.1

Let $B = k[x_1, x_2, x_3]$ and $A = B^{(3)}$. We have $X \stackrel{\text{def}}{=} \text{Proj } A = \text{Proj } B = \mathbb{P}^2$. By Proposition 4.4.1, A has Gorenstein invariant 1.

Unfortunately, we have to deal with the unpleasant notational problem that the shift functors on $\text{coh}(\mathbb{P}^2)$ coming from A and B do not coincide. To be consistent with §§ 4.1 and 4.2, we will denote them respectively by $?(1)$ and $?\{1\}$. Thus, $?(1) = ?\{3\}$. Note that this choice is rather unconventional.


According to Theorem 4.1.2, we have a semi-orthogonal decomposition

$$D^b(\text{coh}(X)) = \langle \mathcal{O}_{\mathbb{P}^2}, \mathcal{T}_0 \rangle.$$

From the fact that $D^b(\text{coh}(X))$ has a strong exceptional collection $\mathcal{O}_{\mathbb{P}^2}, \mathcal{O}_{\mathbb{P}^2}\{1\}, \mathcal{O}_{\mathbb{P}^2}\{2\}$, we deduce that there is a semi-orthogonal decomposition

$$\mathcal{T}_0 = \langle \mathcal{O}_{\mathbb{P}^2}\{1\}, \mathcal{O}_{\mathbb{P}^2}\{2\} \rangle.$$

In particular, $\text{RHom}_{\mathbb{P}^2}(\mathcal{O}_{\mathbb{P}^2}\{1\} \oplus \mathcal{O}_{\mathbb{P}^2}\{2\}, -)$ defines an equivalence between \mathcal{T}_0 and the representations of the quiver Q_3



where $V = kx_1 + kx_2 + kx_3$ and where $\mathcal{O}_{\mathbb{P}^2}\{i\}$ corresponds to the vertex labeled by i . By (4.4), the Auslander–Reiten translate on $\underline{\text{MCM}}_{\text{gr}}(A)$ is given by $?[1](-1)$. In other words, the shift functor on $\underline{\text{MCM}}_{\text{gr}}(A)$ is given by $(\tau[-1])^{-1}$. By Proposition A.8, we find (using $R = \widehat{A}$) that

$$\underline{\text{MCM}}(R) \cong \underline{\text{MCM}}_{\text{gr}}(A)/(1) \cong D^b(\text{mod}(kQ_3))/(\tau[-1]),$$

which is what we wanted to show.

Remark 5.1.1. Note that in this example we had no need for the somewhat subtle formula (4.3).

5.2 Example 1.2

We use similar conventions as in the previous section. Let $B = k[x_1, x_2, x_3, x_4]$ and $A = B^{(2)}$. We have $X = \text{Proj } A \cong \text{Proj } B = \mathbb{P}^3$ and we denote the corresponding shift functors by $?(1), ?\{1\}$ so that $?(1) = ?\{2\}$. By Proposition 4.4.1, A has Gorenstein invariant 2. By Theorem 4.1.2, we have a semi-orthogonal decomposition

$$D^b(\text{coh}(X)) = \langle \mathcal{O}_{\mathbb{P}^3}, \mathcal{O}_{\mathbb{P}^3}\{2\}, \mathcal{T}_{-1} \rangle.$$

Now $D^b(\text{coh}(X))$ has a strong exceptional collection $\mathcal{O}_{\mathbb{P}^3}, \mathcal{O}_{\mathbb{P}^3}\{1\} \cdot \mathcal{O}_{\mathbb{P}^3}\{2\}, \mathcal{O}_{\mathbb{P}^3}\{3\}$. This sequence is geometric [BP94, Proposition 3.3] and hence by every mutation is strongly exceptional [BP94, Theorem 2.3]. We get in particular the following strongly exceptional collection: $\mathcal{O}_{\mathbb{P}^3}, \mathcal{O}_{\mathbb{P}^3}\{2\} \cdot \Omega_{\mathbb{P}^3}^*\{1\}, \mathcal{O}_{\mathbb{P}^3}\{3\}$, where $\Omega_{\mathbb{P}^3}$ is defined by the exact sequence

$$0 \rightarrow \Omega_{\mathbb{P}^3} \rightarrow V \otimes \mathcal{O}_{\mathbb{P}^3}\{-1\} \rightarrow \mathcal{O}_{\mathbb{P}^3} \rightarrow 0, \tag{5.1}$$


where $V = kx_1 + kx_2 + kx_3 + kx_4$. Thus, there is a semi-orthogonal decomposition

$$\mathcal{T}_{-1} = \langle \Omega_{\mathbb{P}^3}^*\{1\}, \mathcal{O}_{\mathbb{P}^3}\{3\} \rangle.$$

An easy computation yields

$$\text{RHom}_{\mathbb{P}^3}(\Omega_{\mathbb{P}^3}^*\{1\}, \mathcal{O}_{\mathbb{P}^3}\{3\}) = \wedge^2 V.$$

$\mathrm{RHom}_{\mathbb{P}^3}(\Omega_{\mathbb{P}^3}^*\{1\} \oplus \mathcal{O}_{\mathbb{P}^3}\{3}, -)$ defines an equivalence between \mathcal{T}_{-1} and the representations of the quiver Q_6 :



Put $W = \wedge^2 V$ and choose an arbitrary trivialization $\wedge^4 V \cong k$. Let $\pi : W \rightarrow W^*$ be the resulting (self-adjoint) isomorphism. We are in the setting of § 3.2 and hence can define $\tau^{1/2}$ as acting on the derived category of Q_6 .

We will now compute $\sigma_{-1}(\Omega_{\mathbb{P}^3}^*\{1\}), \sigma_{-1}(\mathcal{O}_{\mathbb{P}^3}\{3\})$. An easy computation yields

$$\mathrm{RHom}_{\mathbb{P}^3}(\mathcal{O}_{\mathbb{P}^3}\{2\}, \Omega_{\mathbb{P}^3}^*\{1\}) = V^*,$$

$$\mathrm{RHom}_{\mathbb{P}^3}(\mathcal{O}_{\mathbb{P}^3}\{2\}, \mathcal{O}_{\mathbb{P}^3}\{3\}) = V.$$

Using the formula (4.3), we find that

$$\sigma_{-1}(\mathcal{O}_{\mathbb{P}^3}\{3\}) = \mathrm{cone}(V \otimes \mathcal{O}_{\mathbb{P}^3}\{4\} \rightarrow \mathcal{O}_{\mathbb{P}^3}\{5\}) = \Omega_{\mathbb{P}^3}\{5\}[1], \tag{5.2}$$

$$\sigma_{-1}(\Omega_{\mathbb{P}^3}^*\{1\}) = \mathrm{cone}(V^* \otimes \mathcal{O}_{\mathbb{P}^3}\{4\} \rightarrow \Omega_{\mathbb{P}^3}^*\{3\}) = \mathcal{O}_{\mathbb{P}^3}\{3\}[1], \tag{5.3}$$

where in the second line we have used the dual version of (5.1).

Let P_i be the projective representation of Q_6 generated in vertex i . The endofunctor on $D^b(\mathrm{mod}(kQ_6))$ induced by σ_{-1} will be denoted by the same letter. We will now compute it. From (5.3), we deduce immediately that $\sigma_{-1}(P_1) = P_2[1]$. To analyze (5.2), we note that a suitably shifted slice of the Koszul sequence has the form

$$0 \rightarrow \wedge^4 V \otimes \Omega_{\mathbb{P}^3}^*\{1\} \rightarrow \wedge^2 V \otimes \mathcal{O}_{\mathbb{P}^3}\{3\} \rightarrow \Omega_{\mathbb{P}^3}\{5\} \rightarrow 0.$$

Thus, $\Omega_{\mathbb{P}^3}\{5\}$ corresponds to the cone of

$$\wedge^4 V \otimes P_1 \rightarrow \wedge^2 V \otimes P_2,$$

which is easily seen to be equal to $\wedge^4 V \otimes \tau^{-1}P_1$.

If we use our chosen trivialization $\wedge^4 V \cong k$, then we see that at least on objects σ_{-1} coincides with $\tau^{-1/2}[1]$. It is routine to extend this to an isomorphism of functors by starting with a bounded complex of projectives in $\mathrm{mod}(kQ_6)$.

By Proposition A.8, we find (using $R = \widehat{A}$) that

$$\underline{\mathrm{MCM}}(R) \cong \underline{\mathrm{MCM}}_{\mathrm{gr}}(A)/(1) \cong D^b(\mathrm{mod}(kQ_6))/(\tau^{1/2}[-1]),$$

which is what we wanted to show.

6. A remark on gradability of rigid modules

We keep notation as in the previous section. Since in the Iyama–Yoshino examples $\underline{\mathrm{MCM}}_{\mathrm{gr}}(A)$ is the derived category of a hereditary category, the functor

$$\underline{\mathrm{MCM}}_{\mathrm{gr}}(A) \rightarrow \underline{\mathrm{MCM}}_{\mathrm{gr}}(A)/(1)$$

is essentially surjective [Kel05] and hence

$$\underline{\mathrm{MCM}}_{\mathrm{gr}}(A) \rightarrow \underline{\mathrm{MCM}}_{\mathrm{gr}}(\widehat{A})$$

is also essentially surjective. In more complicated examples there is no reason however why this should be the case. Nevertheless, we have the following result, which is probably well known.

PROPOSITION 6.1. *Assume that k has characteristic zero. Let $A = k + A_1 + A_2 + \dots$ be a left noetherian graded k -algebra. Put $R = \widehat{A}$. Let $M \in \text{mod}(R)$ be such that $\text{Ext}_R^1(M, M) = 0$. Then M is the completion of a finitely generated graded A -module N .*

In the rest of this section, we let the notation and hypotheses be as in the statement of the proposition (in particular, k has characteristic zero). We denote the maximal ideal of R by m .

Let E be the Euler derivation on A and R , i.e. on A we have $E(a) = (\text{deg } a)a$ and we extend E to R in the obvious way. If $M \in \text{mod}(R)$, then we will define an Euler connection as a k -linear map $\nabla : M \rightarrow M$ such that $\nabla(am) = E(a)m + a\nabla(m)$. If $M = \widehat{N}$ for N a graded A -module, then M has an associated Euler connection by extending $\nabla(n) = (\text{deg } n)n$ for n a homogeneous element of N .

LEMMA 6.2. *Let M be a finitely generated R module. Then M has an Euler connection if and only if M is the completion of a finitely generated graded A -module.*

Proof. We have already explained the easy direction. Conversely, assume that M has an Euler connection. For each n , we have that $M/m^n M$ is finite dimensional and hence it decomposes into generalized eigenspaces for ∇ :

$$M/m^n M = \prod_{\alpha \in k} (M/m^n M)_\alpha \quad (\text{finite product}).$$

Considering right exact sequences

$$(m/m^2)^{\otimes n} \otimes M/mM \rightarrow M/m^{n+1}M \rightarrow M/m^n M \rightarrow 0,$$

we easily deduce that the multiplicity of a fixed generalized eigenvalue in $M/m^n M$ stabilizes as $n \rightarrow \infty$. Thus, $M = \prod_{\alpha \in k} M_\alpha$, where M_α is a generalized eigenspace with eigenvalue α . We put $N' = \bigoplus_\alpha M_\alpha$. Then N' is noetherian since obviously any ascending chain of graded submodules of N' can be transformed into an ascending chain of submodules in M . In particular, N' is finitely generated and we have $M = \widehat{N}'$.

Now N' is k -graded and not \mathbb{Z} -graded. But we can decompose N' along \mathbb{Z} -orbits and then by taking suitable shifts we obtain a \mathbb{Z} -graded module with the same completion as N' . \square

Proof of Proposition 6.1. Let $\epsilon^2 = 0$ and consider $M[\epsilon]$, where A acts via $a \cdot m = (a + E(a)\epsilon)m$. We have a short exact sequence of A -modules

$$0 \rightarrow M\epsilon \rightarrow M[\epsilon] \rightarrow M \rightarrow 0,$$

which is split by hypotheses. Denote the splitting by $m + \nabla(m)\epsilon$. For $a \in A$, we have

$$am + \nabla(am)\epsilon = (a + E(a)\epsilon)(m + \nabla(m)\epsilon)$$

and hence

$$\nabla(am) = E(a)m + a\nabla(m).$$

Hence, ∇ is an Euler connection and so we may invoke Lemma 6.2 to show that $M = \widehat{N}$. \square

ACKNOWLEDGEMENTS

We thank Osamu Iyama, Idun Reiten, Srikanth Iyengar, and Amnon Neeman for commenting on a preliminary version of this paper. The second author thanks Kyoji Saito, Kazushi Ueda, and Osamu Iyama for pointing out the properties of the Henselization in Remark A.6, and

Apostolos Beligiannis for explaining his work [Bel00]. The third author is a director of research at the FWO. The results of this paper were partially obtained during visits of the third author to the Université Denis Diderot – Paris 7 and the Centre de Recerca Matemàtica in Barcelona. He hereby thanks these institutions for their hospitality.

Appendix A. Generators of singularity categories

Throughout, (A, \mathfrak{m}, k) is a (commutative) local noetherian ring, with maximal ideal \mathfrak{m} and residue field k . The *singularity category* of A is the Verdier quotient

$$D_{\text{Sg}}(A) := D^b(\text{mod } A)/K^b(\text{proj } A)$$

of the bounded derived category of finitely generated A -modules by the full subcategory of perfect complexes. Recall that a functor $F : \mathcal{T} \rightarrow \mathcal{S}$ is an *equivalence up to direct summands* if F is fully faithful and every object $X \in \mathcal{S}$ is a direct summand of $F(Y)$ for some $Y \in \mathcal{T}$. We say that A is a *G-ring* if the canonical morphism from A to its \mathfrak{m} -adic completion $A \rightarrow \widehat{A}$ is regular [Mat89, §32], and that A has an *isolated singularity* if $A_{\mathfrak{p}}$ is regular for every non-maximal prime ideal \mathfrak{p} of A . Our main result about singularity categories is the following.

PROPOSITION A.1. *Let A be a local noetherian ring with an isolated singularity, which is also a G-ring (e.g. A is essentially of finite type over a field). Then the canonical functor*

$$\gamma := - \otimes_A \widehat{A} : D_{\text{Sg}}(A) \rightarrow D_{\text{Sg}}(\widehat{A})$$

is an equivalence up to direct summands.

This is a special case of a general result by Orlov [Orl09b] (which was obtained independently). Our methods are quite different, however.

When A is Gorenstein, there is an equivalence, due to Buchweitz [Buc87], between $D_{\text{Sg}}(A)$ and the stable category of maximal Cohen–Macaulay A -modules $\underline{\text{MCM}}(A)$, so in this case we obtain Proposition 1.6. We remark that, in general, γ is not an equivalence (see e.g. Example A.5).

Let us outline the proof of the proposition. Recall that a *thick subcategory* of a triangulated category \mathcal{T} is a triangulated subcategory closed under retracts. Given an object C of \mathcal{T} , we say that an object X is *finitely built* from C if it belongs to the smallest thick subcategory of \mathcal{T} containing C . If every object of \mathcal{T} has this property, that is, if there are no proper thick subcategories of \mathcal{T} containing C , then C is said to *classically generate* \mathcal{T} .

The local ring A and its completion \widehat{A} have the same residue field k , and it is not difficult to see that γ induces an equivalence between the respective subcategories consisting of objects finitely built from k . The subtlety lies in showing that, because A has an isolated singularity, every object can be finitely built from k . Our proof of this fact uses homotopy colimits, which presents a technical problem since $D_{\text{Sg}}(A)$ lacks infinite coproducts. One approach is to enlarge the category $D_{\text{Sg}}(A)$ by considering the Verdier quotient

$$D'_{\text{Sg}}(A) := D^b(\text{Mod } A)/K^b(\text{Proj } A)$$

of the bounded derived category of all A -modules by the full subcategory of bounded complexes of projective A -modules. By [Orl04, Proposition 1.13], the canonical functor $D_{\text{Sg}}(A) \rightarrow D'_{\text{Sg}}(A)$ is fully faithful, and $D'_{\text{Sg}}(A)$ turns out to contain enough coproducts (and thus homotopy colimits) for our purposes. Throughout, $D(A)$ denotes the (unbounded) derived category of A -modules.

The next proposition follows immediately from the work of Schoutens [Sch03], but we give a direct proof in the special case of an isolated singularity. The result also follows from the general result by Orlov [Orl09b], and Dyckerhoff [Dyc09] has given a proof based on the theory of matrix factorizations in the hypersurface case.

PROPOSITION A.2. *A local noetherian ring (A, \mathfrak{m}, k) has an isolated singularity if and only if $D_{\text{Sg}}(A)$ is classically generated by k .*

Proof. We begin with the easy direction. Suppose that $D_{\text{Sg}}(A)$ is classically generated by k , and let $\mathfrak{p} \neq \mathfrak{m}$ be a prime ideal. The canonical functor $-\otimes_A A_{\mathfrak{p}} : D_{\text{Sg}}(A) \rightarrow D_{\text{Sg}}(A_{\mathfrak{p}})$ is identically zero, because it sends the generator k to zero. The image of this functor contains the residue field $\kappa(\mathfrak{p}) = A/\mathfrak{p} \otimes_A A_{\mathfrak{p}}$, from which we deduce that $\kappa(\mathfrak{p})$ has finite projective dimension over $A_{\mathfrak{p}}$. Hence, $A_{\mathfrak{p}}$ is regular, and we may conclude that A has an isolated singularity.

Now suppose that A has an isolated singularity, and let M in $D^b(\text{mod } A)$ be given. The idea is to write M as a homotopy colimit³ of a sequence of bounded complexes with finite length cohomology; it follows that M is a direct summand of one of the terms in this sequence, from which we conclude that k classically generates $D_{\text{Sg}}(A)$. First, we set up some notation. Given $a \in A$, define complexes

$$K[a] := A \xrightarrow{a} A \quad \text{and} \quad E[a] := A \xrightarrow{\text{can}} A[a^{-1}],$$

both concentrated in degrees zero and one, and observe that the commutative diagram

$$\begin{array}{ccccccc} A & \xrightarrow{1} & A & \xrightarrow{1} & A & \xrightarrow{1} & \cdots \\ a \downarrow & & a^2 \downarrow & & a^3 \downarrow & & \\ A & \xrightarrow{a} & A & \xrightarrow{a} & A & \xrightarrow{a} & \cdots \end{array}$$

is a direct system of complexes $K[a] \rightarrow K[a^2] \rightarrow K[a^3] \rightarrow \cdots$ with colimit $E[a]$. More generally, given a sequence $\mathbf{a} = \{a_1, \dots, a_d\}$ in A , we define $K[\mathbf{a}] := \bigotimes_{j=1}^d K[a_j]$ and $E[\mathbf{a}] := \bigotimes_{j=1}^d E[a_j]$. Setting $\mathbf{a}^i = \{a_1^i, \dots, a_d^i\}$, there is a canonical isomorphism $E[\mathbf{a}] \cong \varinjlim_i K[\mathbf{a}^i]$ and thus a triangle

$$\bigoplus_{i \geq 1} K[\mathbf{a}^i] \xrightarrow{1\text{-shift}} \bigoplus_{i \geq 1} K[\mathbf{a}^i] \longrightarrow E[\mathbf{a}] \longrightarrow \tag{A1}$$

in the derived category $D(A)$. This triangle expresses the fact that $E[\mathbf{a}]$ is the homotopy colimit of $K[\mathbf{a}^i]$ in $D(A)$. For background on homotopy colimits, see [BN93, Nee01].

Now let \mathbf{a} be a system of parameters for A , and extend the augmentation morphism $E[\mathbf{a}] \xrightarrow{\varepsilon} A$ to a triangle $E[\mathbf{a}] \rightarrow A \rightarrow \check{C}[\mathbf{a}] \rightarrow$, where the complex $\check{C}[\mathbf{a}] := \Sigma \ker(\varepsilon)$ is given in each degree by $\check{C}[\mathbf{a}]^t = \bigoplus_{i_0 < \dots < i_t} A[a_{i_0}^{-1}, \dots, a_{i_t}^{-1}]$. Tensoring with M , we obtain a triangle

$$E[\mathbf{a}] \otimes_A M \rightarrow M \rightarrow \check{C}[\mathbf{a}] \otimes_A M \rightarrow \tag{A2}$$

in $D(A)$. Since A has an isolated singularity, $M[a_{i_0}^{-1} \cdots a_{i_t}^{-1}]$ has finite projective dimension over $A[a_{i_0}^{-1} \cdots a_{i_t}^{-1}]$, and hence also over A , for every sequence of indices $i_0 < \dots < i_t$ in $\{1, \dots, d\}$. Here we use the fact that $A[a_{i_0}^{-1} \cdots a_{i_t}^{-1}]$ has finite projective dimension as an A -module.⁴

³To be precise, we do not consider homotopy colimits in $D'_{\text{Sg}}(A)$, since coproducts in this category are rather subtle. Instead, we consider the image under the quotient functor $D^b(\text{Mod } A) \rightarrow D'_{\text{Sg}}(A)$ of homotopy colimits in $D^b(\text{Mod } A)$.

⁴By induction, this reduces to the observation that $\text{pd}_A A[a^{-1}] \leq 1$, which holds because $A[a^{-1}] = A[X]/(aX - 1)$.

We conclude that $\check{C}[\mathbf{a}] \otimes_A M$ is, up to isomorphism in $D(A)$, a bounded complex of projective A -modules, whence the triangle (A2) gives rise to an isomorphism $E[\mathbf{a}] \otimes_A M \cong M$ in $D'_{\text{Sg}}(A)$. Note that the coproduct $\bigoplus_{i \geq 1} K[\mathbf{a}^i] \otimes_A M$ is bounded, so tensoring (A1) with M yields a triangle in $D'_{\text{Sg}}(A)$ of the form

$$\bigoplus_{i \geq 1} K[\mathbf{a}^i] \otimes_A M \xrightarrow{1\text{-shift}} \bigoplus_{i \geq 1} K[\mathbf{a}^i] \otimes_A M \longrightarrow M \longrightarrow . \tag{A3}$$

In what follows, let $\text{Hom}(-, -)$ denote morphism sets in $D'_{\text{Sg}}(A)$. One can check (see Lemma A.4 below) that $\text{Hom}(M, -)$ commutes with coproducts coming from $D^b(\text{Mod } A)$ via the quotient functor, so applying $\text{Hom}(M, -)$ to (A3) and using the argument of [Nee96, Lemma 2.8] we deduce that

$$\text{Hom}(M, M) \cong \varinjlim_i \text{Hom}(M, K[\mathbf{a}^i] \otimes_A M).$$

In particular, the identity $1_M : M \rightarrow M$ corresponds to a split monomorphism $M \rightarrow K[\mathbf{a}^k] \otimes_A M$ in $D'_{\text{Sg}}(A)$ for some $k \geq 1$. The functor $D_{\text{Sg}}(A) \rightarrow D'_{\text{Sg}}(A)$ is fully faithful, so M is also a direct summand of $K[\mathbf{a}^k] \otimes_A M$ in $D_{\text{Sg}}(A)$. The cohomology modules of $K[\mathbf{a}^k] \otimes_A M$ have finite length (\mathbf{a} is a system of parameters), so this complex is an iterated extension in $D^b(\text{mod } A)$ of finite direct sums of copies of k . It is now clear that any thick subcategory of $D_{\text{Sg}}(A)$ containing k must contain M and, since M was arbitrary, this completes the proof. \square

LEMMA A.3. *A morphism $\varphi : M \rightarrow C$ in $D(A)$ with $M \in D^b(\text{mod } A)$ and $C \in K^b(\text{Proj } A)$ factors, in $D(A)$, as $M \rightarrow Q \rightarrow C$ for some $Q \in K^b(\text{proj } A)$.*

Proof. We may, without loss of generality, assume that M is a bounded above complex of finitely generated projective A -modules, that C is a bounded complex of free A -modules, and that φ is a morphism of complexes. Let $n \in \mathbb{Z}$ be such that $C^i = 0$ for $i < n$. The image of $\varphi^n : M^n \rightarrow C^n$ is finitely generated, so let Q^n be a finite free submodule of C^n with the property that φ^n factors as $M^n \rightarrow Q^n \rightarrow C^n$. Similarly, let Q^{n+1} be a finite free submodule of C^{n+1} with the property that $\text{Im}(\varphi^{n+1}) + \partial(Q^n) \subseteq Q^{n+1}$, where ∂ is the differential. Then φ^{n+1} factors as $M^{n+1} \rightarrow Q^{n+1} \rightarrow C^{n+1}$ and the differential restricts to a map $\partial|_Q : Q^n \rightarrow Q^{n+1}$. Proceeding in this way, we construct a bounded complex Q of finite free A -modules and a factorization $M \rightarrow Q \rightarrow C$, as required. \square

LEMMA A.4. *Let $\{X_i\}_{i \in I}$ be a family of bounded complexes of A -modules such that there exist $a, b \in \mathbb{Z}$ with $X_i^k = 0$ for all $k \notin [a, b]$ and $i \in I$. Then, given $M \in D^b(\text{mod } A)$, the canonical map*

$$\bigoplus_i \text{Hom}_{D'_{\text{Sg}}(A)}(M, X_i) \longrightarrow \text{Hom}_{D'_{\text{Sg}}(A)}\left(M, \bigoplus_i X_i\right)$$

is an isomorphism, where $\bigoplus_i X_i$ denotes the degree-wise coproduct of complexes.

Proof. By a standard argument, it is enough to prove that any morphism $M \rightarrow \bigoplus_i X_i$ in $D'_{\text{Sg}}(A)$ factors through a finite subcoproduct. Such a morphism is defined by a diagram in the form of a roof

$$\begin{array}{ccc} & Y & \\ f \swarrow & & \searrow \\ M & & \bigoplus_i X_i \end{array} \tag{A4}$$

in $D^b(\text{Mod } A)$, where the cone of f is a bounded complex C_f of projective A -modules. Extending f to a triangle $Y \rightarrow M \rightarrow C_f \rightarrow$ in $D^b(\text{Mod } A)$, we deduce from Lemma A.3 that $M \rightarrow C_f$ factors as $M \rightarrow Q \rightarrow C_f$ for some $Q \in K^b(\text{proj } A)$. Let C' denote the cone of $M \rightarrow Q$. From the octahedral axiom applied to the pair $(M \rightarrow Q, Q \rightarrow C_f)$, we obtain a commutative diagram in $D^b(\text{Mod } A)$ of the form

$$\begin{array}{ccc} \Sigma^{-1}C' & & \\ \downarrow h & \searrow & \\ Y & \xrightarrow{f} & M \end{array}$$

where the cone of h belongs to $K^b(\text{Proj } A)$. The upshot is that the morphism in $D'_{\text{Sg}}(A)$ represented by the roof in (A4) may also be represented by a roof with $Y \in D^b(\text{mod } A)$ (replace Y with $\Sigma^{-1}C'$). In this case, Y is compact in $D^b(\text{Mod } A)$ by [Rou08, Proposition 6.15], so the morphism $Y \rightarrow \bigoplus_i X_i$ in the roof factors through a finite subcoproduct, which implies that $M \rightarrow \bigoplus_i X_i$ factors through a finite subcoproduct in $D'_{\text{Sg}}(A)$. \square

Proof of Proposition A.1. To begin with, let A denote an arbitrary local noetherian ring, and consider the canonical functor

$$\gamma' := -\otimes_A \widehat{A} : D'_{\text{Sg}}(A) \rightarrow D'_{\text{Sg}}(\widehat{A}).$$

Restriction of scalars defines a functor $(-)_A : D^b(\text{Mod } \widehat{A}) \rightarrow D^b(\text{Mod } A)$ that sends a bounded complex of projective \widehat{A} -modules to a bounded complex of flat A -modules. Since flat A -modules have finite projective dimension by [RG71, Part II, Corollary 3.2.7], there is an induced functor

$$(-)_A : D'_{\text{Sg}}(\widehat{A}) \rightarrow D'_{\text{Sg}}(A)$$

right adjoint to γ' . The unit of this adjunction is the canonical morphism

$$1 \rightarrow (-\otimes_A \widehat{A})_A,$$

which is obviously an isomorphism on k , and thus also an isomorphism on the smallest thick subcategory \mathcal{S} of $D'_{\text{Sg}}(A)$ containing k . By a standard argument of category theory, the restriction of γ' to \mathcal{S} is fully faithful. In particular, γ induces an equivalence of the smallest triangulated subcategory of $D_{\text{Sg}}(A)$ containing k with the smallest triangulated subcategory of $D_{\text{Sg}}(\widehat{A})$ containing k .

Now we assume that A is a G-ring with an isolated singularity. The (only) reason for assuming that A is a G-ring is that this guarantees that the completion \widehat{A} has an isolated singularity [Wie98, Lemma 2.7]. By Proposition A.2, the subcategory \mathcal{S} includes the image of $D_{\text{Sg}}(A)$ under the canonical embedding $D_{\text{Sg}}(A) \rightarrow D'_{\text{Sg}}(A)$, from which we infer that γ is fully faithful. It follows from a second application of Proposition A.2 that the thick closure of $D_{\text{Sg}}(A)$ in $D_{\text{Sg}}(\widehat{A})$ is all of $D_{\text{Sg}}(\widehat{A})$. Since the thick closure of a triangulated subcategory is just the class of all direct summands of objects in the subcategory [Nee01, Remark 2.1.39], γ is an equivalence up to direct summands. \square

It is easy to construct examples where γ is not an equivalence. It suffices to give a Cohen–Macaulay module over the completion of a Gorenstein local ring \widehat{A} which is not *extended* from A , i.e. which is not of the form \widehat{M} for a Cohen–Macaulay A -module.

Example A.5. Let $A = \mathbb{C}[X, Y]_{(X, Y)} / (X^3 + X^2 - Y^2)$ be the local ring of a node, so the completion of A is isomorphic to the reduced ring $S = \mathbb{C}[[U, V]] / (UV)$. This is a singularity

of type (A_1) and, by [Yos90, (9.9)], there are, up to isomorphism, exactly three indecomposable maximal Cohen–Macaulay S -modules, which are

$$S, \mathfrak{p} = US \quad \text{and} \quad \mathfrak{q} = VS.$$

Clearly, $S/\mathfrak{p} \cong \mathfrak{q}$, whence $\mathfrak{q} \cong \Sigma\mathfrak{p}$ in $D_{\text{Sg}}(S)$. Since $\mathfrak{p}, \mathfrak{q}$ are minimal prime ideals, $S_{\mathfrak{p}}$ and $S_{\mathfrak{q}}$ are fields, and it follows from a result of Levy and Odenthal [LO96, Theorem 6.2] that a finitely generated S -module M is extended if and only if $\text{rank}_{S_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \text{rank}_{S_{\mathfrak{q}}}(M_{\mathfrak{q}})$. Hence, \mathfrak{p} and \mathfrak{q} are not extended, and thus not in the essential image of γ , but their direct sum $\mathfrak{p} \oplus \mathfrak{q}$ is extended. This corresponds to the fact that the nodal curve is irreducible, while the curve $XY = 0$ has two irreducible components. Another argument that $\mathfrak{p} \oplus \mathfrak{q} \cong \mathfrak{p} \oplus \Sigma\mathfrak{p}$ in $D_{\text{Sg}}(S)$ belongs to the essential image of γ uses K -theory: simply apply [Nee01, Corollary 4.5.12].

Note that $\{U - V\}$ is a system of parameters for S . It follows from the proof of Proposition A.2 that \mathfrak{p} is a direct summand in $D_{\text{Sg}}(S)$ of $K[(U - V)^n] \otimes \mathfrak{p}$ for some $n \geq 1$. In fact, $K[(U - V)^n] \otimes \mathfrak{p} \xrightarrow{U^n} \mathfrak{p}$ is quasi-isomorphic to $\Sigma^{-1}\mathfrak{p}/\mathfrak{p}^{n+1}$, and \mathfrak{p} is a direct summand of $\Sigma^{-1}\mathfrak{p}/\mathfrak{p}^2$ in $D_{\text{Sg}}(S)$. To see this, observe that there is a triangle in the derived category

$$\mathfrak{p} \xrightarrow{U} \mathfrak{p} \longrightarrow \mathfrak{p}/\mathfrak{p}^2 \longrightarrow \Sigma\mathfrak{p},$$

and $U : \mathfrak{p} \longrightarrow \mathfrak{p}$ is zero in $D_{\text{Sg}}(S)$ (as it factors via S), so we may conclude that $\mathfrak{p} \oplus \Sigma\mathfrak{p} \cong \mathfrak{p}/\mathfrak{p}^2$ in $D_{\text{Sg}}(S)$. Since $\mathfrak{p}/\mathfrak{p}^2$ is isomorphic as an S -module to the residue field \mathbb{C} , we see for a third time that $\mathfrak{p} \oplus \Sigma\mathfrak{p} \cong \mathbb{C}$ is in the essential image of γ .

Remark A.6. Denoting by A^h the Henselization of A , the ring homomorphisms $A \longrightarrow A^h \longrightarrow \widehat{A}$ give rise to a factorization of γ as the composite

$$D_{\text{Sg}}(A) \xrightarrow{\gamma_1} D_{\text{Sg}}(A^h) \xrightarrow{\gamma_2} D_{\text{Sg}}(\widehat{A}),$$

where $\gamma_1 = -\otimes_A A^h$ and $\gamma_2 = -\otimes_{A^h} \widehat{A}$. In the situation of Proposition A.1, γ_2 is an equivalence: up to a shift, every object of $D_{\text{Sg}}(\widehat{A})$ is a finitely generated module M free on the punctured spectrum and, by Elkik’s theorem [Elk74, Théorème 3], such modules can be descended to the Henselization; that is, there exists a finitely generated A^h -module N such that $M \cong \widehat{N}$. In particular, γ is an honest equivalence (not just up to direct summands) when A is Henselian.

Now we give the proof of Proposition 1.5. In [Kra05], Krause produced an embedding $\mu : D_{\text{Sg}}(A) \hookrightarrow K_{\text{ac}}(\text{Inj } A)$, where $K_{\text{ac}}(\text{Inj } A)$ is the homotopy category of $C_{\text{ac}}(\text{Inj } A)$ of acyclic complexes of injective A -modules. This category is compactly generated, and μ induces an equivalence up to direct summands between $D_{\text{Sg}}(A)$ and the full subcategory of compact objects in $K_{\text{ac}}(\text{Inj } A)$.

The embedding μ produces a DG-enhancement for $D_{\text{Sg}}(A)$, where, for $M, N \in D_{\text{Sg}}(A)$, we put

$$\text{RHom}_{D_{\text{Sg}}(A)}(M, N) = \underline{\text{Hom}}_{C_{\text{ac}}(\text{Inj } A)}(\mu(M), \mu(N)).$$

If A is a noetherian \mathbb{Z} -graded ring (not necessarily commutative), then we may define the graded singularity category $D_{\text{Sg}}^{\text{gr}}(A)$ in the obvious way.

Since $D_{\text{Sg}}^{\text{gr}}(A)$ has an analogous DG-enhancement as $D_{\text{Sg}}(A)$, we may define the orbit category $D_{\text{Sg}}^{\text{gr}}(A)/(1)$ (see [Kel05]). By construction, $D_{\text{Sg}}^{\text{gr}}(A)/(1)$ is a triangulated category (with a DG-enhancement) equipped with an exact functor

$$\sigma : D_{\text{Sg}}^{\text{gr}}(A) \rightarrow D_{\text{Sg}}^{\text{gr}}(A)/(1)$$

such that $D_{\text{Sg}}^{\text{gr}}(A)/(1)$ is classically generated by its essential image and such that for $M, N \in D_{\text{Sg}}^{\text{gr}}(A)$ we have

$$\text{Hom}_{D_{\text{Sg}}^{\text{gr}}(A)/(1)}(\sigma M, \sigma N) = \bigoplus_i \text{Hom}_{D_{\text{Sg}}^{\text{gr}}(A)}(M, N(i)).$$

Forgetting the grading yields an exact functor

$$F : D_{\text{Sg}}^{\text{gr}}(A) \rightarrow D_{\text{Sg}}(A),$$

which makes the shift (1) isomorphic to the identity functor. Hence, by the universal property of orbit categories, F factors canonically through

$$\tilde{F} : D_{\text{Sg}}^{\text{gr}}(A)/(1) \rightarrow D_{\text{Sg}}(A).$$

LEMMA A.7. *The functor \tilde{F} is fully faithful.*

Proof. We have to prove that for $M, N \in D_{\text{Sg}}^{\text{gr}}(A)$ we have

$$\text{Hom}_{D_{\text{Sg}}(A)}(M, N) = \bigoplus_i \text{Hom}_{D_{\text{Sg}}^{\text{gr}}(A)}(M, N(i)).$$

By considering cones over suitable truncated projective resolutions, we may assume that M, N are finitely generated graded A -modules.

We then use the well-known formula

$$\text{Hom}_{D_{\text{Sg}}(A)}(M, N) = \text{inj} \lim_n \text{Hom}_A(\Omega^n M, \Omega^n N)$$

and the corresponding formula in the graded case. This reduces us to proving that

$$\text{Hom}_A(\Omega^n M, \Omega^n N) = \bigoplus_i \text{Hom}_{\text{Gr}(A)}(\Omega^n M, \Omega^n N(i)),$$

which follows from the fact that $\Omega^n M$ is finitely generated. □

PROPOSITION A.8. *Let $A = k + A_1 + A_2 \cdots$ be a finitely generated commutative graded k -algebra with the augmentation ideal $m = A_{>0}$ defining an isolated singularity. Then we have equivalences*

$$D_{\text{Sg}}^{\text{gr}}(A)/(1) \xrightarrow{\tilde{F}} D_{\text{Sg}}(A) \xrightarrow{(-)_m} D_{\text{Sg}}(A_m) \xrightarrow{\hat{A} \otimes_A -} D_{\text{Sg}}(\hat{A}).$$

Proof. The third functor is an equivalence because of Proposition A.1. The second functor is an equivalence because of [Orl04]. Finally, in Lemma A.7, we have shown that \tilde{F} is fully faithful. So, we have to show that it is essentially surjective. This is clear by Proposition A.2, since k lies in the essential image of \tilde{F} . □

Again we obtain Proposition 1.5 by invoking Buchweitz’s equivalence $D_{\text{Sg}}(A) \cong \underline{\text{MCM}}(A)$.

REFERENCES

AZ94 M. Artin and J. J. Zhang, *Noncommutative projective schemes*, Adv. Math. **109** (1994), 228–287.
 APR79 M. Auslander, M. Platzeck and I. Reiten, *Coxeter functors without diagrams*, Trans. Amer. Math. Soc. **250** (1979), 1–46.
 AR90 M. Auslander and I. Reiten, *Graded modules and their completions*, in *Topics in algebra, Part 1 (Warsaw, 1988)*, Banach Center Publications, vol. 26, Part 1 (PWN, Warsaw, 1990), 181–192.

- Bel00 A. Beligiannis, *The homological theory of contravariantly finite subcategories: Auslander–Buchweitz contexts, Gorenstein categories and (co-)stabilization*, *Comm. Algebra* **28** (2000), 4547–4596.
- BN93 M. Bökstedt and A. Neeman, *Homotopy limits in triangulated categories*, *Compositio Math.* **86** (1993), 209–234.
- BP94 A. Bondal and A. Polishchuk, *Homological properties of associative algebras: the method of helices*, *Russian Acad. Sci. Izv. Math.* **42** (1994), 219–260.
- Buc87 R.-O. Buchweitz, *Maximal Cohen–Macaulay modules and Tate cohomology over Gorenstein rings* (1987), 155, unpublished manuscript, available at <https://tspace.library.utoronto.ca/handle/1807/16682>.
- BEH87 R.-O. Buchweitz, D. Eisenbud and J. Herzog, *Cohen–Macaulay modules on quadrics, in Singularities, representation of algebras, and vector bundles (Lambrecht, 1985)* (Springer, Berlin, 1987), 58–116.
- Dyc09 T. Dyckerhoff, *Compact generators in categories of matrix factorizations*, Preprint (2009), arXiv:0904.4713v3.
- Elk74 R. Elkik, *Solution d'équations au-dessus d'anneaux henséliens*, in *Quelques problèmes de modules (Sém. Géom. Anal., École Norm. Sup., Paris, 1971–1972)* (Soc. Math. France, Paris, 1974) Astérisque **16**, 116–132.
- Gab80 P. Gabriel, *Auslander–Reiten sequences and representation-finite algebras*, in *Representation theory, I (Proc. Workshop, Carleton University, Ottawa, Ontario, 1979)*, *Lecture Notes in Mathematics*, vol. 831 (Springer, Berlin, 1980), 1–71.
- OY08 O. Iyama and Y. Yoshino, *Mutation in triangulated categories and rigid Cohen–Macaulay modules*, *Invent. Math.* **172** (2008), 117–168.
- Kel94 B. Keller, *Deriving DG-categories*, *Ann. Sci. École Norm. Sup. (4)* **27** (1994), 63–102.
- Kel98 B. Keller, *On the construction of triangle equivalences*, in *Derived equivalences for group rings*, *Lecture Notes in Mathematics*, vol. 1685 (Springer, Berlin, 1998), 155–176.
- Kel05 B. Keller, *On triangulated orbit categories*, *Doc. Math.* **10** (2005), 551–581 (electronic).
- KR08 B. Keller and I. Reiten, *Acyclic Calabi–Yau categories (with an appendix by Michel Van den Bergh)*, *Compositio Math.* **144** (2008), 1332–1348.
- Kra05 H. Krause, *The stable derived category of a Noetherian scheme*, *Compositio Math.* **141** (2005), 1128–1162.
- LO96 L. Levy and C. Odenthal, *Krull–Schmidt theorems in dimension 1*, *Trans. Amer. Math. Soc.* **348** (1996), 3391–3455.
- Mat89 H. Matsumura, *Commutative ring theory* (Cambridge University Press, Cambridge, 1989).
- Nee96 A. Neeman, *The Grothendieck duality theorem via Bousfield's techniques and Brown representability*, *J. Amer. Math. Soc.* **9** (1996), 205–236.
- Nee01 A. Neeman, *Triangulated categories*, *Annals of Mathematics Studies*, vol. 148 (Princeton University Press, Princeton, NJ, 2001).
- Orl93 D. Orlov, *Projective bundles, monoidal transformations and derived functors of coherent sheaves*, *Russian Acad. Sci. Izv. Math.* **41** (1993), 133–141.
- Orl04 D. Orlov, *Triangulated categories of singularities and D-branes in Landau–Ginzburg models*, *Tr. Mat. Inst. Steklova* **246** (2004), 240–262, *Algebr. Geom. Metody, Svyazi i Prilozh.*
- Orl09a D. Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, in *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin, Vol. II*, *Progress in Mathematics*, vol. 270 (Birkhäuser, Boston, MA, 2009), 503–531 (English summary).
- Orl09b D. Orlov, *Formal completions and idempotent completions of triangulated categories of singularities*, Preprint (2009), arXiv:0901.1859v1.

- RG71 M. Raynaud and L. Gruson, *Critères de platitude et de projectivité. Techniques de ‘platification’ d’un module*, Invent. Math. **13** (1971), 1–89.
- Ric89 J. Rickard, *Morita theory for derived categories*, J. London Math. Soc. (2) **39** (1989), 436–456.
- Rou08 R. Rouquier, *Dimensions of triangulated categories*, J. K-Theory **1** (2008), 193–256.
- Sch03 H. Schoutens, *Projective dimension and the singular locus*, Comm. Algebra **31** (2003), 217–239.
- Ser55 J. P. Serre, *Faisceaux algébriques cohérents*, Ann. of Math. (2) **61** (1955), 197–278.
- Wie98 R. Wiegand, *Local rings of finite Cohen–Macaulay type*, J. Algebra **203** (1998), 156–168.
- Yos90 Y. Yoshino, *Cohen–Macaulay modules over Cohen–Macaulay rings*, London Mathematical Society Lecture Note Series, vol. 146 (Cambridge University Press, Cambridge, 1990).

Bernhard Keller keller@math.jussieu.fr

UFR de Mathématiques, Université Denis Diderot – Paris 7, 2, place Jussieu,
75251 Paris Cedex 05, France

Daniel Murfet murfet@math.uni-bonn.de

Hausdorff Center for Mathematics, Endenicher Allee 62, D-53115 Bonn, Germany

Michel Van den Bergh michel.vandbenbergh@uhasselt.be

Departement WNI, Universiteit Hasselt, Universitaire Campus, Building D, 3590 Diepenbeek,
Belgium