

Astro+: Design, construction, and scientific exploitation of a large-scale massive star spectroscopic database

Klaus Rübke¹, Amparo Marco¹, Ignacio Negueruela², Artemio Herrero^{3,4}, Sergio Simon-Diaz^{3,4}, Hugo Tabernero⁵ and Lee Patrick^{2,6}

¹Departamento de Física, Ingeniería de Sistemas y Teoría de la Señal, Universidad de Alicante, Carretera de San Vicente s/n, E03690, San Vicente del Raspeig, Spain email: klaus.rubke@ua.es

²Departamento de Física Aplicada, Facultad de Ciencias, Universidad de Alicante, Carretera de San Vicente s/n, E03690, San Vicente del Raspeig, Spain

³Instituto de Astrofísica de Canarias, Vía Láctea s/n, E38200, La Laguna, Tenerife, Spain

⁴Universidad de La Laguna, Departamento de Astrofísica, E38206 La Laguna, Tenerife, Spain

⁵ Centro de Astrobiología, CSIC-INTA. Campus ESAC. C. bajo del castillo s/n. E-28 692 Villanueva de la Cañada, Madrid, Spain

 6 School of Physical Sciences, The Open University, Walton Hall, Milton Keynes MK7 6AA, UK

Abstract. Massive stars are the objects that condition the evolution of the interstellar medium by the amount of energy released during their lives and especially by their death as a supernova explosion. All the data provided by the previous and ongoing missions of ground and space telescopes have saturated us with the amount of information, which is no longer easy to process daily by human routines. To this end, we present the development of a massive star spectroscopic interactive database designed for scientific research.

Keywords. Database, Massive-stars, Automatic-programs, parameter-determination.

1. Introduction


We have created a web application Astro+, a dedicated massive star spectra storage site, to have a centralised place for all the spectra available from, 1.the users who want to contribute, 2.the data already available and 3.the data coming from the different new missions. Our application is divided into three main parts, the upload, the search and the science modules. Within the first, we find the upload process, where the user will be able to upload spectra in both FITS and ASCII format.



© The Author(s), 2024. Published by Cambridge University Press on behalf of International Astronomical Union. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

2. Upload method

FITS has different configurations (array, binary data, etc.), and so we allow the user options to set the FITS configuration. On the other hand, the ASCII upload is quite simple, a separate header and spectrum need to be provided, following the standard template required. Once the upload is done, the user can track the upload and correct any errors by using the log we provide. The entered parameters can also be checked, and the user can reprocess the already uploaded file without the need to re-upload the files and then submit the upload for review.



3. Visualization/Search

To inspect the uploaded spectrum, we provide a friendly and easy way to visualize the spectrum, complemented with visual identification using the Aladin API. We add the data from Simbad and Gaia databases through a correlation based on ID or coordinates to check if the star was correctly identified. From Simbad we also adopt the name of the star as the main identification ID to our database.

The search tool, can be used to search through the available spectra in the database by ID, coordinates or by using SQL syntax for more specific star types.

4. Science/AI

By a battery of programs created in Python, based on the best model determination methods (Simón-Díaz & Herrero 2007), (Simón-Díaz et al. 2011), (Simón-Díaz & Herrero 2014) and (Simón-Díaz et al. 2015) on a grid FASTWIND (Santolaya-Rey et al. 1997), (Puls et al. 2005), we can obtain the most critical parameters, such as $V_{\rm rad}$, $V \sin i$, $T_{\rm eff}$ and log g, in a completely automatic way, for early or late massive stars. We are now contrasting preliminary results for Blue/Hot stars with those obtained by (Holgado et al. 2018) from which we are finding errors within the expected, $T_{\rm eff} \sim 1000[\rm K]$ and log $g \sim 0.3[\rm dex]$.

Acknowledgements

This research is partially supported by the Spanish Government Ministerio de Ciencia, Innovación y Universidades under grant PGC2018-093741-B-C21/22 (MICIU/AEI/ FEDER, UE) and the Generalitat Valenciana under grant PROMETEO/2019/041.

Supplementary material

To view supplementary material for this article, please visit http://dx.doi.org/ 10.1017/S1743921322003295.

References

Holgado, G., et al. 2018, A&A, 613, A65

Puls, J., Urbaneja, M. A., Venero, R., Repolust, T., Springmann, U., Jokuthy, A., & Mokiem, M. R. 2005, A&A, 435, 669 Santolaya-Rey, A. E., Puls, J., & Herrero, A. 1997, A&A, 323, 488

- Simón-Díaz, S., Castro, N., Garcia, M., Herrero, A., & Markova, N. 2011, Bulletin de la Societe Royale des Sciences de Liege, 80, 514
- Simón-Díaz, S., & Herrero, A. 2007, A&A, 468, 1063
- —. 2014, A&A, 562, A135
- Simón-Díaz, S., et al. 2015, in Highlights of Spanish Astrophysics VIII, 576-581