

The entropy of polynomial diffeomorphisms of \mathbb{C}^2

JOHN SMILLIE†

Department of Mathematics, Cornell University, Ithaca NY 14853, USA

(Received 21 November 1988)

In this note we answer a question raised by Friedland and Milnor in [FM] concerning the topological entropy of polynomial diffeomorphisms of \mathbb{C}^2

Friedland and Milnor prove that a polynomial diffeomorphism is conjugate to a diffeomorphism of one of three types: affine, elementary or cyclically reduced. The first two families of maps are very simple from a dynamical point of view. The third family contains diffeomorphisms which are dynamically very interesting. The Hénon map is an example of a cyclically reduced diffeomorphism of degree 2.

Topological entropy is most naturally defined for maps of compact spaces. Since \mathbb{C}^2 is not compact, Friedland and Milnor consider the map g , the extension of g to the one-point compactification of \mathbb{C}^2 . They prove that if g is a cyclically reduced diffeomorphism of (algebraic) degree d then the inequality $h(g) \leq \log d$ holds. They raise the question of whether the inequality can be replaced by an equality. We show that it can.

THEOREM *If g is cyclically reduced then $h(g) = \log d$*

The Hénon map has been intensively studied as a map from \mathbb{R}^2 to itself and yet many important problems remain. In particular, the dependence of the entropy of g on the parameter values determining g is quite mysterious. The above result suggests that the dynamics of the Hénon map when considered as a diffeomorphism of \mathbb{C}^2 may be simpler than when considered as a diffeomorphism of \mathbb{R}^2 .

Let $\text{Per}_n(g)$ be the set of periodic points of period n . Let

$$H(g) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log^+ |\text{Per}_n(g)|$$

COROLLARY $H(g) = \log d$

Proof of Corollary This follows by combining the above theorem with the result of [FM] that $h(g) \leq H(g) \leq \log d$.

Friedland and Milnor show that every cyclically reduced polynomial diffeomorphism is conjugate to a composition of generalized Hénon maps of the form $g(x, y) = (y, p(y) - \delta x)$ where p is a polynomial and δ is a nonzero complex number. The degree of g is the degree of p . The degree of a composition of generalized Hénon

† Partially supported by NSF grant #DMS-8803228 and the Sloan Foundation

maps is the product of the degrees of the factors. We begin by proving some basic facts about these maps. The proof of the theorem follows. A version of Lemma 2 first appears in [DN].

LEMMA 1 (see [FM] Lemma 3.4) *For every generalized Hénon map $(x, y) \mapsto (y, z) = (y, p(y) - \delta x)$ there exists a constant κ so that $|y| > \kappa$ implies that either $|z| > |y|$ or $|x| > |y|$ or both.*

We fix the following notation. Let $g = g_1 \circ g_2 \circ \dots \circ g_n$ be a composition of generalized Hénon maps. Let d be the degree of g . Choose κ large enough so that Lemma 1 holds for each g_i . Let

$$\begin{aligned} V^- &= \{(x, y) \mid |y| > \kappa \text{ and } |y| > |x|\} \\ V^+ &= \{(x, y) \mid |x| > \kappa \text{ and } |x| > |y|\} \\ V &= \{(x, y) \mid |x| \leq \kappa \text{ and } |y| \leq \kappa\} \end{aligned}$$

LEMMA 2

- (1) $g(V^-) \subset V^-$
- (2) $g(V^- \cup V) \subset V^- \cup V$
- (3) $g^{-1}(V^+) \subset V^+$
- (4) $g^{-1}(V^+ \cup V) \subset V^+ \cup V$

Proof It suffices to prove each assertion when $g(x, y) \mapsto (y, z)$ is itself a generalized Hénon map.

- (1) Let (x, y) be an element of V^- then $|y| > \kappa$ and $|y| > |x|$. By Lemma 1 $|z| > |y|$ and, since $|y| > \kappa$, we conclude that $|z| > \kappa$. This implies that $g(x, y) = (y, z)$ is in V^- .
- (2) By (1) it suffices to consider the case when (x, y) is an element of V . We will show that $g(x, y) = (y, z)$ is in $V \cup V^-$. Consider two cases. If $|z| \leq \kappa$ then, since $|y| \leq \kappa$, (y, z) is in V . If $|z| > \kappa$ then, since $|y| < \kappa$, we conclude that $|z| > |y|$ so (y, z) is in V^- .
- (3) Let (y, z) be an element of V^+ we want to show that $g^{-1}(y, z) = (x, y)$ is in V^+ . Since $|y| > \kappa$ and $|y| > |z|$ Lemma 1 gives $|x| > |y|$ and, since $|y| > \kappa$ and $|x| > |y|$, we conclude that $|x| > \kappa$. This implies that (x, y) is in V^+ .
- (4) By (3) it suffices to consider the case when (y, z) is an element of V . We will show that (x, y) is in $V^+ \cup V$. If $|x| \leq \kappa$ then since $|y| \leq \kappa$ we conclude that (x, y) is in V . If $|x| > \kappa$ then, since $|y| < \kappa$, we conclude that $|x| > |y|$ and (x, y) is in V^+ .

Notation Let $D_r \subset \mathbb{C}$ be the disk of radius r centered at the origin. Let $\iota: \mathbb{C} \rightarrow \mathbb{C}^2$ be defined by $\iota(z) = (0, z)$. Let $\pi: \mathbb{C}^2 \rightarrow \mathbb{C}$ be defined by $\pi(x, y) = y$.

LEMMA 3 *The set V^- is homotopy equivalent to S^1 . The map $\iota: \partial D_{2\kappa} \rightarrow V^-$ is a homotopy equivalence. The topological degree of the map induced by g on V^- is the algebraic degree of g .*

Proof Let \mathbb{C}_κ be the y -axis. Let $\phi_t(x, y) = ((1-t)x, y)$ for $t \in [0, 1]$. Now $\phi_t(V^-) \subset V^-$, ϕ_0 is the identity on V^- and ϕ_1 is the projection from V^- to $V^- \cap \mathbb{C}_\kappa$. Thus ϕ provides a retraction from V^- to $V^- \cap \mathbb{C}_\kappa$. The set $V^- \cap \mathbb{C}_\kappa$ is the image of $\iota: \mathbb{C} - D_\kappa \rightarrow V^-$. Both $\iota: \mathbb{C} - D_{2\kappa} \rightarrow V^-$ and $\pi: V^- \rightarrow \mathbb{C} - D_\kappa$ are homotopy equivalences.

To prove the last assertion it suffices to consider a single generalized Hénon map $g_t(x, y) \mapsto (y, p(y) - \delta x)$. If we can prove it for a single such map it will follow for a composition of generalized Hénon maps because both the algebraic and homological degrees of generalized Hénon multiply under composition. We compute the degree of the map from V^- to itself by computing the degree of the map $\pi \circ g \circ \iota$. This is an equivalent problem because π and ι are homotopy equivalences. This map is given by $y \mapsto p(y)$. Let d_t be the algebraic degree of g_t , then d_t is the degree of p_t . If L is sufficiently large then the topological degree of the map on $\mathbb{C} - D_L$ induced by p is the degree of the polynomial p . The inclusion $\mathbb{C} - D_k \subset \mathbb{C} - D_L$ is a homotopy equivalence. Thus p has degree d_t on $\mathbb{C} - D_k$.

LEMMA 4 *Let $f : (D, \partial D) \rightarrow (V^- \cup V, V^-)$ be a holomorphic map. Let $\deg(f)$ denote the topological degree of $f : \partial D \rightarrow V^-$. Then $\text{area}(f(D) \cap V) \geq \text{area}(D_k) \cdot \deg(f)$.*

Proof The projection map π sends v to D_k . The induced map from $f(D) \cap V$ to D_k is a proper map and therefore a branched cover. We see that the covering degree is $\deg(f)$ by noting that $\pi f(\partial D)$ wraps $\deg(f)$ times around D_k . Let U be the set obtained from D_k by removing the critical points of the projection and removing arcs connecting the critical points to the boundary of D_k . The area of U is the same as the area of D_k and $f(D) \cap \pi^{-1}U$ consists of $\deg(f)$ components each mapped bijectively onto U by π . Now π does not increase lengths and hence does not increase area so the area of each component is at least $\text{area}(U) = \text{area}(D_k)$. Thus the area of $f(D) \cap V$ is at least $\text{area}(D_k) \cdot \deg(f)$.

Proof of Theorem 1 Let $K^+ \subset \mathbb{C}^2$ be the set of points with bounded forward orbits and let K^- be the set of points with bounded backwards orbits. Let $K = K^+ \cap K^-$. When g is cyclically reduced an argument from [FM] Lemma 3.5 proves that $K^+ \subset V \cup V^-$, $K^- \subset V \cup V^+$ hence $K \subset V$. The same argument shows that all points outside of K are wandering. The set K is compact and is in fact the maximal compact invariant subset of \mathbb{C}^2 .

Friedland and Milnor give $\log d$ as an upper bound for the entropy of $h(g)$. The inequality $h(g) \geq h(g|K)$ is a basic property of entropy. It suffices to prove the lower bound $h(g|K) \geq \log(d)$.

Lemmas 3 and 4 imply that the area of $g^n \iota(D_{2k}) \cap V$ is at least constant d^n . Thus the volume growth, as defined in [Y], of the submanifold $\iota(D_{2k})$ is at least $\log d$. We wish to apply the result of Yomdin ([Y], see also [G]) which says that, for C^∞ maps of compact manifolds, volume growth of submanifolds is a lower bound for entropy. We cannot apply this theorem directly to \mathbb{C}^2 because it is not compact. We cannot apply this theorem directly to K because it is not a manifold and we do not have information on the area of $g^n \iota(D_{2k}) \cap K$. We proceed by an indirect course, we approximate the set K^+ by manifolds with boundary V_n defined below.

Let $d_n(x, y) = \max_{i=0, \dots, n-1} d(g^i(x), g^i(y))$. For X a compact subset of \mathbb{C}^2 we denote by $M(n, \epsilon, X)$ the minimum number of ϵ -balls in the d_n metric needed to cover X . Let $v(n)$ be the area of $g^n \iota(D_{2k}) \cap V$. Let $V_n = V \cap g^{-n}(V)$. Let $v^0(n, \epsilon)$ be the maximum of the area of $g^n \iota(S')$ where S' is $\iota^{-1}(S)$ for S an ϵ -ball of V_n in the d_n

metric If we choose a minimal covering of V_n by ϵ -balls S_i , then the area of $g^n \iota(D_{2\kappa}) \cap V$ is bounded above by the sum of the areas of $g^n \iota(S'_i)$ The sum of areas is bounded above by the number of balls times the maximum area This gives

$$v(n) \leq M(n, \epsilon, V_n) v^0(n, \epsilon)$$

Taking limits gives

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log v(n) \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log M(n, \epsilon, V_n) + \limsup_{n \rightarrow \infty} \frac{1}{n} \log v^0(n, \epsilon)$$

We evaluate $v(n)$ By Lemma 3 the topological degree of the map $g^n \iota$ on $\partial D_{2\kappa}$ is d^n By Lemma 4 we have

$$\text{area}(g^n \iota(D_{2\kappa}) \cap V) \geq \text{constant} \quad \deg(g^n \iota) = \text{constant} \quad d^n$$

Thus the left hand side is greater than or equal to $\log d$ and we have

$$\log d \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log M(n, \epsilon, V_n) + \limsup_{n \rightarrow \infty} \frac{1}{n} \log v^0(n, \epsilon)$$

Taking limits as ϵ goes to zero gives

$$\log d \leq \lim_{\epsilon \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{1}{n} \log M(n, \epsilon, V_n) + \lim_{\epsilon \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{1}{n} \log v^0(n, \epsilon)$$

Yomdin shows ([Y] Theorem 1.8) that

$$\lim_{\epsilon \rightarrow \infty} \limsup_{n \rightarrow \infty} n^{-1} \log v^0(n, \epsilon)$$

is zero for C^∞ maps This result is stated for compact manifolds but it holds in our situation. The following modification is required in the proof A bound of the form B^k on the norm of the first derivative of the k th iterate of the map is needed In our case if B is a bound for the norm of the derivative of $g|V$ then B^k is a bound for the norm of the derivative of $g^k|V_k$

It remains for us to relate the quantity

$$\lim_{\epsilon \rightarrow \infty} \limsup_{n \rightarrow \infty} n^{-1} \log M(n, \epsilon, V_n)$$

to the entropy of $g|K$ Let \bar{V} denote the quotient space $(V \cup V^-)/V^-$ Let m be the point corresponding to V^- We define a metric $\bar{d}(x, y)$ on \bar{V} by the formula

$$\begin{aligned} \bar{d}(x, y) &= \min \{d(x, y), d(x, V^-) + d(y, V^-)\} \\ \bar{d}(x, m) &= d(x, V^-) \end{aligned}$$

Since the set V^- is g invariant, g extends to a continuous map \bar{g} from \bar{V} to itself We have

$$\begin{aligned} h(\bar{g}) &= \lim_{\epsilon \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{1}{n} \log M(n, \epsilon, \bar{V}) \\ &\geq \lim_{\epsilon \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{1}{n} \log M(n, \epsilon, V_n) \\ &\geq \log d \end{aligned}$$

The first equality is the definition of entropy. The second inequality follows because $V_n \subset V$ and if V_n is sufficiently far from $V \cap V^-$ (relative to the size of ε) then the metrics d and \bar{d} are the same when restricted to V_n . Thus an (n, ε) cover of \bar{V} with respect to the \bar{d} metric yields an (n, ε) cover of V_n with respect to d .

By a result of Bowen [B] the entropy of a map is equal to the entropy of the restriction of the map to the nonwandering set. In this case we have $h(\bar{g}) = h(\bar{g}|K^+ \cup \{m\})$ because the nonwandering set is contained in $K^+ \cup \{m\}$. Now

$$h(\bar{g}|K^+ \cup \{m\}) = h(\bar{g}|K^+) + h(\bar{g}|\{m\}) = h(\bar{g}|K^+)$$

On the set K^+ the maps g and \bar{g} are identical. Thus $h(\bar{g}) = h(g|K^+)$. The nonwandering set of $g|K^+$ is contained in $g|K$ so applying Bowen's result again we have $h(g|K^+) = h(g|K)$. Combining these results gives

$$h(g|K) = h(g|K^+) = h(\bar{g}) \geq \log d$$

This completes the proof of the theorem □

REFERENCES

- [B] R. Bowen. Topological entropy and axiom A. *Proc Symp Pure Math* **14** Amer Math Soc (1970), 23-41.
- [DN] R. Devaney & Z. Nitecki. Shift automorphisms in the Hénon map. *Comm Math Phys* **67** (1979), 137-146.
- [FM] S. Friedland & J. Milnor. Dynamical properties of plane polynomial automorphisms. *Ergod Th & Dynam Sys* **9** (1989), 67-99.
- [G] M. Gromov. Entropy, homology and semialgebraic geometry (after Y. Yomdin). *Seminaire N Bourbaki* **663** (1985-86).
- [HO] J. H. Hubbard & R. Oberste-Vorth. Hénon mappings in the complex domain. In preparation.
- [Y] Y. Yomdin. Volume growth and entropy. *Israel J Math* **57** (1987), 285-300.