

## CORRIGENDUM ET ADDENDUM: THE FRATTINI SUBALGEBRA OF A BERNSTEIN ALGEBRA

by JESÚS LALIENA

(Received 5th March 1993)

In a previous paper it is supposed that if  $A$  is a Bernstein algebra, every maximal subalgebra,  $M$ , verifies that  $\dim M = \dim A - 1$ . This is not true in general. Therefore Proposition 2 in this paper is not correct. However other results there, where this assertion was used, are correct but their proofs need some modifications now.

1991 Mathematics subject classification: 17 DXX

### 1. Maximal subalgebras of Bernstein algebras

If we take the commutative algebra  $A$  over a field  $K$  spanned by  $\{e, u_1, u_2, v_1, v_2\}$  such that  $eu_i = 1/2u_i$ ,  $u_1v_1 = u_1$ ,  $u_1v_2 = -u_2$ ,  $u_2v_i = u_i$ ,  $i = 1, 2$  and the other products equals to zero, we have a Bernstein algebra. In this algebra, the subalgebra  $M$  spanned by  $\{e, v_1, v_2\}$  is maximal subalgebra. So if  $M$  is maximal subalgebra of a Bernstein algebra  $A$ , we do not always have  $\dim M = \dim A - 1$ . But we will see that if  $A$  is also genetic then  $\dim M = \dim A - 1$  for every  $M$  maximal subalgebra.

**Lemma 1.** *Let  $A$  be a Bernstein algebra,  $e$  a nonzero idempotent in  $A$  such that  $A = Ke \oplus U_e \oplus V_e$ , and  $M$  a maximal subalgebra of  $A$ . Then  $U_e^2, U_e^3, (U_e^2)^2 \subseteq M$ .*

**Proof.** Let  $N = U_e + U_e^2$ . We have that  $B = Ke + N$  is a subalgebra of  $A$  and a Bernstein algebra. From [1] it is known that a Bernstein algebra  $A$  with  $A^2 = A$  is genetic. Since  $B^2 = B$ , we have that  $N$  is nilpotent, and from [3]  $F(N) = N^2$ . But  $N^2 = U_e^2 + U_e^3 + (U_e^2)^2$  is an ideal of  $A$  because of [4] (or checking it directly). Therefore, using [3], we have  $N^2 \subseteq F(A)$ , that is  $U_e^2, U_e^3, (U_e^2)^2 \subseteq M$  for every maximal subalgebra  $M$ .

**Lemma 2.** *Let  $M$  be a maximal subalgebra of a Bernstein algebra  $A$  and let  $e$  be an idempotent in  $M$ . Then either  $V_e \subseteq M$  or  $U_e \subseteq M$ .*

**Proof.** Clearly  $M = Ke + U'_e + V'_e$  with  $U'_e \subseteq U_e$ ,  $V'_e \subseteq V_e$ . Now  $M + U_e = M$  or  $M + U_e \subseteq A$ . The former implies that  $U'_e = U_e$ ; the latter implies that  $V'_e = V_e$ .

Now it is easy to prove the following results.

**Theorem 3.** Let  $A$  be a genetic Bernstein algebra. If  $M$  is a maximal subalgebra of  $A$ , then  $\dim M = \dim A - 1$ . Therefore a vector subspace  $M$  of  $A$  is a maximal subalgebra if and only if

either (i)  $M = \text{Ker } w$ ,

or (ii)  $M$  has a nonzero idempotent  $e$  such that  $M$  is one of the following subalgebras

(a)  $M = K.e \oplus U_e \oplus V'_e$  with  $V'_e \leq V_e$  such that  $\dim V'_e + 1 = \dim V_e$  and  $U_e^2 \leq V'_e$ ; in this case  $M$  is an ideal,

(b)  $M = K.e \oplus U'_e \oplus V_e$  with  $U'_e \leq U_e$ ,  $\dim U'_e + 1 = \dim U_e$ ,  $U_e V_e + V_e^2 \leq U'_e$ .

**Proposition 4.** Let  $A$  be a Bernstein algebra and  $M$  a vector subspace of  $A$ . Then  $M$  is a maximal subalgebra if  $M$  is one of the following subspaces:

(i)  $M = \text{Ker } w$ ,

(ii)  $M$  has a nonzero idempotent  $e$  such that

(a)  $M = K.e \oplus U_e \oplus V'_e$  with  $V'_e$  such that  $\dim V'_e + 1 = \dim V_e$  and  $U_e^2 \leq V'_e$ . In this case  $M$  is an ideal.

(b)  $M = K.e \oplus U'_e \oplus V_e$  with  $U'_e \leq U_e$ ,  $\dim U'_e + 1 = \dim U_e$ ,  $U'_e V_e + V_e^2 \leq M$

(c)  $M = K.e \oplus U'_e \oplus V_e$  with  $U'_e \leq U_e$ , such that  $\dim U'_e + 1 < \dim U_e$ ,  $U'_e V_e + V_e^2 \leq U'_e$ .

**Theorem 5.** Let  $A$  be a Bernstein algebra. Then  $F(A)$  is an ideal.

**Proof.** Let  $N = U_e + U_e^2$ ; then from the proof of Lemma 1,  $N^2 \leq F(A)$ . Since  $F(A/N) = 0$ , then  $F(A) \leq N$ . Now suppose that  $AF(A) \not\leq F(A)$ . Then there is a maximal subalgebra  $M$  of  $A$  with  $AF(A) \not\leq M$ . But  $AF(A) \leq AN \leq N$ , so  $N \not\leq M$  and  $A = M + N$ . Thus  $AF(A) = MF(A) + NF(A) \leq M + N^2 \leq M$ , a contradiction.

#### REFERENCES

1. A. N. GRISHKOV, On the Genetic Property of Bernstein algebras, *Soviet Math. Dokl.* **35** (1987), 489–492.
2. J. LALIENA, The Frattini subalgebra of a Bernstein algebra, *Proc. Edinburgh Math. Soc.* **34** (1992), 397–403.
3. D. TOWERS, A Frattini theory for algebras, *Proc. London Math. Soc.* (3) **27** (1973), 440–462.
4. A. WÖRZ-BUSEKROS, Algebras in Genetics (Springer-Verlag. Lectures Notes in Biomathematics **36**, Berlin Heidelberg 1980).

DEPARTAMENTO DE MATEMÁTICAS  
UNIVERSIDAD DE LA RIOJA  
26004-LOGROÑO  
SPAIN