

Welfare assessment of captive Asian elephants (*Elephas maximus*) and Indian rhinoceros (*Rhinoceros unicornis*) using salivary cortisol measurement

A Menargues^{*†}, V Urios[†] and M Mauri[‡]

[†] Estacion Biologica Terra Natura, Fundacion Terra Natura-CIBIO, Universidad de Alicante, Apdo correos 99, E-03080, Alicante, Spain

[‡] Hospital General Universitario de Alicante, E-03010, Alicante, Spain

* Contact for correspondence and requests for reprints: a.menargues@ua.es

Abstract

The measurement of salivary cortisol allows non-invasive assessment of welfare in captive animals. We utilised this technique to test the effect of zoo opening on six Asian elephants and two Indian rhinoceros at the Terra Natura Zoological Park, Alicante, Spain, during pre-opening, opening and post-opening periods. Salivary cortisol concentrations were found to be significantly higher during the opening period than during pre- and post-opening periods for both species. This method could prove a useful tool in monitoring the success of decisions taken to improve the welfare of captive animals.

Keywords: animal welfare, Asian elephant, Indian rhinoceros, management, salivary cortisol, stress

Introduction

One of the most important objectives of zoological parks is maintaining the psychological and physiological well-being of animals in their care. Changes in the animals' life such as acute environmental or social upheaval (eg introduction to novel environments and new group formation) can be extremely stressful (Soltis *et al* 2003). Although no widely-held definition exists, stress generally refers to a variety of responses to internal or external stimuli (stressors) that modify the homeostasis of an individual. These stimuli can be physical, physiological, behavioural or psychological factors (Dantzer & Mormede 1983; Stratakin & Chrousos 1995). Immediately following exposure to a stressor, activation of the hypothalamic-pituitary-adrenal axis occurs and glucocorticoids are released from the adrenal cortex (Munck *et al* 1984). Thus, corticosteroids and glucocorticoids are secreted in response to stress and these can provide information regarding the physiological well-being of animals (Whitten *et al* 1998).

Plasma cortisol has been used to evaluate animal stress responses to capture and translocation (Morton *et al* 1995; Denhard *et al* 2001), disturbances within their environment (Zoldag *et al* 1983; Powell *et al* 2006) or psychogenic stress (Haemisch 1990) but capture and handling, along with blood sample collection via venipuncture, can be stressors themselves, leading to increased plasma cortisol concentration (Cook *et al* 1973; Dawson & Howe 1983; Sabatino *et al* 1991; Whitten *et al* 1998; Schoech *et al* 1999; Romero & Romero 2002).

Non-invasive glucocorticoid collection techniques allow accurate monitoring of stress without the bias of capture-

induced or disturbance-induced increases in cortisol levels (Hamilton & Weeks 1985; Harper & Austad 2000; Millsbaugh *et al* 2001; Washburn & Millsbaugh 2002). Measurement of glucocorticoids via analysis of faecal samples has been used to monitor stress responses in different species (Wasser *et al* 2000; Von der Ohe & Servheen 2002; Shepherdson *et al* 2004; McKenzie & Deane 2005; Lane 2006). Another such method is saliva collection which can be used with habituated or laboratory-housed animals. The use of salivary cortisol levels to monitor stress has been validated in humans (see Kirchbaum & Hellhammer 1994 for a review) and other species such as dogs (*Canis familiaris*) (Vincent & Michell 1992; Beerda *et al* 1996), rhesus monkeys (*Macaca mulatta*) (Boyce *et al* 1995), squirrel monkeys (*Saimiri sciureus*) (Fuchs *et al* 1997), and tree shrews (*Tupaia belangeri*) (Ohl *et al* 1999). This technique has also been used in Indian rhinoceros (*Rhinoceros unicornis*) (Kuckelkorn & Dathe 1990) and in Asian elephants (*Elephas maximus*) (Dathe *et al* 1992). Studies in pigs (*Sus scrofa*) have revealed that salivary cortisol is related to free plasma cortisol and represents approximately 10% of the concentration in plasma (Haeckel 1989; Parrot *et al* 1989). Saliva samples are easy to collect and store and sampling can be performed frequently, which is of particular importance as it allows acute stress to be monitored by measuring short-term changes in cortisol levels (Queyras & Carossi 2004).

In this study we used salivary cortisol to assess the degree to which zoo opening affected stress levels of Asian elephants and Indian rhinoceros and also sought to assess

the viability of this method in terms of monitoring animal welfare in captivity and providing a tool for testing the success of management decisions designed specifically to improve or maintain welfare.

Materials and methods

Location

The study was conducted at Terra Natura Zoological Park, Alicante, Spain. Salivary cortisol concentrations were recorded from Asian elephants and Indian rhinoceros for three separate timeperiods: (1) pre-opening, ie during the month prior to opening (15th February–14th March 2005), at which time animals were confined to their cages and internal courtyards; (2) opening, ie when the zoo opened its doors for the very first time and throughout the month that followed (15th March–14th April 2005), during which time the animals spent their days in dry meadows and had direct visual contact with not only zoo visitors but also the final remnants of the construction work which had yet to be completed and (3) post-opening, ie the following month (15th April–14th May 2005), during which time animals were found in their prairies and were in direct visual contact with visitors. By this point the construction work, which had consisted of excavation, meadow and artificial lake construction, tree planting, power line, water and sewage line installation and the construction of ancillary buildings (eg stables, offices, restaurants, shops, etc), was completed.

Animals

Six adult female Asian elephants, ranging from 23 to 45 years of age and two juvenile female Indian rhinoceros aged three and five years, were studied. Three of the elephants (Tania, Jasmin and Motki) arrived at Terra Natura Zoological Park in August 2004 from Austria (Gänserndorf Safari Park) whilst the others (Petita, Baby and Kaisoso) arrived in December 2004, having previously resided at the Vergel Safari Park (Vergel, Alicante, Spain). The rhinoceros arrived in March 2003: Nisha from Stuttgart Zoo and Shiwa from Munich Zoo. At night, each rhinoceros was housed in an individual cage measuring $4 \times 5 \times 2.50$ m (length \times width \times height) and during the day (0900 to 1800h), they ranged freely in a 2,000 m² meadow. One of the elephants (Baby) was housed individually in a cage measuring $5 \times 6 \times 3.50$ m during the night due to conflict with other elephants, while the rest of the herd was housed in two cages measuring $5 \times 8 \times 3.50$ m; one with a group of two individuals (Kaisoso and Petita) and the other with a group of three (Tania, Modki and Jasmin). During the day time was spent in three different 3,000 m² dry meadows. The opening period represented the first time these animals had encountered the meadows as they had been confined to the internal courtyards since arriving. Animals were fed on a diet of oats, branches, fruit and vegetables with *ad libitum* water. All the animals were trained and managed using free-contact by keepers with whom they became familiar. This made it possible to use saliva sampling as a non-invasive method for assessing animal welfare.

Sample collection

Saliva samples were taken on a daily basis at 0730h during February, March, April and May 2005. Samples were always collected at this time to avoid variations in saliva cortisol concentration, ie significantly large variation in concentrations can be seen during early morning, within relatively short time intervals. Samples were collected using the Salivette® kit (Sarstedt, AG & Co, Nümbrecht, Germany) and centrifuged at 2,000 rpm for two minutes at 15°C. The eluted saliva was stored at –20°C until assaying.

Hormone analysis

Salivary cortisol was measured in duplicate, using a modification of the solid-phase RIA (Coat-A-Count®, Siemens Medical Solutions Diagnostics, Los Angeles, CA, USA) and the tubes were counted on a Packard Cobra Auto Gamma Counter (Auto-Gamma® 5000 series, Cobra 5005, Packard Instruments Company, Meriden, CT, USA). A minimum of 0.4 ml of saliva was required for the duplicate assay. Modifications were made in the form of increasing the volume of the sample to 200 µl, in the standard curve, through 1:10 dilutions of the standards and increasing the incubation period to 24 h to increase assay sensitivity (López-Mondejar *et al* 2006).

Assay validation

The saliva cortisol assays of the Asian elephants and Indian Rhinoceros were validated through demonstration of the correlation between serial dilutions of pooled saliva and the standard curve (Meyer *et al* 2004). Analytical sensitivity, or minimum detection limit, was calculated by interpolation of the mean minus two standard deviations of ten replicates of the zero calibrator. Assay precision was assessed by calculating intra- and inter-assay coefficients of variation (CVs) of the percentage bound of the internal controls. Specificity was extracted from the information supplied by the manufacturer's kit.

Statistical analysis

Data were analysed by repeated measures analysis of variance (ANOVA), using the cortisol concentration data obtained from each individual in order to test for differences between individual elephants and periods. The same procedure was carried out for rhinoceros. The Huynh and Feldt correction was used in Mauchly's Test of Sphericity in elephant analysis. The Bonferroni multiple comparison test was used to test for pairwise differences in mean cortisol values among pre-opening, opening and post-opening periods, for both species. Statistical analyses were conducted using SPSS, version 13.0.


Results

Assay validation


The dilution curve for the pooled Indian rhinoceros saliva correlated positively with that of the Asian elephants ($r = 0.997$, $df = 6$, $P < 0.001$). The dilution curve for the pooled saliva correlated positively with the standard curve for the rhinoceros ($r = 0.962$, $df = 6$, $P < 0.001$), and the elephants ($r = 0.946$, $df = 6$, $P < 0.001$).

Figure 1

Mean salivary cortisol concentration during pre-opening, opening and post-opening period in (a) Asian elephants.



Mean salivary cortisol concentration during pre-opening, opening and post-opening period in (b) Indian rhinoceros.



Sensitivity was 0.82 nmol l⁻¹ of cortisol, intra- and inter-assay coefficients of variation were 8% (n = 10) for Asian elephants and 5.8% (n = 10) for Indian rhinoceros. The cortisol antibody crossreactivity was: 100% with cortisol, 11.4% with 11-deoxycortisol, 2.3% with prednisone, 1% with corticosterone and cortisone, < 1% with aldosterone, 11-deoxycorticosterone, dexamethasone, oestriol, oestrone, flumethasone, methotrexate, pregnenolone, progesterone, tetrahydrocortisol and triamcinolone.

Hormone analysis

The mean concentration of salivary cortisol in both elephants and rhinoceros was increased during the opening period, compared to pre- and post-opening periods (Figure 1). Table 1 shows that for both species the lowest concentration of salivary cortisol was obtained during the post-opening period.

Repeated measures ANOVA revealed significant changes in the cortisol concentration in Asian elephants ($F_{9,203, 377.341} = 5.964$, $P < 0.001$) and Indian rhinoceros ($F_{2, 82} = 4.694$, $P = 0.012$) between the pre-opening, opening and post-opening periods. Bonferroni multiple comparison tests (Figure 2) revealed that cortisol concentrations during the opening period were significantly higher than during pre- and post-opening periods for both species, but no significant differences were seen between pre- and post-opening periods' cortisol concentration for both species.

Discussion


Glucocorticoids reflect physical and behavioural stress situations and their concentration in blood plasma is quantitatively related to the degree of stress (Dathe *et al* 1992). Saliva sampling is a preferred, non-invasive means of assessing free

Table 1 Means of salivary cortisol of all Asian elephant and Indian rhinoceros individuals during the zoo pre-opening, opening and post-opening periods.

Species	Animal	Period	Mean (nmol l ⁻¹)	SD	n
<i>Elephas maximus</i>	Baby	Pre-opening	2.49	0.39	27
		Opening	4.59	1.78	29
		Post-opening	1.97	0.87	29
		Total	3.03	1.64	85
	Jasmin	Pre-opening	2.36	0.50	27
		Opening	3.75	1.57	29
		Post-opening	1.78	0.64	29
		Total	2.64	1.32	85
	Kaisoso	Pre-opening	2.68	0.65	27
		Opening	4.95	1.90	29
		Post-opening	1.73	0.92	29
		Total	3.13	1.87	85
	Modki	Pre-opening	2.33	0.37	27
		Opening	4.17	2.15	29
		Post-opening	1.85	0.62	29
		Total	2.80	1.65	85
	Petita	Pre-opening	2.71	0.57	27
		Opening	4.35	1.76	29
		Post-opening	2.08	0.68	29
		Total	3.06	1.49	85
	Tania	Pre-opening	2.17	0.39	27
		Opening	3.32	1.38	29
		Post-opening	2.26	0.80	29
		Total	2.59	1.08	85
	Total	Pre-opening	2.46	0.49	189
		Opening	4.26	1.77	196
		Post-opening	1.96	0.73	210
		Total	2.87	1.51	595
<i>Rhinoceros unicornis</i>	Nisha	Pre-opening	2.15	0.59	27
		Opening	4.18	1.02	29
		Post-opening	2.36	0.99	29
		Total	2.91	1.28	85
	Shiwa	Pre-opening	2.24	0.59	27
		Opening	3.47	0.89	29
		Post-opening	1.71	0.45	29
		Total	2.48	1.00	85
	Total	Pre-opening	2.22	0.49	54
		Opening	3.69	0.80	56
		Post-opening	1.89	0.56	60
		Total	2.59	1.00	170

Figure 2

Vertical bars indicate means of cortisol concentration during pre-opening, opening and post-opening period. Bars with the same letter do not differ significantly.



cortisol concentration (Boyce *et al* 1995; Tiefenbacher *et al* 2003) because, depending on the collection method used, it is less stressful to animals than collecting blood samples (Cross *et al* 2004). Salivary cortisol concentration can be used for assessing plasma cortisol concentration since it accurately reflects the concentration of the biologically-active free fraction of the hormone in plasma (Haeckel 1989) and represents approximately 10% of the concentration in plasma (Haeckel 1989; Parrot *et al* 1989).

Our aim in this study was to assess a potential tool for indicating the welfare of animals during zoo openings. This was carried out through measurement of salivary cortisol concentrations in Asian elephants and Indian rhinoceros. This non-invasive method of examining stress responses in these species was selected because these animals were familiarised with their keepers and trained and managed using free-contact.

One question that needed to be addressed was the amount of cortisol concentration that could be attributed to stress-induction in the animal. The term 'stress' is generally considered to be a reference to the negative consequences of an animal's failure to cope with factors in its environment (Wielebnowski *et al* 2002) and increases in cortisol concentration are normally established as a negative experience (Norcross & Newman 1999; Cross *et al* 2004; Dehnhard 2007). That said, not all stressors have negative impacts on animal health (Moodie & Chamove 1990) and, in certain circumstances, stress is a necessity in ensuring survival and allowing adaptation to environmental change (Wielebnowski *et al* 2002).

So-called 'negative stress' can be identified when animals appear to incur biological costs such as impaired reproduction (Swaigood 2007) or immune function (Siegel *et al* 1987; Herbert & Cohen 1993) from responding to stressors (Moberg 2000). Due to the limitations imposed by a study only lasting three months, we were unable to confirm whether or not the cortisol concentrations obtained in these three periods could have biological costs. Thus, in order to determine which concentration represented 'negative stress', it was necessary to assess the normal range of cortisol concentration for these species; not only to establish a baseline range of salivary cortisol to detect deviations which we could attribute to stressful situations, but to also monitor their evolution in individuals and in groups (Dathe *et al* 1992). Furthermore, in mammals, the production of basal cortisol generally peaks at the beginning of daily activity, which can be explained by a circadian rhythmicity that has evolved to help the organism predict daily environmental changes, such as food seeking and consumption (Moore-Ede & Sulzman 1977). To ensure results were not influenced by this circadian rhythmicity, samples were always collected at the same time (0730h).

Once the normal cortisol concentration range of both species had been established, we observed that during the opening period, salivary cortisol concentrations of both species increased significantly, compared to pre- and post-opening periods. We considered these higher values to be 'negative stress' and they may have been attributable to the continuation of building work, causing increased disturbance during this period. Some studies have documented negative responses of giant pandas (*Ailuropoda melanoleuca*) to construction noise (Powell *et al* 2006). In

addition, during this opening period, the animals were introduced into their dry meadows for the first time whereas previously they had been confined to their cages and internal courtyards. Furthermore, during the opening period, animals had their first direct visual contact with visitors that year. Several studies have shown correlations between increased cortisol concentration in animals and zoo visitors (Glatston *et al* 1984; Thompson 1989; Carlstead *et al* 1999) as well as with changes in animals' enclosures (Ohl *et al* 1999; Cross *et al* 2004).

After the opening period, salivary cortisol values decreased, due to possible habituation to their new meadows and visitors, during the month following the opening. Furthermore, Bonferroni multiple comparison tests reveal that cortisol levels were slightly lower during the post-opening period than during the pre-opening period. This suggests animal installation conditions improved after the zoo opening.

Moreover, we found high inter-individual variation in salivary cortisol concentrations which reflects an individual's ability to cope with the captive situation (Zayan 1991; Carlstead & Shepherdson 2000). An intra-specific metabolic variation has been noted in other species, such as cattle and other ruminants (Palme *et al* 1999, 2000; Morrow *et al* 2002), elephants (Stead *et al* 2000), rhinoceros (Brown *et al* 2001), clouded leopard (Wielebnowski *et al* 2002) and other mammals and avian species (Wasser *et al* 2000). In spite of these inter-individual differences, it was observed in all individuals that higher cortisol concentration values were found in the opening period; this time being particularly stressful for all animals concerned.

Some studies have documented positive responses to environmental enrichment (Wells & Hepper 2000; Wells 2004, 2005; Graham *et al* 2005; Mallapur *et al* 2005; Wells *et al* 2007) and cortisol concentration could be used to test the success of these enrichment techniques (Schapiro *et al* 1993; Carlstead & Shepherdson 2000; de Groot *et al* 2000; Ernst *et al* 2006). We propose that this non-invasive method of measuring salivary cortisol in Asian elephants and Indian rhinoceros could be a useful tool for zoo management and an objective measure to test environmental enrichment techniques and improve animal welfare. This method could be equally applicable to other captive animals which are trained for this technique and have direct contact with keepers.

Conclusions

The findings of this study indicate that measurement of salivary cortisol concentrations could be a useful tool for assessing animal welfare and for testing the suitability of the captive environment.

Acknowledgements

We are extremely grateful to Terra Natura Foundation that has financed this study, to the zoo's veterinarians, Manuel Lopez and Rocio Canales for their collaboration and to animal keepers of Terra Natura Zoo for their assistance in taking animal samples. We also want to thank the University General Hospital of Alicante for allowing us access to the hormone laboratory in order to analyse saliva samples and to

Rocio Alfayate and Mara Gozalvez. Finally we want to thank Alvaro Soutullo, Luis Cadahia, Ruben Limiñana, Miguel Gallardo and Ramon Garcia Navarro for the assessment of the statistical analysis and the revision of the translation.

References

- Beerda B, Schilder MBH, Janssen NSCRM and Mol JA** 1996 The use of saliva cortisol, urinary cortisol, and catecholamine measurements for a non-invasive assessment of stress responses in dogs. *Hormones and Behaviour* 30: 272-279
- Boyce WT, Champoux M, Suomi SJ and Gunnar MR** 1995 Salivary cortisol in nursery-reared rhesus monkeys: reactivity to peer interactions and altered circadian activity. *Developmental Psychobiology* 28: 257-267
- Brown JL, Bellem AC, Fouraker M, Wildt DE and Roth TL** 2001 Comparative analysis of gonadal and adrenal activity in the black and white rhinoceros in North America by non-invasive endocrine monitoring. *Zoo Biology* 20: 463-486
- Carlstead K and Shepherdson D** 1994 Effects of environmental enrichment on reproduction. *Zoo Biology* 13: 447-459
- Carlstead K, Mellen J and Kleiman DG** 1999 Black rhinoceros (*Diceros bicornis*) in US Zoos: I Individual behaviour profiles and their relationship to breeding success. *Zoo Biology* 18: 17-34
- Carlstead K and Shepherdson D** 2000 Alleviating stress in zoo animals with environmental enrichment. In: Moberg GP and Mench JA (eds) *The Biology of Animal Stress* pp 337-354. CABI Publishing: New York, USA
- Cook DM, Kendall JW, Greer MA and Kramer RM** 1973 The effect of acute or chronic ether stress on plasma ACTH concentration in the rat. *Endocrinology* 93: 1019-1024
- Cross N, Pines MK and Rogers LJ** 2004 Saliva sampling to assess cortisol levels in unrestrained common marmosets and the effect of behavioural stress. *American Journal of Primatology* 62: 107-114
- Dantzer R and Mormede P** 1983 Stress in farm animals: a need for reevaluation. *Journal of Animal Science* 57: 6-18
- Dathe HH, Kuckelkorn B and Minnemann D** 1992 Salivary cortisol assessment for stress detection in the Asian Elephant (*Elephas maximus*): a pilot study. *Zoo Biology* 11: 285-289
- Dawson A and Howe PD** 1983 Plasma corticosterone in wild starlings (*Sturnus vulgaris*) immediately following capture and relation to bodyweight during the annual cycle. *General and Comparative Endocrinology* 51: 303-308
- de Groot J, de Jong IC, PELLE IT and Colas JM** 2000 Immunity in barren and enriched housed pigs differing in baseline cortisol concentration. *Physiology and Behavior* 71: 217-223
- Dehnhard M, Clauss M, Lechner-Doll M, Meyer HHD and Palme R** 2001 Non-invasive monitoring of adrenocortical activity in roe deer (*Capreolus capreolus*) by measurement of fecal cortisol metabolites. *General and Comparative Endocrinology* 123: 111-120
- Dehnhard M** 2007 Characterisation of the sympathetic nervous system of Asian (*Elephas maximus*) and African (*Loxodonta africana*) elephants based on urinary catecholamine analysis. *General and Comparative Endocrinology* 151: 274-284
- Ernst K, Tuchscherer M, Kanitz E, Puppe B and Manteuffel G** 2006 Effects of attention and rewarded activity on immune parameters and wound healing in pigs. *Physiology and Behavior* 89: 448-456
- Fuchs E, Kirschbaum C, Benisch D and Bieser A** 1997 Salivary cortisol: a non-invasive measure of hypothalamo-pituitary-adrenocortical activity in the squirrel monkey, *Saimiri sciureus*. *Laboratory Animals* 31: 306-311
- Glatston AR, Geilvoet-Soetman E, Hora-Peck E and van Hoff J** 1984 The influence of the zoo environment on social behaviour of groups of cotton-top tamarin, *Sanguinus oedipus oedipus*. *Zoo Biology* 3: 241-253

- Graham L, Wells DL and Hepper PG** 2005 The influence of olfactory stimulation on the behaviour of dogs housed in a rescue shelter. *Applied Animal Behaviour Science* 91: 143-153
- Haeckel R** 1989 Application of saliva in laboratory medicine. Report on the workshop conference. *Journal of Clinical Chemistry and Clinical Biochemistry* 27: 233-252
- Haemisch A** 1990 Coping with social conflict, and short term changes of plasma cortisol titers in familiar and unfamiliar environments. *Physiology and Behaviour* 47: 1265-1270
- Hamilton GD and Weeks Jr HP** 1985 Cortisol and aldosterone comparisons of cottontail rabbits collected by shooting, trapping, and falconry. *Journal of Wildlife Diseases* 31: 40-42
- Harper JM and Austad SN** 2000 Fecal glucocorticoids: a non-invasive method of measuring adrenal activity in wild and captive rodents. *Physiological and Biochemical Zoology* 73: 12-22
- Herbert TB and Cohen S** 1993 Stress and immunity in humans: a meta-analytic review. *Psychosomatic Medicine* 55: 364-379
- Kirschbaum C and Hellhammer DH** 1994 Salivary cortisol in psychoneuroendocrine research: recent developments and applications. *Psychoneuroendocrinology* 19: 313-333
- Kuckelkorn B and Dathe HH** 1990 Trächtigkeits-diagnose beim Panzernashorn (*Rhinoceros unicornis*) anhand von Progesteronbestimmungen im Speichel. *Der Zoologische Garten NF* 60: 333-340. [Title translation: Pregnancy diagnosis in the Asian rhino on the basis of progesterone determination in saliva]
- Lane** 2006 Can non-invasive glucocorticoid measures be used as reliable indicators of stress in animals. *Animal Welfare* 15: 331-342
- López-Mondejar P, Fuentes MA, Mauri M, Mora A, Pérez Soto M, Vargas F and Martín Hidalgo A** 2006 Determinación de cortisol salivar en el diagnóstico de la enfermedad de Cushing pediátrico. *Anales de Pediatría* 64: 270-272. [Title translation: Salivary cortisol determination in pediatric Cushing illness diagnosis]
- Mallapur A, Sinha A and Waran N** 2005 Influence of visitor presence on the behaviour of captive lion-tailed macaques (*Macaca silenus*) housed in Indian zoos. *Applied Animal Behaviour Science* 94: 341-352
- McKenzie S and Deane EM** 2005 Faecal corticosteroid levels as an indicator of well-being in the tammar wallaby, *Macropus eugenii*. *Comparative Biochemistry and Physiology* 140: 81-87
- Meyer JM, Walker SL, Freeman EW, Steinetz BG and Brown JL** 2004 Species and fetal gender effects on the endocrinology of pregnancy in elephants. *General and Comparative Endocrinology* 138: 263-270
- Millspaugh JJ, Woods RJ, Hunt KE, Raedeke KJ, Brundige GC, Washburn BE and Wasser SK** 2001 Using fecal glucocorticoid assays to study the physiological stress response of elk. *Wildlife Society Bulletin* 29: 899-907
- Moberg GP** 2000 Biological response to stress: implications for animal welfare. In: Moberg GP and Mench JA (eds) *The Biology of Animal Stress* pp 1-22. CABI Publishing: New York, USA
- Moodie EM and Chamove AS** 1990 Brief threatening events are beneficial for captive tamarins. *Zoo Biology* 9: 275-286
- Moore-Ede MC and Sulzman FM** 1977 The physiological basis of circadian time-keeping in primates. *Physiologist* 20: 17-25
- Morrow CJ, Kolver ES, Verkerk GA and Matthews L** 2002 Fecal glucocorticoid metabolites of adrenal activity in dairy cattle. *General and Comparative Endocrinology* 126: 229-241
- Morton DJ, Anderson E, Foggin CM, Kock MD and Tiran EP** 1995 Plasma cortisol as an indicator of stress due to capture and translocation in wildlife species. *Veterinary Record* 136: 60-63
- Munck A, Guyre PM and Holbrook NJ** 1984 Physiological functions of glucocorticoids in stress and their relation to pharmacological effects. *Endocrine Reviews* 5: 25-44
- Norcross JL and Newman JD** 1999 Effects of separation and novelty on distress vocalizations and cortisol in the common marmoset (*Callithrix jacchus*). *American Journal of Primatology* 47: 209-222
- Ohi F, Kirschbaum C and Fuchs E** 1999 Evaluation of hypothalamo-pituitary-adrenal activity in the tree shrew (*Tupaia belangeri*) via salivary cortisol measurement. *Laboratory Animals* 33: 269-274
- Palme R, Robia C, Messmann S, Hofer J and Mostl E** 1999 Measurement of faecal cortisol metabolites in ruminants: a non-invasive parameter of adrenocortical function. *Wiener Tierärztliche Monatsschrift* 86: 237-241
- Palme R, Robia C, Baumgartner W and Mostl E** 2000 Transport stress in cattle as reflected by an increase in faecal cortisol metabolite concentrations. *The Veterinary Record* 146: 108-109
- Parrot RF, Misson BH and Baldwin BA** 1989 Salivary cortisol in pigs following adrenocorticotropic hormone stimulation: comparison with plasma levels. *British Veterinary Journal* 145: 362-366
- Powell DM, Carlstead K, Tarou LR, Brown JL and Monfort SL** 2006 Effects of construction noise on behaviour and cortisol levels in a pair of captive giant pandas (*Ailuropoda melanoleuca*). *Zoo Biology* 25: 391-408
- Queyras A and Carossi M** 2004 Non-invasive techniques for analysing hormonal indicators of stress. *Ann Ist Super Sanità* 40: 211-221
- Romero LM and Romero RC** 2002 Corticosterone responses in birds: the importance of rapid initial sampling. *Condor* 104: 129-135
- Sabatino F, Masoro EJ, McMahan A and Kuhn RW** 1991 Assessment of the role of glucocorticoid system in aging process and in the action of food restriction. *Journal of Gerontology* 46: 171-179
- Schapiro S, Bloomsmith MA, Kessel AL and Shively CA** 1993 Effects of enrichment and housing on cortisol response in juvenile rhesus monkeys. *Applied Animal Behaviour Science* 37: 251-263
- Schoech SJ, Ketterson ED and Nolan Jr V** 1999 Exogenous testosterone and the adrenocortical response in dark-eyed junco. *Auk* 116: 64-72
- Shepherdson DJ, Carlstead KC and Wielebnowski NC** 2004 Cross-institutional assessment of stress responses in zoo animals using longitudinal monitoring of faecal corticoids. *Animal Welfare* 13: 105-115
- Siegel RA, Ducker EM, Pahnke U and Wuttke W** 1987 Stress-induced changes in cholecystokinin and substance P concentrations in discrete regions of the rat hypothalamus. *Neuroendocrinology* 46: 75-81
- Soltis J, Wegner FH and Newman JD** 2003 Adult cortisol response to immature offspring play in captive squirrel monkeys. *Physiology and Behaviour* 80: 217-223
- Stead SK, Meltzer DGA and Palme R** 2000 The measurement of glucocorticoid concentrations in the serum and faeces of captive African elephants (*Loxodonta africana*) after ACTH stimulation. *Journal of the South African Veterinary Medical Association* 71: 192-196
- Stratakis CA and Chrousos GP** 1995 Neuroendocrinology and pathophysiology of the stress system: In stress, basic mechanism and clinical implications. *Annals of the New York Academy of Sciences* 771: 1-18
- Swaisgood R** 2007 Current status and future directions of applied behavioural research for animal welfare conservation. *Applied Animal Behaviour Science* 102: 139-162
- Thompson VD** 1989 Behavioural response of 12 ungulate species in captivity to the presence of humans. *Zoo Biology* 8: 275-297
- Tiefenbacher S, Lee B, Meyer JS and Spelman RD** 2003 Non-invasive technique for repeated sampling of salivary free cortisol in awake, unrestrained squirrel monkeys. *American Journal of Primatology* 60: 69-75
- Vincent IC and Michell AR** 1992 Comparison of cortisol concentrations in saliva and plasma of dogs. *Research in Veterinary Science* 53: 342-345

- Von der Ohe CG and Servheen C** 2002 Measuring stress in mammals using faecal glucocorticoids: opportunities and challenges. *Wildlife Society Bulletin* 30: 1215-1225
- Washburn BE and Millspaugh JJ** 2002 Effects of simulated environmental conditions on glucocorticoid metabolite measurements in white-tailed deer faeces. *General and Comparative Endocrinology* 127: 217-222
- Wasser SK, Hunt KE, Brown JL, Cooper K, Crockett CM, Bechert U, Millspaugh JJ, Larson S and Monfort SL** 2000 A generalized fecal glucocorticoid assay for use in a diverse array of non domestic mammalian and avian species. *General and Comparative Endocrinology* 120: 260-275
- Wells DL and Hepper PG** 2000 The influence of environmental change on the behaviour of sheltered dogs. *Applied Animal Behaviour Science* 68: 151-162
- Wells DL** 2004 A review of environmental enrichment for kennelled dogs, *Canis familiaris*. *Applied Animal Behaviour Science* 85: 307-317
- Wells DL** 2005 A note on the influence of visitors on the behaviour and welfare of zoo-housed gorillas. *Applied Animal Behaviour Science* 93: 13-17
- Wells DL, Hepper PG, Coleman D and Challis MG** 2007 A note on the effect of olfactory stimulation on the behaviour and welfare of zoo-housed gorillas. *Applied Animal Behaviour Science* 106: 155-160
- Whitten PL, Brockman DK and Stavisky RC** 1998 Recent advances in non-invasive techniques to monitor hormone-behaviour interactions. *Yearbook of Physical Anthropology* 41: 1-23
- Wielebnowski NC, Fletchall N, Carlstead K, Busso JM and Brown JL** 2002 Non-invasive assessment of adrenal activity associated with husbandry and behavioural factors in the North American clouded leopard population. *Zoo Biology* 21: 77-98
- Zayan R** 1991 The specificity of social stress. *Behavioural Process* 25: 81-93
- Zoldag L, Heuwiesser W, Grunert E and Stephan E** 1983 Steroid hormone profile in pregnant cows after exposure to noise stress, with particular reference to corticosteroids. *Zentral Veterinärmed* 30: 737-748