



On the genericity of cuspidal automorphic forms of $\mathrm{SO}(2n + 1)$, II

Dihua Jiang and David Soudry

ABSTRACT

This paper is a continuation of our previous work (D. Jiang and D. Soudry, *On the genericity of cuspidal automorphic forms on SO_{2n+1}* , J. reine angew. Math., to appear). We extend Moeglin’s results (C. Moeglin, J. Lie Theory **7** (1997), 201–229, 231–238) from the even orthogonal groups to odd orthogonal groups and complete our proof of the CAP conjecture for irreducible cuspidal automorphic representations of $\mathrm{SO}_{2n+1}(\mathbb{A})$ with special Bessel models. We also give a characterization of the vanishing of the central value of the standard L -function of $\mathrm{SO}_{2n+1}(\mathbb{A})$ in terms of theta correspondence. As a result, we obtain the weak Langlands functorial transfer from $\mathrm{SO}_{2n+1}(\mathbb{A})$ to $\mathrm{GL}_{2n}(\mathbb{A})$ for irreducible cuspidal automorphic representations of $\mathrm{SO}_{2n+1}(\mathbb{A})$ with special Bessel models.

1. Introduction

Let G be a reductive algebraic group defined over a number field k . Assume that G is k -quasi-split. It is known that irreducible cuspidal automorphic representations π of $G(\mathbb{A})$, where \mathbb{A} is the ring of adèles of k , may not be generic, i.e. may not have a nonzero Whittaker Fourier coefficient. It is interesting to know how close, in terms of near equivalence, the non-generic cuspidal automorphic representations are to irreducible generic cuspidal automorphic representations. Two irreducible automorphic representations π_1 and π_2 are nearly equivalent if, at almost all places ν of k , the local components $\pi_{1,\nu}$ and $\pi_{2,\nu}$ are equivalent, as representations of $G(k_\nu)$. In [JS], we state the CAP conjecture, which says that, for an irreducible cuspidal automorphic representation π of $G(\mathbb{A})$, there exists generic cuspidal data (P, σ) , where $P = MN$ is a parabolic k -subgroup of G and σ is an irreducible generic cuspidal automorphic representation of the Levi subgroup $M(\mathbb{A})$, such that π is nearly equivalent to an irreducible constituent of the induced representation $\mathrm{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})}(\sigma)$ (see [JS, Conjecture 1.1]). Further remarks on this conjecture and its relation to the Arthur conjecture can be found in [JS].

In [JS], we study the CAP conjecture for $G = \mathrm{SO}_{2n+1}$, the k -split odd special orthogonal group. More precisely, we study the CAP conjecture for irreducible cuspidal automorphic representations of $\mathrm{SO}_{2n+1}(\mathbb{A})$ with a nonzero special Bessel model. From its definition [JS, § 2] a special Bessel model is attached to the sub-regular nilpotent orbit of the corresponding Lie algebra, while a Whittaker model is attached to the regular nilpotent orbit. Hence, the family of irreducible cuspidal automorphic representations of $\mathrm{SO}_{2n+1}(\mathbb{A})$ that have special Bessel models is expected to be very close to irreducible generic cuspidal automorphic representations. Indeed, in Proposition 2.2 and

Received 14 August 2006, accepted in final form 24 November 2006.

2000 Mathematics Subject Classification 22E, 11F.

Keywords: cuspidal automorphic forms, Langlands functoriality.

The research of the first author is partially supported by NSF grant DMS 0400414 and by the distinguished visiting professorship at The Institute of Mathematics, CAS. The research of the second author is supported by a grant from the Israel Science Foundation.

This journal is © Foundation Compositio Mathematica 2007.

Theorems 4.1 and 4.6 of [JS], we show almost completely that the CAP conjecture holds for irreducible cuspidal automorphic representations of $\mathrm{SO}_{2n+1}(\mathbb{A})$ that have special Bessel models. What remained to be done was to check the genericity of the cuspidal data involved in the CAP conjecture for this case. In order to determine the genericity of the cuspidal data, we need to know certain basic results about the theta correspondence for the reductive dual pair $(\widetilde{\mathrm{Sp}}_{2m}(\mathbb{A}), \mathrm{SO}_{2n+1}(\mathbb{A}))$. Such results for the theta correspondence for the pair $(\mathrm{Sp}_{2m}(\mathbb{A}), \mathrm{SO}_{2n}(\mathbb{A}))$ were established by Mœglin in [Moe97a, Moe97b]. One of the main ingredients of the proofs of Mœglin’s theorems is the regularized Siegel–Weil formula [KR94] and its variants. Thanks to the work of Ichino [Ioh01] on the regularized Siegel–Weil formula for $\widetilde{\mathrm{Sp}}_{2m}(\mathbb{A})$, we are able to apply the arguments in Mœglin’s proofs to our current case.

In order to make the paper self-contained, as much as possible, we summarize in §§ 2.1 and 2.2 the details of basic facts about both the local and global Rao cocycles and metaplectic groups. In § 2.3, we prove a variant of the regularized Siegel–Weil formula for $\widetilde{\mathrm{Sp}}_{2n}(\mathbb{A})$ (Corollary 2.2) based on the work of Ichino [Ioh01] and the local theory of the Siegel–Weil formula [Ral84, KR90a, Zhu07], and then extend the theorem of Mœglin on the regularized Siegel–Weil formula for SO_{2n} to any number field, by using the regularization in terms of local p -adic Hecke elements (Theorem 2.4).

In § 3, we establish certain basic properties of the theta correspondence for the reductive dual pair $(\widetilde{\mathrm{Sp}}_{2m}, \mathrm{SO}_{2n+1})$, which are analogues of Mœglin’s theorems for the reductive dual pair $(\mathrm{Sp}_{2m}, \mathrm{SO}_{2n})$. Since our proofs are very similar to Mœglin’s in [Moe97a, Moe97b], we either indicate briefly, for the sake of completeness, Mœglin’s proof for our case or just refer directly to [Moe97a, Moe97b], for the parts which carry over, word for word. To state these theorems, we need further notation.

Let Z be a symplectic vector space of dimension $2n$ defined over k , and let V be a quadratic vector space of odd dimension m defined over k . For a positive integer b , denote by Z_b the direct sum of the space Z and b copies of two-dimensional k -symplectic vector spaces. Similarly, denote by V_b the direct sum of the space V and b copies of k -hyperbolic planes. Denote by $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$ (or $\widetilde{\mathrm{Sp}}(Z_b)_{\mathbb{A}}$) the corresponding adelic metaplectic group attached to Z (or Z_b , respectively) and, similarly, by $\mathrm{O}(V)_{\mathbb{A}}$ (or $\mathrm{O}(V_b)_{\mathbb{A}}$) the adelic orthogonal group attached to V (or V_b , respectively). Let ψ be a non-trivial character of $k \backslash \mathbb{A}$. We denote by $\theta_{\psi, Z}^V$ the ψ -theta correspondence from $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$ to $\mathrm{O}(V)_{\mathbb{A}}$, and by $\theta_{\psi, V}^Z$ the ψ -theta correspondence in the other direction.

THEOREM 1.1. *Let $\tilde{\pi}$ be an irreducible, genuine, cuspidal, automorphic representation of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$. Assume that $\theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ is cuspidal.*

(1) *The following identity holds*

$$\theta_{\psi, V}^Z(\theta_{\psi^{-1}, Z}^V(\tilde{\pi})) = \tilde{\pi}.$$

(2) *Let b be a positive integer. Then $\theta_{\psi, V}^{Z_b}(\theta_{\psi^{-1}, Z}^V(\tilde{\pi}))$ is orthogonal to all cusp forms on $\widetilde{\mathrm{Sp}}(Z_b)_{\mathbb{A}}$.*

(3) *Let b be a positive integer, and let Z' be a symplectic subspace of Z , such that $Z = Z'_b$. Put, for brevity, $Z' = Z_{-b}$. Then*

$$\theta_{\psi, V}^{Z_{-b}}(\theta_{\psi^{-1}, Z}^V(\tilde{\pi})) = 0.$$

THEOREM 1.2. *Let σ be an irreducible, cuspidal, automorphic representation of $\mathrm{O}(V)_{\mathbb{A}}$. Assume that $\theta_{\psi^{-1}, V}^Z(\sigma)$ is cuspidal.*

(1) *The following identity holds*

$$\theta_{\psi, Z}^V(\theta_{\psi^{-1}, V}^Z(\sigma)) = \sigma.$$

(2) *Let b be a positive integer. Then $\theta_{\psi, Z}^{V_b}(\theta_{\psi^{-1}, V}^Z(\sigma))$ is orthogonal to all cusp forms on $\mathrm{O}(V_b)_{\mathbb{A}}$.*

- (3) Let b be a positive integer, and let V_{-b} be a quadratic subspace of V , such that $V = (V_{-b})_b$. Then

$$\theta_{\psi, Z}^{V_{-b}}(\theta_{\psi^{-1}, V}^Z(\sigma)) = 0.$$

The following theorem is deduced from Theorems 1.1 and 1.2, in exactly the same way as in [Moe97a, § 3].

THEOREM 1.3. *Let $\tilde{\pi}$ (respectively, σ) be as in Theorem 1.1 (respectively, Theorem 1.2).*

- (1) *The representation $\theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ (respectively, $\theta_{\psi^{-1}, V}^Z(\sigma)$) is irreducible, and for all positive integers b , $\theta_{\psi^{-1}, Z}^{V_b}(\tilde{\pi})$ (respectively, $\theta_{\psi^{-1}, V}^{Z_b}(\sigma)$) is orthogonal to all cusp forms on $O(V_b)_{\mathbb{A}}$ (respectively, $\widetilde{Sp}(Z_b)_{\mathbb{A}}$).*
- (2) *Let $\tilde{\pi}'$ (respectively, σ') be an irreducible cuspidal automorphic representation of $\widetilde{Sp}(Z_b)_{\mathbb{A}}$ (respectively, $O(V_b)_{\mathbb{A}}$), where b is a non-negative integer; we assume that $\tilde{\pi}'$ is genuine. Then $\theta_{\psi^{-1}, Z_b}^V(\tilde{\pi}') = \theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ (respectively, $\theta_{\psi^{-1}, V_b}^Z(\sigma') = \theta_{\psi^{-1}, V}^Z(\sigma)$), if and only if $b = 0$ and $\tilde{\pi} = \tilde{\pi}'$ (respectively, $\sigma = \sigma'$).*
- (3) *Let $\tilde{\pi}, \tilde{\pi}'$ (respectively, σ, σ') be two isomorphic, irreducible cuspidal automorphic representations of $\widetilde{Sp}(Z)_{\mathbb{A}}$ (respectively, of $O(V)_{\mathbb{A}}$). We assume that $\tilde{\pi}, \tilde{\pi}'$ are genuine. Then*

$$\theta_{\psi^{-1}, Z}^V(\tilde{\pi}) \neq 0 \Leftrightarrow \theta_{\psi^{-1}, Z}^V(\tilde{\pi}') \neq 0,$$

and

$$\theta_{\psi^{-1}, V}^Z(\sigma) \neq 0 \Leftrightarrow \theta_{\psi^{-1}, V}^Z(\sigma') \neq 0.$$

In § 4, we discuss several applications of Theorems 1.1, 1.2, and 1.3. First we establish a criterion for the non-vanishing of the central value of the standard L -function, $L(\sigma, \frac{1}{2})$ attached to an irreducible generic cuspidal automorphic representation σ of $SO_{2n+1}(\mathbb{A})$ in terms of the ψ -theta lift of σ to the Witt tower $\widetilde{Sp}_{2m}(\mathbb{A})$ (Theorems 4.1 and 4.2). We prove the following.

THEOREM 1.4. *Let σ be an irreducible cuspidal automorphic representation of $SO_{2n+1}(\mathbb{A})$. Assume that σ is generic, i.e. has a nonzero Whittaker Fourier coefficient. Then the ψ -theta lift of σ to $\widetilde{Sp}_{2n}(\mathbb{A})$, $\tilde{\theta}_{\psi, n}^n(\sigma)_+$, is non-trivial if and only if $L(\sigma, \frac{1}{2}) \neq 0$, where $L(\sigma, s)$ is the standard L -function attached to σ .*

See § 4 for comments on the relation of this theorem to Furusawa’s work [Fur95] and to the work of Howe and Piatetski-Shapiro [HP83].

In the rest of § 4 we complete our proof of the CAP conjecture for irreducible cuspidal automorphic representations of $SO_{2n+1}(\mathbb{A})$ with special Bessel models. In fact, we obtain additional information about the generic cuspidal data in terms of the central value of standard L -functions. We state this now. Here, for $\lambda \in k^*$, which is not a square, α_{λ} denotes the global Hilbert symbol (\cdot, λ) , and χ_{λ} denotes the character of $SO_{2n+1}(\mathbb{A})$, which is the composition of the spinor norm with α_{λ} . For other unexplained notation, see [JS].

THEOREM 1.5. *Let σ be an irreducible cuspidal automorphic representation of $SO_{2n+1}(\mathbb{A})$. Assume that σ has a nonzero Bessel model of special type, i.e. of type $(D_{\lambda}, 1, \psi_{n, n-1; \lambda})$. Then either σ is nearly equivalent to an irreducible generic cuspidal automorphic representation, or σ is CAP with respect to generic cuspidal data. More precisely, the following hold.*

- (1) *If the special Bessel model is k -split, i.e. $\lambda \in k^{\times}$ is a square, then σ is generic.*
- (2) *Assume that the special Bessel model is not k -split. Then the first occurrence, $m_{0, \psi}(\sigma)$, in the ψ -theta lifting tower satisfies*

$$n - 1 \leq m_{0, \psi}(\sigma) \leq n.$$

- (a) Assume that $m_{0,\psi}(\sigma) = n$. Then either σ is nearly equivalent to an irreducible generic cuspidal automorphic representation σ' , such that $L(\sigma' \otimes \chi_\lambda, \frac{1}{2}) \neq 0$, or σ is CAP with respect to the generic cuspidal data of the form

$$(P_1; \alpha_\lambda | \cdot |^{\frac{1}{2}} \otimes \sigma_{n-1}),$$

where P_1 is the standard parabolic subgroup, whose Levi part is isomorphic to $GL_1 \times SO_{2n-1}$, and σ_{n-1} is irreducible generic cuspidal automorphic representation of $SO_{2n-1}(\mathbb{A})$, such that $L(\sigma_{n-1} \otimes \chi_\lambda, \frac{1}{2}) = 0$.

- (b) Assume that $m_{0,\psi}(\sigma) = n - 1$. Then σ is a CAP representation. It is CAP either with respect to the generic cuspidal data

$$(P_1; | \cdot |^{\frac{1}{2}} \otimes \sigma_{n-1}),$$

such that $L(\sigma_{n-1} \otimes \chi_\lambda, \frac{1}{2}) \neq 0$; or with respect to the generic cuspidal data

$$(P_{1,1}; | \cdot |^{\frac{1}{2}} \otimes \alpha_\lambda | \cdot |^{\frac{1}{2}} \otimes \sigma_{n-2}),$$

where $P_{1,1}$ is the standard parabolic subgroup, whose Levi part is isomorphic to $GL_1 \times GL_1 \times SO_{2n-3}$, and $L(\sigma_{n-2} \otimes \chi_\lambda, \frac{1}{2}) = 0$.

It is not hard now to figure out from Theorem 1.5 the possible global Arthur parameters for irreducible automorphic cuspidal representations of $SO_{2n+1}(\mathbb{A})$, which have a Bessel model of special type. We will show, in a future work, that the relevant global Arthur packets can be constructed by theta correspondences.

2. Notation and preliminaries

2.1 The metaplectic group over a local field

Let F be a local field of characteristic 0 and let W be a vector space of dimension $2f$ over F , equipped with a non-degenerate anti-symmetric form $\langle \cdot, \cdot \rangle$. Let $Sp(W)$ denote the corresponding symplectic group acting from the right on W . For $F \neq \mathbb{C}$, $Sp(W)$ has a unique (up to isomorphism) non-trivial twofold cover $\widetilde{Sp}(W)$ (the metaplectic group). In this paper, we use the realization of $\widetilde{Sp}(W)$ in terms of a Rao normalized cocycle c_W (see [Ran93]), which may be simply called a Rao cycle. Thus, the metaplectic group $\widetilde{Sp}(W)$ is realized as

$$\{(g, \epsilon) \mid g \in Sp(W), \epsilon = \pm 1\},$$

with the group law

$$(g_1, \epsilon_1)(g_2, \epsilon_2) = (g_1 g_2, \epsilon_1 \epsilon_2 c_W(g_1, g_2)).$$

Let us recall briefly some facts about the Rao cocycle c_W . See [Ran93] for the details. The definition of a Rao (normalized) cocycle depends on a choice of a symplectic basis

$$B = \{\epsilon_1, \dots, \epsilon_f, \epsilon_{-f}, \dots, \epsilon_{-1}\}$$

of W , i.e. $W^+ = \text{Span}_k\{\epsilon_1, \dots, \epsilon_f\}$ and $W^- = \text{Span}_k\{\epsilon_{-1}, \dots, \epsilon_{-f}\}$ are totally isotropic subspaces of W , and $\langle \epsilon_i, \epsilon_{-j} \rangle = \delta_{ij}$, for $1 \leq i, j \leq f$. In particular, W^+ and W^- are transversal Lagrangians of W . (So, we should denote the cocycle by c_B , but this will make future notation quite cumbersome. Later on, we even drop W from c_W .)

Let $P \subset Sp(W)$ be the Siegel parabolic subgroup, which preserves W^- . Let us now list some properties of the Rao cocycle [Ran93].

(a) There is a function $x : Sp(W) \rightarrow F^*/(F^*)^2$, satisfying (1)–(5) as follows.

- (1) We have $x(p) \equiv \det_{W^-}(p) \pmod{(F^*)^2}$, $p \in P$.
- (2) We have $x(p_1hp_2) = x(p_1)x(h)x(p_2)$, for $p_1, p_2 \in P$ and $h \in Sp(W)$.
- (3) For $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(W)$, according to the polarization $W = W^+ \oplus W^-$, with $\det C \neq 0$,

$$x \begin{pmatrix} A & B \\ C & D \end{pmatrix} \equiv \det C \pmod{(F^*)^2}.$$

(4) Let (\cdot, \cdot) denote the Hilbert symbol of F . Then, for $p_1, p_2 \in P$, $h_1, h_2 \in Sp(W)$,

$$\frac{c_W(p_1h_1, h_2p_2)}{c_W(h_1, h_2)} = \frac{(x(p_1), x(h_1))(x(p_2), x(h_2))}{(x(p_1), x(p_2))(x(p_1p_2), x(h_1h_2))}$$

and

$$\frac{c_W(h_1p_1^{-1}, p_1h_2)}{c_W(h_1, h_2)} = (x(p_1), -x(h_1)x(h_2)).$$

(5) For each subset $S \subset \{1, \dots, f\}$, consisting of j elements, let τ_S be the element of $Sp(W)$, defined by $\epsilon_i\tau_S = -\epsilon_{-i}$ and $\epsilon_{-i}\tau_S = \epsilon_i$, for $i \in S$, and, otherwise, τ_S fixes the remaining elements of the basis B . Then the double coset $P\tau_S P$ depends only on $j = |S|$. Let τ_j be a choice of τ_S , for $|S| = j$. Then $\{\tau_j\}_{j=0}^f$ is a set of representatives for $P \backslash Sp(W) / P$. We have

$$x(\tau_j) = 1.$$

Note that the Rao function $g \mapsto x(g)$ is unique by [Ran93, Lemma 5.1].

(b) Let W_1, W_2 be two symplectic spaces of dimensions $2n_1, 2n_2$, respectively. Denote the corresponding symplectic forms by \langle, \rangle_{W_1} and \langle, \rangle_{W_2} . Let $W = W_1 \oplus W_2$ be the direct sum of these two spaces, with the symplectic form $\langle, \rangle = \langle, \rangle_{W_1} \oplus \langle, \rangle_{W_2}$. Let, for $l = 1, 2$, $i_l : W_l \rightarrow W$ be the embedding of W_l in the l th coordinate, and let $j_l : Sp(W_l) \rightarrow Sp(W)$ be the corresponding embedding of symplectic groups. Let $j : Sp(W_1) \times Sp(W_2) \rightarrow Sp(W)$ be the corresponding direct sum embedding, i.e.

$$j(g_1, g_2) = j_1(g_1)j_2(g_2) = j_2(g_2)j_1(g_1), \quad \text{for } g_l \in Sp(W_l).$$

Choose symplectic bases

$$B_l = \{\epsilon_1^{(l)}, \dots, \epsilon_{n_l}^{(l)}, \epsilon_{-n_l}^{(l)}, \dots, \epsilon_{-1}^{(l)}\}$$

for W_l , $l = 1, 2$. Let $B = B_1 \cup B_2$, and order B as follows. Take, first, the image, by i_1 , of the first n_1 elements of B_1 , then the image, by i_2 , of all elements of B_2 , then the image, by i_1 , of the last n_1 elements of B_1 . This is a symplectic basis of W . Let c_l be the Rao normalized cocycle on $Sp(W_l)$, corresponding to B_l ($l = 1, 2$), and let c_W be the Rao normalized cocycle on $Sp(W)$, corresponding to B . Denote by x_l the corresponding x -functions on $Sp(W_l)$ ($l = 1, 2$). Then [Ran93, Corollary 5.6] states that, for $g_1, g'_1 \in Sp(W_1)$ and $g_2, g'_2 \in Sp(W_2)$,

$$\frac{c_W(j(g_1, g_2), j(g'_1, g'_2))}{c_1(g_1, g'_1)c_2(g_2, g'_2)} = (x_1(g_1), x_2(g_2))(x_1(g'_1), x_2(g'_2))(x_1(g_1g'_1), x_2(g_2g'_2)). \tag{2.1}$$

In particular, for $g, h \in Sp(W_l)$, with $l = 1, 2$

$$c_W(j_l(g), j_l(h)) = c_l(g, h). \tag{2.2}$$

Thus, the restriction of the Rao normalized cocycle c_W to $Sp(W_l)$ is the Rao normalized cocycle c_l . Hence, we can lift, for $l = 1, 2$, j_l to an embedding $\tilde{j}_l : \widetilde{Sp}(W_l) \rightarrow \widetilde{Sp}(W)$ by $\tilde{j}_l(g, \epsilon) = (j_l(g), \epsilon)$. We have

$$\tilde{j}_1(\tilde{g}_1)\tilde{j}_2(\tilde{g}_2) = \tilde{j}_2(\tilde{g}_2)\tilde{j}_1(\tilde{g}_1),$$

and this defines a homomorphism $\tilde{j} : \widetilde{\text{Sp}}(W_1) \times \widetilde{\text{Sp}}(W_2) \longrightarrow \widetilde{\text{Sp}}(W)$, defined by $\tilde{j}(\tilde{g}_1, \tilde{g}_2) = \tilde{j}_1(\tilde{g}_1)\tilde{j}_2(\tilde{g}_2)$. Its kernel is $\{(I_{W_1}, \epsilon), (I_{W_2}, \epsilon) \mid \epsilon = \pm 1\}$.

(c) Assume that our local field F is non-archimedean of odd residual characteristic. Denote by \mathcal{O} its ring of integers and by $W(\mathcal{O})$ the lattice over \mathcal{O} spanned by the symplectic basis B . Let $K \subset \text{Sp}(W)$ be the stabilizer of $W(\mathcal{O})$. This is a maximal compact subgroup (standard with respect to B). It is known that there is an embedding $\varepsilon : K \longrightarrow \widetilde{\text{Sp}}(W)$ of the form $\varepsilon(r) = (r, \epsilon(r))$. See [MVW87, p. 43] for a proof. Note that this means that, for $r_1, r_2 \in K$,

$$c_W(r_1, r_2) = \frac{\epsilon(r_1 r_2)}{\epsilon(r_1)\epsilon(r_2)}.$$

This embedding is unique, if the residual field has at least four elements. It is not hard to see that, for $r \in P(\mathcal{O})$,

$$\varepsilon(r) = (r, 1). \tag{2.3}$$

(d) We will be interested in dual pairs $\text{Sp}(Z) \times \text{O}(V)$, inside $\text{Sp}(W)$, where $\text{O}(V)$ is the orthogonal group corresponding to a non-degenerate symmetric bilinear form Q on a vector space V of odd dimension $m = 2l + 1$, over F , and $\text{Sp}(Z)$ is the symplectic group, corresponding to a $2n$ -dimensional vector space over F , equipped with a symplectic form $\langle \cdot, \cdot \rangle_Z$. We view $\text{O}(V)$ as acting from the left on V and $\text{Sp}(Z)$ as acting from the right on Z . Thus, $W = Z \otimes V$ is of dimension $2mn$ over F and is equipped with the symplectic form $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_Z \otimes Q$. Let

$$\alpha : \text{Sp}(Z) \times \text{O}(V) \longrightarrow \text{Sp}(W)$$

be the natural homomorphism; it is given by $(z \otimes v)\alpha(g, h) = zg \otimes h^{-1}v$. Let d_Q and h_Q be the discriminant and the Hasse invariant, respectively, of Q . Choose an orthogonal basis $\{e_1, \dots, e_m\}$ of V , and put $a_i = Q(e_i, e_i)$, for $1 \leq i \leq m$. Choose a symplectic basis $B_Z = \{z_1, \dots, z_n, z_{-n}, \dots, z_{-1}\}$, for Z . Denote by B_Z^+ the first n elements of B_Z and by B_Z^- the last n elements. We also denote by X and Y the transversal Lagrangian subspaces of Z spanned by B_Z^+ and B_Z^- , respectively. Let c denote the corresponding Rao normalized cocycle on $\text{Sp}(Z)$. Let

$$B = B_Z^+ \otimes e_1 \cup \dots \cup B_Z^+ \otimes e_m \cup a_m^{-1} B_Z^- \otimes e_m \cup \dots \cup a_1^{-1} B_Z^- \otimes e_1.$$

This is a symplectic basis for W . Let c_W denote the corresponding normalized Rao cocycle on $\text{Sp}(W)$. Then, from [Kud96, pp. 20 and 35] one can get the following equivalence of c and the restriction of c_W on $\alpha_1(\text{Sp}(Z)) \times \alpha_1(\text{Sp}(Z))$, where $\alpha_1(g) = \alpha(g, I_V)$. (From [Kud96] one sees clearly that the double cover of $\text{Sp}(W)$ does not split over $\text{Sp}(Z)$ exactly due to the fact that m is odd; α_1 is equivalent to c^m .) We have, for $g_1, g_2 \in \text{Sp}(Z)$,

$$c_W(\alpha_1(g_1), \alpha_1(g_2)) = \frac{e(g_1 g_2)}{e(g_1)e(g_2)} c(g_1, g_2), \tag{2.4}$$

where

$$e(g) = h_Q^{u(g)}(x(g), (-1)^{\frac{1}{2}m(m-1)} d_Q^{u(g)+1})(d_Q, (-1)^{\frac{1}{2}u(g)(u(g)-1)}). \tag{2.5}$$

Here $u(g)$ is determined by writing g in the form $p_1 \tau_S p_2$, with $|S| = u(g)$, as in property (a) part (5). Note, also, that

$$x(\alpha_1(g)) \equiv d_Q^{u(g)} x(g)^m \pmod{(F^*)^2}. \tag{2.6}$$

The restriction of c_W to $\alpha_2(\text{O}(V)) \times \alpha_2(\text{O}(V))$, where $\alpha_2(h) = \alpha(I_Z, h)$ is very simple (and the following relation is valid also if m were even). For $h_1, h_2 \in \text{O}(V)$,

$$c_W(\alpha_2(h_1), \alpha_2(h_2)) = (\det h_1, \det h_2)^n. \tag{2.7}$$

Let $\widetilde{\text{O}}(V)$ be the group of all pairs $\{(h, \epsilon) \mid h \in \text{O}(V), \epsilon = \pm 1\}$, with the group law

$$(h_1, \epsilon_1)(h_2, \epsilon_2) = (h_1 h_2, \epsilon_1 \epsilon_2 (\det h_1, \det h_2)).$$

Then $\tilde{\alpha} : \widetilde{\mathrm{Sp}}(Z) \times \widetilde{\mathrm{O}}(V) \longrightarrow \widetilde{\mathrm{Sp}}(W)$, given by

$$\tilde{\alpha}((g, \epsilon_1), (h, \epsilon_2)) = (\alpha(g, h), \epsilon_1 \epsilon_2^n e(g)(d_Q^{u(g)} x(g), \det h)^n) \tag{2.8}$$

is a homomorphism, which lifts α . This follows from (2.4), (2.5), (2.7), and the fact that

$$c_W(\alpha_1(g), \alpha_2(h)) = c_W(\alpha_2(h), \alpha_1(g)) = (d_Q^{u(g)} x(g), \det h)^n, \tag{2.9}$$

which follows from (2.1) and (2.6).

We denote $\tilde{\alpha}_1(g, \epsilon) = \tilde{\alpha}((g, \epsilon), (I_V, 1))$ and $\tilde{\alpha}_2(h, \epsilon) = \tilde{\alpha}((I_Z, 1), (h, \epsilon))$. These homomorphisms from $\widetilde{\mathrm{Sp}}(Z)$ and $\widetilde{\mathrm{O}}(V)$ into $\widetilde{\mathrm{Sp}}(W)$ lift α_1 and α_2 , respectively. For other dual pairs in the symplectic group, see [Kud94].

(e) We keep the notation of property (d). Let ψ be a non-trivial character of F . Let ω_ψ be the Weil representation of $\widetilde{\mathrm{Sp}}(W)$, corresponding to ψ . We realize it in the Schrödinger model $S(X \otimes V)$. We will identify $X \otimes V \cong V^n$, through $z_1 \otimes v_1 + \dots + z_n \otimes v_n \longmapsto (v_1, \dots, v_n)$. Then we have the following formulae (see [Kud96, p. 37]).

$$\begin{aligned} \omega_\psi \left(\tilde{\alpha}_1 \left(\begin{pmatrix} a & \\ & a^* \end{pmatrix}, \epsilon \right) \right) \phi(v_1, \dots, v_n) &= \epsilon \gamma(\det a, \psi^{\frac{1}{2}})^{-1} (\det a, (-1)^{\frac{1}{2}m(m-1)} d_Q) \\ &\quad \cdot |\det a|^{m/2} \phi((v_1, \dots, v_n)a); \tag{2.10} \\ \omega_\psi \left(\tilde{\alpha}_1 \left(\begin{pmatrix} I_n & x \\ & I_n \end{pmatrix}, \epsilon \right) \right) \phi(v_1, \dots, v_n) &= \epsilon \psi(\frac{1}{2} \mathrm{tr}(\mathrm{Gr}(v_1, \dots, v_n)xw_n)) \phi(v_1, \dots, v_n); \\ \omega_\psi(\tilde{\alpha}_2(h, \epsilon)) \phi(v_1, \dots, v_n) &= \epsilon^n \gamma(\det h^{-n}, \psi^{\frac{1}{2}})^{-1} \phi(h^{-1}v_1, \dots, h^{-1}v_n). \end{aligned}$$

Here, the elements of $\mathrm{Sp}(Z)$ are written as matrices with respect to the basis B_Z . The element w_n in (2.10) is the $n \times n$ matrix which has 1 in the second main diagonal and zeroes elsewhere. Gr denotes the Gram matrix. Finally, $\gamma(t, \psi^x)$ is the Weil factor, with respect to the character ψ^x ($x \neq 0$). As is traditional, we normalize $\omega_\psi \circ \tilde{\alpha}$ so that the action of $\mathrm{O}(V)$ becomes linear. Thus, we define the following representation, which we still denote by ω_ψ , of $\widetilde{\mathrm{Sp}}(Z) \times \mathrm{O}(V)$,

$$\omega_\psi(\tilde{g}, h) = \gamma(\det h^{-n}, \psi^{\frac{1}{2}}) \omega_\psi(\tilde{\alpha}(\tilde{g}, (h, 1))), \tag{2.11}$$

for $\tilde{g} \in \widetilde{\mathrm{Sp}}(Z)$ and $h \in \mathrm{O}(V)$. Note that the first two formulae in (2.10) remain the same, and the last formula in (2.10) becomes

$$\omega_\psi((I_Z, 1), h) \phi(v_1, \dots, v_n) = \phi(h^{-1}v_1, \dots, h^{-1}v_n). \tag{2.12}$$

Sometimes, when we want to emphasize the dual pair, we will re-denote ω_ψ in (2.11) by $\omega_{\psi, Z, Q}$ or $\omega_{\psi, Z \otimes V}$.

2.2 The metaplectic group over an adèle ring

Let k be a number field. Let W be a vector space, of dimension $2f$ over k , equipped with a symplectic form $\langle \cdot, \cdot \rangle$. Let

$$B = \{\epsilon_1, \dots, \epsilon_f; \epsilon_{-f}, \dots, \epsilon_{-1}\}$$

be a symplectic basis of W , over k , as before. For each place ν of k , let $W_\nu = k_\nu \otimes_k W$. We will re-denote $k_\nu \otimes \epsilon_j = k_\nu \epsilon_j$, for $|j| = 1, \dots, f$. Denote by $\langle \cdot, \cdot \rangle_\nu$ the corresponding symplectic form on W_ν . Here B serves as a symplectic basis of W_ν as well. Denote by c_{W_ν} the Rao normalized cocycle on $\mathrm{Sp}(W_\nu)$ corresponding to B . For each finite odd place ν , we have an embedding $\epsilon_\nu : K_\nu \longrightarrow \widetilde{\mathrm{Sp}}(W_\nu)$ as in property (c) of § 2.1.

(a) Let \mathbb{A} be the ring of adèles of k and let $\widehat{\mathrm{Sp}}(W)_{\mathbb{A}}$ be the restricted product $\prod'_{\nu} \widetilde{\mathrm{Sp}}(W_{\nu})$ with respect to $\{\varepsilon_{\nu}(K_{\nu}) \mid \nu \text{ is finite and odd}\}$. Let

$$C' = \left\{ \prod_{\nu} (I, \varepsilon_{\nu}) \in \widehat{\mathrm{Sp}}(W)_{\mathbb{A}} \mid \prod_{\nu} \varepsilon_{\nu} = 1 \right\}.$$

Then

$$\widetilde{\mathrm{Sp}}(W)_{\mathbb{A}} := C' \backslash \widehat{\mathrm{Sp}}(W)_{\mathbb{A}} \tag{2.13}$$

is a twofold central cover of $\mathrm{Sp}(W)_{\mathbb{A}}$. The projection

$$\mathrm{pr} : \widetilde{\mathrm{Sp}}(W)_{\mathbb{A}} \longrightarrow \mathrm{Sp}(W)_{\mathbb{A}}$$

satisfies

$$\mathrm{pr} \left(C' \prod_{\nu \in S} (g_{\nu}, \varepsilon_{\nu}) \prod_{\nu \notin S} \varepsilon_{\nu}(k_{\nu}) \right) = \prod_{\nu \in S} g_{\nu} \prod_{\nu \notin S} k_{\nu},$$

where S is a finite, large enough set of places. The kernel of the projection is

$$C_2 = \left\{ C' \prod_{\nu} (I, \varepsilon_{\nu}) \in \widetilde{\mathrm{Sp}}(W)_{\mathbb{A}} \right\} \cong \{\pm 1\} = \mu_2.$$

Note that the natural projection $\widehat{\mathrm{Sp}}(W)_{\mathbb{A}} \longrightarrow \widetilde{\mathrm{Sp}}(W)_{\mathbb{A}}$ is injective on each $\widetilde{\mathrm{Sp}}(W_{\nu})$ and gives a commutative diagram

$$\begin{array}{ccc} \widetilde{\mathrm{Sp}}(W_{\nu}) & \longrightarrow & \widetilde{\mathrm{Sp}}(W)_{\mathbb{A}} \\ \downarrow & & \downarrow \\ \mathrm{Sp}(W_{\nu}) & \longrightarrow & \mathrm{Sp}(W)_{\mathbb{A}} \end{array}$$

where the vertical arrows are the natural projections.

(b) Let $\gamma \in \mathrm{Sp}(W)$. Let S be a finite set of places, including those at infinity and the even ones, such that $\gamma \in K_{\nu}$, for all $\nu \notin S$. Then there is a finite set of places $S' \supset S$, such that

$$\varepsilon_{\nu}(\gamma) = (\gamma, 1), \tag{2.14}$$

for all $\nu \notin S'$. This follows from the decomposition $\gamma = p_1 \tau_j p_2$ as in property (a) part (5) of §2.1, where now p_1, p_2 are in P , the Siegel k -parabolic subgroup of $\mathrm{Sp}(W)$ which stabilizes the subspace $W_{\nu}^{-} = \mathrm{Span}_k\{\varepsilon_{-1}, \dots, \varepsilon_{-f}\}$. Thus, we can take S' , such that $p_1, p_2 \in P(k_{\nu}) \cap K_{\nu}$, so that $x(p_i) \in \mathcal{O}_{\nu}^*$, for all $\nu \notin S'$ ($i = 1, 2$). By parts (2), (4) and (5) of property (a) in §2.1, and by (2.3), we get that there is $a_j = \pm 1$, such that, for $\nu \notin S'$,

$$\varepsilon_{\nu}(\gamma) = \varepsilon_{\nu}(p_1) \varepsilon_{\nu}(\tau_j) \varepsilon_{\nu}(p_2) = (p_1, 1) (\tau_j, a_j) (p_2, 1) = (\gamma, a_j).$$

To show that $a_j = 1$, we write τ_j as a product of elements of the form τ_2 . By [Ran93, Corollary 5.5], we may assume that $\tau_j = \tau_2$ and, then, it is a simple calculation in a SL_2 to show that $a_j = 1$.

(c) As a corollary, we get that the map

$$\gamma \mapsto C' \prod_{\nu} (\gamma, 1), \tag{2.15}$$

from (the k -rational points) $\mathrm{Sp}(W)$ goes to $\widetilde{\mathrm{Sp}}(W)_{\mathbb{A}}$. It is an injective group homomorphism, since for all $\gamma_1, \gamma_2 \in \mathrm{Sp}(W)$, we have

$$\prod_{\nu} c_{W_{\nu}}(\gamma_1, \gamma_2) = 1,$$

as follows from Rao’s explicit formula in [Ran93, Theorem 5.3]. We view $\mathrm{Sp}(W)$ as a subgroup of $\widetilde{\mathrm{Sp}}(W)_{\mathbb{A}}$, via (2.15). We re-denote the right-hand side of (2.15) by $(\gamma, 1)$.

(d) Let P_ν be the Siegel parabolic subgroup of $\mathrm{Sp}(W_\nu)$ stabilizing the subspace W_ν^- as in part (6) of this section and let $P_\mathbb{A}$ be the restricted product of all P_ν , with respect to $\{P_\nu \cap K_\nu \mid \nu < \infty\}$. Denote by $\tilde{P}_\mathbb{A}$, the inverse image of $P_\mathbb{A}$ inside $\widetilde{\mathrm{Sp}}(W)_\mathbb{A}$. By (2.13), it is easy to see that $\tilde{P}_\mathbb{A} \cong \{(p, \epsilon) \in P_\mathbb{A} \times \{\pm 1\}\}$, with the group law

$$(p_1, \epsilon_1)(p_2, \epsilon_2) = \left(p_1 p_2, \epsilon_1 \epsilon_2 \prod_\nu (\det p_{1,\nu}, \det p_{2,\nu})_\nu \right), \tag{2.16}$$

where the determinants of $p_{i,\nu}$ are taken as linear operators on the vector space W_ν^- . The isomorphism is given by

$$C' \prod_{\nu \in S} (p_\nu, \epsilon_\nu) \prod_{\nu \notin S} \varepsilon_\nu(p_\nu) \longrightarrow \left(\prod_\nu p_\nu, \prod_\nu \epsilon_\nu \right). \tag{2.17}$$

Here S is a large enough finite set, and $p_\nu \in P_\nu \cap K_\nu$, for $\nu \notin S$, so that by (2.3), $\varepsilon_\nu(p_\nu) = (p_\nu, 1)$ for $\nu \notin S$. In this case, we also re-denote, for brevity, the elements of $\tilde{P}_\mathbb{A}$ via (2.16) and (2.17) by (p, ϵ) , where $p \in P_\mathbb{A}$ and $\epsilon = \pm 1$.

(e) Assume that W is the (orthogonal) direct sum of two symplectic subspaces W_1 and W_2 . Then, as in property (6) of § 2.1, we have the direct sum embedding $j : \mathrm{Sp}(W_1) \times \mathrm{Sp}(W_2) \longrightarrow \mathrm{Sp}(W)$, and its restriction to each component is given by $j_l : \mathrm{Sp}(W_l) \longrightarrow \mathrm{Sp}(W)$, $l = 1, 2$. Similarly, we have the corresponding local embeddings $j_\nu, j_{l,\nu}$, and global embeddings $j_\mathbb{A}, j_{l,\mathbb{A}}$. Let us fix symplectic k -bases B_1, B_2 and $B = B_1 \cup B_2$ of W_1, W_2 , and W , respectively, as in property (b) of § 2.1. Then for each place ν , we have obtained the embeddings $\tilde{j}_{l,\nu} : \widetilde{\mathrm{Sp}}(W_{l,\nu}) \longrightarrow \widetilde{\mathrm{Sp}}(W_\nu)$, $l = 1, 2$, and the homomorphism $\tilde{j}_\nu : \widetilde{\mathrm{Sp}}(W_{1,\nu}) \times \widetilde{\mathrm{Sp}}(W_{2,\nu}) \longrightarrow \widetilde{\mathrm{Sp}}(W_\nu)$, which lift $j_{l,\nu}, l = 1, 2$, and j_ν , respectively. Let us denote, for finite and odd places ν , the standard maximal compact subgroup of $\mathrm{Sp}(W_{l,\nu})$ by $K_{l,\nu}$ ($l = 1, 2$). Consider the embeddings, as in property (c) of § 2.1, $\varepsilon_{l,\nu} : K_{l,\nu} \longrightarrow \widetilde{\mathrm{Sp}}(W_{l,\nu})$. Then, by (2.2) (if the residue field contains at least four elements),

$$\tilde{j}_{l,\nu}(\varepsilon_{l,\nu}(r)) = \varepsilon_\nu(j_{l,\nu}(r)),$$

for $l = 1, 2$, and $r \in K_{l,\nu}$. Thus, the collection $\{\tilde{j}_{l,\nu}\}_\nu$ defines an embedding $\tilde{j}_{l,\mathbb{A}} : \widetilde{\mathrm{Sp}}(W_l)_\mathbb{A} \longrightarrow \widetilde{\mathrm{Sp}}(W)_\mathbb{A}$, which lifts $j_{l,\mathbb{A}}$, and a homomorphism $\tilde{j}_\mathbb{A} : \widetilde{\mathrm{Sp}}(W_1)_\mathbb{A} \times \widetilde{\mathrm{Sp}}(W_2)_\mathbb{A} \longrightarrow \widetilde{\mathrm{Sp}}(W)_\mathbb{A}$, which lifts $j_\mathbb{A}$. We have

$$\tilde{j}_\mathbb{A}(\tilde{g}_1, \tilde{g}_2) = \tilde{j}_{1,\mathbb{A}}(\tilde{g}_1) \tilde{j}_{2,\mathbb{A}}(\tilde{g}_2).$$

Note that (with our conventions) for $\gamma_l \in \mathrm{Sp}(W_l)$ ($l = 1, 2$),

$$\tilde{j}_\mathbb{A}((\gamma_1, 1), (\gamma_2, 1)) = (j(\gamma_1, \gamma_2), 1).$$

(f) Let $\alpha : \mathrm{Sp}(Z) \times \mathrm{O}(V) \longrightarrow \mathrm{Sp}(W)$ be a dual pair, as in property (d) of § 2.1, and we use the same notation, except that here we assume that the vector space Z (respectively V) and the corresponding non-degenerate anti-symmetric (respectively symmetric) form are defined over the number field k . Recall that $\dim_k V = m = 2l + 1$ is odd. Let ψ be a non-trivial character of $k \setminus \mathbb{A}$. For each local place ν of k , consider the local Weil representation ω_{ψ_ν} of $\widetilde{\mathrm{Sp}}(Z_\nu) \times \mathrm{O}(V_\nu)$, realized in the Schrödinger model $\mathcal{S}(X_\nu \otimes V_\nu) \cong \mathcal{S}(V_\nu^n)$. Then the restricted tensor product $\omega_\psi = \otimes' \omega_{\psi_\nu}$ defines the global Weil representation of $\widetilde{\mathrm{Sp}}(Z)_\mathbb{A} \times \mathrm{O}(V)_\mathbb{A}$ on the Schrödinger model $\mathcal{S}(V_\mathbb{A}^n)$. It pulls back to $\widetilde{\mathrm{Sp}}(Z)_\mathbb{A} \times \mathrm{O}(V)_\mathbb{A}$. (Of course, it is obtained from the Weil representation of $\widetilde{\mathrm{Sp}}(Z \otimes V)_\mathbb{A}$.) Let θ be the theta distribution on $\mathcal{S}(V_\mathbb{A}^n)$,

$$\theta(\phi) = \sum_{x \in V^n} \phi(x)$$

and denote, for $(\tilde{g}, h) \in \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}} \times \mathrm{O}(V)_{\mathbb{A}}$ and $\phi \in S(V_{\mathbb{A}}^n)$,

$$\theta_{\psi}^{\phi}(\tilde{g}, h) = \theta(\omega_{\psi}(\tilde{g}, h)\phi),$$

and sometimes, more precisely,

$$\theta_{\psi, Z \otimes V}^{\phi}(\tilde{g}, h) = \theta(\omega_{\psi, Z \otimes V}(\tilde{g}, h)\phi).$$

Due to the $\mathrm{Sp}(Z \otimes V)$ -invariance of θ , we know that θ_{ψ}^{ϕ} is $\mathrm{Sp}(Z) \times \mathrm{O}(V)$ -left invariant. The function $\theta_{\psi}^{\phi}(\tilde{g}, h)$ is of moderate growth in each variable [HP83]. Thus, for an irreducible automorphic representation σ of $\mathrm{O}(V)_{\mathbb{A}}$, if either the quadratic form Q on V is anisotropic or σ is cuspidal, then the integrals

$$\theta_{\psi}^{\phi}(\varphi_{\sigma})(\tilde{g}) = \int_{\mathrm{O}(V) \backslash \mathrm{O}(V)_{\mathbb{A}}} \theta_{\psi}^{\phi}(\tilde{g}, h)\varphi_{\sigma}(h) dh \tag{2.18}$$

converge absolutely, for each φ_{σ} , in the space of σ , and $\phi \in S(V_{\mathbb{A}}^n)$. These automorphic functions $\theta_{\psi}^{\phi}(\varphi_{\sigma})$ generate a subspace of automorphic forms on $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, which is denoted by $\tilde{\theta}_{\psi, V}^Z(\sigma)$. It is stable under right translations by $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$. We use the same notation, for the automorphic representation of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, thus obtained. This is the automorphic representation of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$ obtained by the ψ -theta correspondence from σ , which may of course be zero. Similarly, we consider the ψ -theta correspondence in the reverse direction, starting with an irreducible genuine cuspidal automorphic representation $\tilde{\pi}$ of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$. The subspace of automorphic functions on $\mathrm{O}(V)_{\mathbb{A}}$ thus obtained is denoted by $\theta_{\psi, Z}^V(\tilde{\pi})$. It is an automorphic representation of $\mathrm{O}(V)_{\mathbb{A}}$ in the space generated by the integrals

$$\theta_{\psi}^{\phi}(\varphi_{\tilde{\pi}})(h) = \int_{C_2 \mathrm{Sp}(Z) \backslash \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}} \theta_{\psi}^{\phi}(\tilde{g}, h)\varphi_{\tilde{\pi}}(\tilde{g}) d\tilde{g}, \tag{2.19}$$

as ϕ and $\varphi_{\tilde{\pi}}$ vary in $S(V_{\mathbb{A}}^n)$ and the space of $\tilde{\pi}$, respectively.

2.3 The Siegel–Weil formula

We keep the notation of §2.2. We establish some simple variants of the regularized Siegel–Weil formulae based on the works of Kudla and Rallis [KR94], Ichino [Ioh01], and Mœglin [Moe97a, Moe97b].

Let (V, Q) be the nondegenerate quadratic vector space over k with Witt index r . If $r = 0$, the ψ -theta lifting of the trivial representation of $\mathrm{O}(V)_{\mathbb{A}}$ to $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$ does not need regularization. In the following we may assume that $r > 0$. We first recall Ichino’s formulation [Ioh01] of the Siegel–Weil formula. Assume that $\dim_k V = m$ is odd, and that

$$m = \dim_k V < \frac{1}{2} \dim_k Z + 1 = n + 1. \tag{2.20}$$

Since V has Witt index $r > 0$, we may write $V = V_0 \oplus H_r$, where V_0 is k -anisotropic and H_r is the direct sum of r hyperbolic planes. Let (V', Q') be the nondegenerate quadratic space over k in the same Witt class of (V, Q) , and of dimension $m' = 2n + 2 - m$. In this case we may write

$$V' = V_0 \oplus H_{n+1-r-\dim_k V_0}.$$

From (2.20), we have

$$n + 1 - r - \dim_k V_0 > m - r - \dim_k V_0 = r > 0.$$

Ichino’s theorem now applies with our m' replacing m in [Ioh01, p. 203]. Indeed, let $r' = n + 1 - r - \dim_k V_0$ denote the Witt index of V' . Then we clearly have $n + 1 < m' \leq 2n + 2$, and $m' - r' \leq n + 1$, which are two of the conditions in Ichino’s theorem (the regularized Siegel–Weil formula).

We recall some details of the regularized Siegel–Weil formula from [Ioh01]. In the case under consideration, the theta integral

$$\int_{\mathrm{O}(V)\backslash\mathrm{O}(V)_{\mathbb{A}}} \theta_{\psi}^{\phi}(\tilde{g}, h) dh \tag{2.21}$$

diverges in general. In [KR94], the regularization is carried out using a certain differential operator from the universal enveloping algebra at real archimedean local places. In [Ioh01], the regularization is obtained by using a certain element of a local p -adic Hecke algebra. The advantage of the approach in [Ioh01], whose possibility was mentioned in [KR94], is that it applies to any number field.

Let us describe Ichino’s regularization. Fix a finite place ν_0 of k , where the number of elements of the residue field is congruent to 1 modulo 4. Assume also that ψ_{ν_0} is unramified, and the quadratic form is unimodular at ν_0 . Assume that $\phi \in \mathcal{S}(V_{\mathbb{A}}^n)$ is fixed by $\omega_{\psi_{\nu_0}}(\varepsilon_{\nu_0}(K_{\nu_0}^Z \times K'_{\nu_0}))$. Here, $K_{\nu_0}^Z$ is a (standard) maximal compact subgroup of $\mathrm{Sp}(Z_{\nu_0})$, and K'_{ν_0} is a hyper-special maximal compact subgroup of $\mathrm{O}(V_{\nu_0})$. Then there is an element α_{ν_0} , which depends only on the place ν_0 , in the spherical Hecke algebra (genuine functions) of $\widetilde{\mathrm{Sp}}(Z_{\nu_0})$ with respect to $K_{\nu_0}^Z$, such that $\theta_{\psi}^{\omega_{\psi_{\nu_0}}(\alpha_{\nu_0}, 1)\phi}(\tilde{g}, h)$ is rapidly decreasing in h , and hence the integral

$$\int_{\mathrm{O}(V)\backslash\mathrm{O}(V)_{\mathbb{A}}} \theta_{\psi}^{\omega_{\psi_{\nu_0}}(\alpha_{\nu_0}, 1)\phi}(\tilde{g}, h) dh \tag{2.22}$$

converges absolutely. There is a non-zero constant c_{ν_0} , such that if the integral (2.21) converges absolutely, then

$$\int_{\mathrm{O}(V)\backslash\mathrm{O}(V)_{\mathbb{A}}} \theta_{\psi}^{\phi}(\tilde{g}, h) dh = c_{\nu_0}^{-1} \int_{\mathrm{O}(V)\backslash\mathrm{O}(V)_{\mathbb{A}}} \theta_{\psi}^{\omega_{\psi_{\nu_0}}(\alpha_{\nu_0}, 1)\phi}(\tilde{g}, h) dh.$$

In other words, the regularization (2.23) is a natural extension of (2.21), in its domain of convergence. Moreover, there is an element $\alpha'_{\nu_0} = \theta(\alpha_{\nu_0})$ in the spherical Hecke algebra of $\mathrm{O}(V_{\nu_0})$, with respect to K'_{ν_0} , such that

$$\omega_{\psi_{\nu_0}}(\alpha_{\nu_0}, 1) = \omega_{\psi_{\nu_0}}(1, \alpha'_{\nu_0}), \tag{2.23}$$

as endomorphisms of $\mathcal{S}(V_{\mathbb{A}}^n)^{\omega_{\psi_{\nu_0}}(\varepsilon_{\nu_0}(K_{\nu_0}^Z \times K'_{\nu_0}))}$. Here θ is the homomorphism from the Hecke algebra of $\widetilde{\mathrm{Sp}}(Z_{\nu_0})$ to the Hecke algebra of $\mathrm{O}(V_{\nu_0})$, given in [Ioh01, Proposition 1.1]. For $\phi \in \mathcal{S}(V_{\mathbb{A}}^n)$ and $\tilde{g} \in \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, the regularized theta integral is denoted by

$$I_{\mathrm{reg}, V, \psi}(\tilde{g}, \phi) = c_{\nu_0}^{-1} \int_{\mathrm{O}(V)\backslash\mathrm{O}(V)_{\mathbb{A}}} \theta_{\psi}^{\omega_{\psi_{\nu_0}}(\alpha_{\nu_0}, 1)\phi}(\tilde{g}, h) dh. \tag{2.24}$$

It is independent of the choice of ν_0 , as above, or the choice of α_{ν_0} . Finally, for $\phi' \in \mathcal{S}((V'_{\mathbb{A}})^n)$, $s \in \mathbb{C}$, and $\tilde{g} \in \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, the Siegel section attached to ϕ' is denoted by

$$f_s^{\phi'}(\tilde{g}) = |a(\tilde{g})|^{s-(n+1-m)/2} \omega_{\psi, Z}^{V'}(\tilde{g}, I_{V'})\phi'(0). \tag{2.25}$$

Here $a(\tilde{g})$ is defined as follows: if $g \in \mathrm{Sp}(Z)_{\mathbb{A}}$ is the projection of \tilde{g} and $g = pk$ is its Iwasawa decomposition with $p \in P_{\mathbb{A}}$ and $k \in K_{\mathbb{A}}^Z$ (with self-evident notation), then $a(\tilde{g}) = a(g) = \prod_{\nu} \det_{Y_{\nu}} p_{\nu}$. By comparing with [Ioh01, p. 202] we note that

$$\frac{m' - n - 1}{2} = \frac{n + 1 - m}{2}.$$

By (2.10) and using part (d) of § 2.2, we have

$$f_s^{\phi'} \left(\left(\left(\begin{pmatrix} a & x \\ & a^* \end{pmatrix}, \epsilon \right) \tilde{g} \right) \right) = \epsilon \chi_{V', \psi}(\det a) |\det a|^{s+(n+1)/2} f_s^{\phi'}(\tilde{g}), \tag{2.26}$$

where

$$\chi_{V',\psi}(x) = \prod_{\nu} \chi_{V',\psi_{\nu}}(x_{\nu}) = \prod_{\nu} \gamma(x_{\nu}, \psi_{\nu}^{\frac{1}{2}})^{-1}(x_{\nu}, (-1)^{\frac{1}{2}m'(m'-1)}d_{Q'})_{\nu}. \tag{2.27}$$

Note that

$$\chi_{V,\psi} = \chi_{V',\psi}. \tag{2.28}$$

Indeed, $(-1)^{\frac{1}{2}m'(m'-1)}d_{Q'} = (-1)^{\frac{1}{2}m(m-1)}d_Q$. We view $\chi_{V,\psi}$ as a character of $\tilde{P}_{\mathbb{A}}$. From (2.27), we see that

$$f_s^{\phi'} \in \text{Ind}_{\tilde{P}_{\mathbb{A}}}^{\tilde{\text{Sp}}(Z)_{\mathbb{A}}}(\chi_{V,\psi}|\det|^s).$$

Consider the Eisenstein series, which is the analytic continuation of the following absolutely convergent series when $\text{Re}(s) > n + 1/2$,

$$E(\tilde{g}, f_s^{\phi'}) = \sum_{\gamma \in P \backslash \text{Sp}(Z)} f_s^{\phi'}((\gamma, 1)\tilde{g}).$$

It has at most a simple pole at $s = (n + 1 - m)/2$. Finally, Ichino defines a map $r_{V',V} : \mathcal{S}(V'_{\mathbb{A}})^n \rightarrow \mathcal{S}(V_{\mathbb{A}}^n)$, which is $\tilde{\text{Sp}}(Z)_{\mathbb{A}}$ -intertwining (this is ‘ $\pi_{Q'}^Q, \pi_K$ ’ in [Ioh01]). Now, we can state the following.

THEOREM 2.1 (Ichino [Ioh01]). *In the notation above, assume that $m < n + 1$ (and m is odd). Then there is a non-zero constant c_0 , such that, for all $\phi' \in \mathcal{S}((V'_{\mathbb{A}})^n)$ and $\tilde{g} \in \tilde{\text{Sp}}(Z)_{\mathbb{A}}$,*

$$\text{Res}_{s=(n+1-m)/2} E(\tilde{g}, f_s^{\phi'}) = c_0 I_{\text{reg},V,\psi}(\tilde{g}, r_{V',V}(\phi')). \tag{2.29}$$

We derive the following corollary, which will be used in the following sections.

COROLLARY 2.2. *Assume that $m < n + 1$ (m odd). Then, for all $\phi \in \mathcal{S}(V_{\mathbb{A}}^n)$, there is $\phi' \in \mathcal{S}((V'_{\mathbb{A}})^n)$, such that for all $\tilde{g} \in \tilde{\text{Sp}}(Z)_{\mathbb{A}}$,*

$$I_{\text{reg},V,\psi}(\tilde{g}, \phi) = \text{Res}_{s=(n+1-m)/2} E(\tilde{g}, f_s^{\phi'}). \tag{2.30}$$

Proof. By checking the definition of $r_{V',V}$ (which is ‘ $\pi_{Q'}^Q, \pi_K$ ’ in [Ioh01, p. 203]), it is easy to see that at almost all finite places ν , the local map $r_{V',V_{\nu}}$ takes the characteristic function of the standard lattice in $(V'_{\nu})^n$ to the characteristic function of the standard lattice of V_{ν}^n . Let ν be an arbitrary place of k , and $\phi = \otimes_{\nu'} \phi_{\nu'}$ be a factorizable function in $\mathcal{S}(V_{\mathbb{A}}^n)$, whose local components $\phi_{\nu'}$ are fixed at all places, except at the given local place ν , where we let it vary. Consider the linear functional

$$l_{\nu}(\phi_{\nu}) = I_{\text{reg},V,\psi}(\tilde{1}, \phi).$$

Here, $\tilde{1}$ denotes the identity element of $\tilde{\text{Sp}}(Z)_{\mathbb{A}}$. We recall from (2.24) in the case $\nu_0 = \nu$ that

$$l_{\nu}(\omega_{\psi_{\nu},Z_{\nu},Q_{\nu}}(\tilde{g}_{\nu}, h_{\nu})\phi_{\nu}) = I_{\text{reg},V,\psi}(\tilde{g}_{\nu}, \phi).$$

In particular, l_{ν} is $O(V_{\nu})$ -invariant, i.e. it factors through the space of co-invariants $S(V_{\nu}^n)_{O(V_{\nu})}$. This last space is isomorphic to $R_n(V_{\nu})$, which is the space of functions on $\tilde{\text{Sp}}(Z_{\nu})$ generated by $f^{\phi_{\nu}}(\tilde{g}) = \omega_{\psi_{\nu}}(\tilde{g}, 1)\phi_{\nu}(0)$, as ϕ_{ν} varies in $\mathcal{S}(V_{\nu}^n)$. Note that

$$R_n(V_{\nu}) \subset \text{Ind}_{\tilde{P}_{\nu}}^{\tilde{\text{Sp}}(Z_{\nu})}(\chi_{V_{\nu},\psi_{\nu}}|\det|^{(m-n-1)/2}).$$

The isomorphism

$$d'_{\nu} : \mathcal{S}(V_{\nu}^n)_{O(V_{\nu})} \rightarrow R_n(V_{\nu})$$

is induced from $d_{\nu}(\phi_{\nu}) = f^{\phi_{\nu}}$. See [Ral84] for ν finite, [KR90a] for ν real, and [Zhu07] for ν complex. Thus, if $f^{\phi_{\nu}} = 0$, then $l_{\nu}(\omega_{\psi_{\nu}}(\tilde{g}, 1)\phi_{\nu}) = 0$ for all $\tilde{g} \in \tilde{\text{Sp}}(Z_{\nu})$. Consider the map

$$d_{\nu} \circ r_{V',V_{\nu}} : \mathcal{S}(V_{\nu}^m) \rightarrow R_n(V_{\nu}).$$

This map is $\widetilde{\mathrm{Sp}}(Z_\nu)$ -intertwining and $\mathrm{O}(V_\nu)$ -invariant [Ioh01, Lemma 5.5]. By [Ioh01, Lemma 5.1] for ν finite, [KR90a, Corollary 2.7] for ν real, and [Zhu07] for ν complex, and by our assumption that $m < n + 1$, $R_n(V_\nu)$ is an irreducible $\widetilde{\mathrm{Sp}}(Z_\nu)$ -module. Hence the map $d_\nu \circ r_{V'_\nu, V_\nu}$ is surjective. It follows that for a given $\phi_\nu \in \mathcal{S}(V_\nu^n)$, there is $\phi'_\nu \in \mathcal{S}((V'_\nu)^n)$ such that $f^{\phi_\nu} = f^{r_{V'_\nu, V_\nu}(\phi'_\nu)}$, and hence

$$l_\nu(\omega_{\psi_\nu}(\tilde{g}_\nu, 1)\phi_\nu) = l_\nu(\omega_{\psi_\nu}(\tilde{g}_\nu, 1)r_{V'_\nu, V_\nu}(\phi'_\nu)),$$

for all $\tilde{g}_\nu \in \mathrm{Sp}(\tilde{Z}_\nu)$. Therefore, for a given factorizable $\phi = \bigotimes_\nu \phi_\nu \in \mathcal{S}(V_\mathbb{A}^n)$, there is a factorizable $\phi' = \bigotimes_\nu \phi'_\nu \in \mathcal{S}((V'_\mathbb{A})^n)$, such that

$$I_{\mathrm{reg}, V, \psi}(\tilde{g}, \phi) = I_{\mathrm{reg}, V, \psi}(\tilde{g}, r_{V', V}(\phi')),$$

for all $\tilde{g} \in \mathrm{Sp}(\tilde{Z})_\mathbb{A}$. Now, the identity (2.31) follows from (2.30). □

Next, we are going to establish a regularized Siegel–Weil formula characterizing residues of Siegel Eisenstein series on the k -split even orthogonal group $\mathrm{O}(l, l)$. This was done by Mœglin for totally real number fields by using the regularization at real archimedean local place [Moe97a, § 3.2]. We use the regularization in terms of elements in a p -adic Hecke algebra as in [Ioh01], which extends the results of Mœglin to all number fields. This result will be used in the next section. We give the details very briefly. Once we introduce the regularization, the rest follows exactly as in [Moe97a, § 3.2].

Let (U, b) be a non-degenerate quadratic space over k , which is k -split and even dimensional. Put $\dim_k U = 2l$. In this case, the double cover $\widetilde{\mathrm{Sp}}(U \otimes Z)_\mathbb{A}$ splits over $\mathrm{Sp}(Z)_\mathbb{A} \times \mathrm{O}(U)_\mathbb{A}$. Denote, again, by ω_ψ or $\omega_{\psi, Z \otimes U}$ the Weil representation of $\mathrm{Sp}(Z)_\mathbb{A} \times \mathrm{O}(U)_\mathbb{A}$ obtained by composing the ψ -Weil representation of $\widetilde{\mathrm{Sp}}(U \otimes Z)_\mathbb{A}$ with an embedding

$$\mathrm{Sp}(Z)_\mathbb{A} \times \mathrm{O}(U)_\mathbb{A} \rightarrow \widetilde{\mathrm{Sp}}(U \otimes Z)_\mathbb{A}.$$

Consider the corresponding theta series $\theta_\psi^\phi(g, h)$, where ϕ is in a corresponding Schrödinger model $\mathcal{S}(U_\mathbb{A}^n)$. The theta integral involved in the regularized Siegel–Weil formula characterizing residues of Siegel Eisenstein series on $\mathrm{O}(U)_\mathbb{A}$ is given by

$$\int_{\mathrm{Sp}(Z) \backslash \mathrm{Sp}(Z)_\mathbb{A}} \theta_\psi^\phi(g, h) dg,$$

which is divergent in general if $n < l$, so that regularization is necessary.

First, we fix a local finite place ν_0 , such that ψ_{ν_0} is unramified and the local quadratic form b_{ν_0} is k_{ν_0} -unimodular. Assume that ϕ is fixed by $\omega_{\psi_{\nu_0}}(K_{\nu_0}^Z \times K_{\nu_0}^U)$, where $K_{\nu_0}^Z$ (respectively, $K_{\nu_0}^U$) is a (standard) maximal compact subgroup of $\mathrm{Sp}(Z_{\nu_0})$ (respectively, $\mathrm{O}(U_{\nu_0})$). (Thus, the choice of ν_0 depends on ϕ .) The local Howe duality asserts that $\omega_{\psi_{\nu_0}}(\mathcal{H}_{\mathrm{O}(U_{\nu_0})})$ and $\omega_{\psi_{\nu_0}}(\mathcal{H}_{\mathrm{Sp}(Z_{\nu_0})})$ coincide as algebras of operators on $\mathcal{S}(U_{\nu_0}^n)^{\omega_{\psi_{\nu_0}}(K_{\nu_0}^Z \times K_{\nu_0}^U)}$, where $\mathcal{H}_{\mathrm{Sp}(Z_{\nu_0})}$ (respectively, $\mathcal{H}_{\mathrm{O}(U_{\nu_0})}$) is the spherical Hecke algebra of $\mathrm{Sp}(Z_{\nu_0})$ (respectively, $\mathrm{O}(U_{\nu_0})$) with respect to $K_{\nu_0}^Z$ (respectively, $K_{\nu_0}^U$); see [How79] and [MVW87]. Following the explicit calculation of the local theta correspondence (Howe duality) for unramified representations by Rallis in [Ral82], one has the following.

PROPOSITION 2.3. *Assume that $n \leq l$. Let $\theta = \theta_{l, n, \nu_0}$ be the Hecke algebra homomorphism*

$$\mathbb{C}[q^{\pm s_1}, \dots, q^{\pm s_l}]^{W_{\mathrm{O}(U)}} \cong \mathcal{H}_{\mathrm{O}(U_{\nu_0})} \xrightarrow{\theta} \mathcal{H}_{\mathrm{Sp}(Z_{\nu_0})} \cong \mathbb{C}[q^{\pm t_1}, \dots, q^{\pm t_n}]^{W_{\mathrm{Sp}(Z)}},$$

which is given by

$$\theta(q^{s_i}) = \begin{cases} q^{t_i}, & \text{if } 1 \leq i \leq n, \\ q^{-l+i}, & \text{if } n < i \leq l. \end{cases}$$

Then for all $\alpha \in \mathcal{H}_{\mathcal{O}(U_{\nu_0})}$,

$$\omega_{\psi_{\nu_0}, Z_{\nu_0}, b_{\nu_0}}(1, \alpha) = \omega_{\psi_{\nu_0}, Z_{\nu_0}, b_{\nu_0}}(\theta(\alpha), 1)$$

as endomorphisms of $\mathcal{S}(U_{\nu_0}^n)^{\omega_{\psi_{\nu_0}}(K_{\nu_0}^Z \times K_{\nu_0}^U)}$.

We have

$$\mathbb{C}[q^{\pm s_1}, \dots, q^{\pm s_l}]^{W_{\mathcal{O}(U)}} \cong \mathbb{C}[X_1, \dots, X_l]^{S_l},$$

where $X_i = q^{s_i} + q^{-s_i}$, $i = 1, \dots, l$; S_l is the symmetric group on l letters. Similarly, for $Y_i = q^{t_i} + q^{-t_i}$, $i = 1, \dots, n$,

$$\mathbb{C}[q^{\pm t_1}, \dots, q^{\pm t_n}]^{W_{\text{Sp}(Z)}} \cong \mathbb{C}[Y_1, \dots, Y_n]^{S_n}.$$

Let $\sigma_1, \dots, \sigma_l$ denote the elementary symmetric polynomials in X_1, \dots, X_l . We have, as in [KR94, Corollary 5.1.2] and [Ioh01, Lemma 1.3] that, for $n < l$, there is a unique element $\alpha_{l,n,\nu_0} \in \mathcal{H}_{\mathcal{O}(U_{\nu_0})}$, of the form

$$\alpha_{l,n,\nu_0} = \sigma_{n+1} - \sum_{i=1}^n a_i \sigma_i,$$

such that

$$\theta_{l,n,\nu_0}(\alpha_{l,n,\nu_0}) = 0.$$

Define, for $n \leq l$,

$$\alpha_{\nu_0} = \alpha_{l,n-1,\nu_0}.$$

This defines an element of $\mathcal{H}_{\mathcal{O}(U_{\nu_0})}$. We have $\theta_{l,n-1,\nu_0}(\alpha_{\nu_0}) = 0$, which means that

$$\alpha_{\nu_0}(Y_1, \dots, Y_{n-1}, q^{-l+n} + q^{l-n}, q^{-l+n+1} + q^{l-n-1}, \dots, q^{-1} + q, 2) = 0.$$

Note that α_{ν_0} is of degree n and is S_l -invariant. Consider the element $\theta(\alpha_{\nu_0}) = \theta_{l,n,\nu_0}(\alpha_{\nu_0})$. Then

$$\theta(\alpha_{\nu_0})(Y_1, \dots, Y_n) = \alpha_{\nu_0}(Y_1, \dots, Y_n, q^{-l+n+1} + q^{l-n-1}, \dots, q^{-1} + q, 2).$$

This is a polynomial of degree n in $\mathbb{C}[Y_1, \dots, Y_n]^{S_n} = \mathcal{H}_{\text{Sp}(Z_{\nu_0})}$. Since it is symmetric and satisfies $\theta(\alpha_{\nu_0})(Y_1, \dots, Y_{n-1}, q^{-l+n} + q^{l-n}) = 0$, we conclude, as in [KR94, Lemma 5.5.4] and as in [Ioh01, (1.1)] that

$$\theta(\alpha_{\nu_0})(Y_1, \dots, Y_n) = \prod_{i=1}^n (Y_i - (q^{-l+n} + q^{l-n})).$$

Let $\theta(\alpha_{\nu_0})$ act on the trivial representation of $\text{Sp}(Z_{\nu_0})$ by the scalar $c_{\alpha_{\nu_0}}$. Then

$$c_{\alpha_{\nu_0}} = \theta(\alpha_{\nu_0})(q^{-n} + q^n, q^{-n+1} + q^{n-1}, \dots, q^{-1} + q).$$

We conclude that, for $2n < l$, $c_{\alpha_{\nu_0}} \neq 0$.

Let us return to the global set-up. Recall that we choose the finite place α_{ν_0} , dependent on the function $\phi \in \mathcal{S}(U_{\mathbb{A}}^n)$. Assume that $n \leq l$, so that $\alpha_{\nu_0} = \alpha_{l,n-1,\nu_0}$ is defined. Then, as in [KR94, Proposition 5.3.1] and [Ioh01, Proposition 1.5], we have that $\theta_{\psi}^{\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi}(g, h)$ is rapidly decreasing in $g \in \text{Sp}(Z) \backslash \text{Sp}(Z_{\mathbb{A}})$, for all $h \in \mathcal{O}(U_{\mathbb{A}})$. For this, it is enough to take g as a Siegel domain and $h \in \prod_{\nu \neq \nu_0} \mathcal{O}(U_{\nu})$ (since $\mathcal{O}(U) \prod_{\nu \neq \nu_0} \mathcal{O}(U_{\nu})$ is dense in $\mathcal{O}(U_{\mathbb{A}})$). Then, it is enough to show that $\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi_{\nu_0}(y_1, \dots, y_n) = 0$, for linearly dependent $y_1, \dots, y_n \in U_{\nu_0}$. We assume, for simplicity, that ϕ is decomposable. As in [Ioh01, p. 210] we can find $a \in \text{GL}_n(\mathcal{O}_{\nu_0})$ and $x_1, \dots, x_{n-1} \in U_{\nu_0}$ such that $(y_1, \dots, y_n) = (x_1, \dots, x_{n-1}, 0)a$, and then, since ϕ_{ν_0} is fixed by $K_{\nu_0}^Z \times 1$ and the action of α_{ν_0} (via $\omega_{\psi_{\nu_0}}$) commutes with the action of (the Siegel parabolic subgroup of) $\text{Sp}(Z_{\nu_0})$, we get that

$$\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi_{\nu_0}(y_1, \dots, y_n) = \omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi_{\nu_0}(x_1, \dots, x_{n-1}, 0).$$

The last expression is zero, since $\theta_{l,n-1,\nu_0}(\alpha_{\nu_0}) = 0$. Indeed, let

$$\{z_1, \dots, z_n, z_{-n}, \dots, z_{-1}\},$$

be a symplectic basis of Z , as in property (d) of § 2.1. Put

$$Z' = \text{Span}_k\{z_1, \dots, z_{n-1}, z_{-n+1}, \dots, z_{-1}\}.$$

Denote by ω'_ψ the Weil representation for the dual pair $\text{Sp}(Z') \times \text{O}(U)$, and let us realize ω'_ψ in the Schrödinger model $\mathcal{S}(U_{\mathbb{A}}^{n-1})$. Denote the restriction of ϕ_{ν_0} to $U_{\nu_0}^{n-1}$ by ϕ'_{ν_0} . Then

$$\begin{aligned} \omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi_{\nu_0}(x_1, \dots, x_{n-1}, 0) &= \omega'_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi'_{\nu_0}(x_1, \dots, x_{n-1}) \\ &= \omega'_{\psi_{\nu_0}}(\theta_{l,n-1,\nu_0}(\alpha_{\nu_0}), 1)\phi'_{\nu_0}(x_1, \dots, x_{n-1}) \\ &= 0. \end{aligned}$$

This proves that

$$\theta_\psi^{\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi}(g, h) = \sum_{x \in U^n, \text{rank}(x)=n} \omega_\psi(g, h)\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0}\phi(x)),$$

where, for $x = (x_1, \dots, x_n)$, $\text{rank}(x)$ is the dimension of the subspace of U , spanned by x_1, \dots, x_n . Now, the rapid decrease in g follows as in [KR94, Proposition 5.3.1]. Moreover, if

$$\int_{\text{Sp}(Z)\backslash\text{Sp}(Z_{\mathbb{A}})} \theta_\psi^\phi(g, h) dg$$

converges absolutely, then

$$\int_{\text{Sp}(Z)\backslash\text{Sp}(Z_{\mathbb{A}})} \theta_\psi^{\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi}(g, h) dg = c_{\alpha_{\nu_0}} \int_{\text{Sp}(Z)\backslash\text{Sp}(Z_{\mathbb{A}})} \theta_\psi^\phi(g, h) dg.$$

As before, we define, for $2n < l$ (so that $c_{\alpha_{\nu_0}} \neq 0$)

$$I_{\text{reg},\psi}(h, \phi) = c_{\alpha_{\nu_0}}^{-1} \int_{\text{Sp}(Z)\backslash\text{Sp}(Z_{\mathbb{A}})} \theta_\psi^{\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi}(g, h) dg.$$

This definition is independent of the choice of the place ν_0 .

Next, consider a Siegel parabolic subgroup $P \subset \text{O}(U)$, whose Levi part is isomorphic to GL_l . Take a standard section ξ_s in the space of the normalized induced module $\text{Ind}_P^{\text{O}(U)_{\mathbb{A}}}(|\det|^s)$ and form an Eisenstein series $E(h, \xi_s)$ as usual. By [KR90b, Theorem 1.0.1], the Eisenstein series $E(h, \xi_s)$ converges absolutely when the real part of s is greater than $(l - 1)/2$; it has a meromorphic continuation to the complex plane \mathbb{C} and, after normalization, it has at most simple poles occurring only at $s = (l - 1)/2 - j \neq 0$ with $j \in \{0, 1, \dots, l - 1\}$.

Assume now that $n < (l - 1)/2$. Then $s = (l - 1)/2$ is a simple pole of the normalized Eisenstein series, and it is also a pole of $E(h, \xi_s)$. Now, we can repeat the proof of Mœglin, in [Moe97a, § 3.2], to get the following version of the regularized Siegel–Weil formula, which is valid for arbitrary number fields.

THEOREM 2.4. *Assume that $n < (l - 1)/2$. Then, for every Schwartz function ϕ as above, there is a section ξ_s as above, such that*

$$\int_{\text{Sp}(Z)\backslash\text{Sp}(Z_{\mathbb{A}})} \theta_\psi^{\omega_{\psi_{\nu_0}}(1, \alpha_{\nu_0})\phi}(g, h) dg = \text{Res}_{s=(l-1)/2-n} E(h, \xi_s). \tag{2.31}$$

3. The theta correspondence

We are going to prove Theorems 1.1, 1.2, and 1.3 as stated in the introduction. The main task is to give an explicit description of various spaces of automorphic functions via the theta correspondences. In § 3.1, we start with an irreducible genuine cuspidal automorphic representation $\tilde{\pi}$ of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, and in § 3.3, we start with an irreducible cuspidal automorphic representation σ of $\mathrm{O}(V)_{\mathbb{A}}$. Theorem 1.1 is proven in § 3.2 and the proofs of Theorems 1.2 and 1.3 are discussed in § 3.3.

3.1 Certain subspace of automorphic forms on $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$

Let $\tilde{\pi}$ be an irreducible genuine cuspidal automorphic representation of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$. The ψ^{-1} -theta lift of $\tilde{\pi}$ to $\mathrm{O}(V)_{\mathbb{A}}$ is denoted by $\theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ as before. Assume that $\theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ is a cuspidal representation of $\mathrm{O}(V)_{\mathbb{A}}$. For any integer $a \geq 0$, define a symplectic vector space Z_a by

$$Z_a = Z \oplus l_a, \tag{3.1}$$

where l_a denotes the $2a$ -dimensional symplectic space over k . We consider the ψ -theta lift $\theta_{\psi, V}^{Z_a}(\theta_{\psi^{-1}, Z}^V(\tilde{\pi}))$, which is a subspace of automorphic functions on the metaplectic group $\widetilde{\mathrm{Sp}}(Z_a)_{\mathbb{A}}$.

To fix a Rao normalized cocycle for $\widetilde{\mathrm{Sp}}(Z_a)_{\mathbb{A}}$ (at each place), we fix a symplectic k -basis

$$B'_a = \{z'_1, \dots, z'_a, z'_{-a}, \dots, z'_{-1}\}$$

of l_a , and a symplectic k -basis

$$B_Z = \{z_1, \dots, z_n, z_{-n}, \dots, z_{-1}\}$$

of Z . Then we take the symplectic k -basis

$$B_a = \{z'_1, \dots, z'_a, z_1, \dots, z_n, z_{-n}, \dots, z_{-1}, z'_{-a}, \dots, z'_{-1}\}$$

for Z_a . Put

$$l_a^{\pm} = \mathrm{Span}_F\{z'_{\pm 1}, \dots, z'_{\pm a}\}.$$

Denote by P_a the standard Siegel parabolic subgroup of $\mathrm{Sp}(Z_a \oplus Z)$, with respect to the basis $i_1(B_a) \cup i_2(B_Z)$, ordered as in property (b) of § 2.1, where i_1 (respectively, i_2) is the embedding of Z_a (respectively, Z) in the first (respectively, second) coordinate of $Z_a \oplus Z$. Consider, also, the homomorphism as in part (e) of § 2.2 (which we now shorten to \tilde{j} , instead of $\tilde{j}_{\mathbb{A}}$),

$$\tilde{j} : \widetilde{\mathrm{Sp}}(Z_a)_{\mathbb{A}} \times \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}} \rightarrow \widetilde{\mathrm{Sp}}(Z_a \times Z)_{\mathbb{A}}.$$

Let c be the k -linear automorphism of Z , such that $z_i c = z_i$ and $z_{-i} c = -z_{-i}$ for all $1 \leq i \leq n$. Clearly, $c \in \mathrm{GSp}(Z)$, and has similitude factor -1 . Let c act by conjugation on $\mathrm{Sp}(Z_{\nu})$, i.e.

$$g \mapsto g^c = c g c^{-1}.$$

Then one can lift this conjugation to an automorphism of $\widetilde{\mathrm{Sp}}(Z_{\nu})$ by

$$(g, \epsilon) = \tilde{g} \mapsto \tilde{g}' = (g, \epsilon)' = (g^c, \epsilon(x(g), (-1)^{u(g)+1})(-1, -1)^{\frac{1}{2}u(g)(u(g)-1)}),$$

in the notation of (2.5). See [Szp06] for details. This gives an automorphism of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, $\tilde{g} \mapsto \tilde{g}'$, which lifts the conjugation c on $\mathrm{Sp}(Z)_{\mathbb{A}}$. Finally, we denote by K_Z the standard (with respect to B_Z) maximal compact subgroup of $\mathrm{Sp}(Z)_{\mathbb{A}}$, and we let \tilde{K}_Z denote its inverse image in $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$. We use similar notation for other symplectic spaces.

PROPOSITION 3.1. *Let $\tilde{\pi}$ be an irreducible genuine cuspidal automorphic representation of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$ with $\dim_k Z = 2n$. Assume that $\theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ is cuspidal on $\mathrm{O}(V)_{\mathbb{A}}$ with $\dim_k V = m$. Let $a \geq 1$ be an integer such that $m < n + a + 1$. Then the subspace $\theta_{\psi, V}^{Z_a}(\theta_{\psi^{-1}, Z}^V(\tilde{\pi}))_0$ consisting of all \tilde{K}_{Z_a} -finite*

functions of the space $\theta_{\psi,V}^{Z_a}(\theta_{\psi^{-1},Z}^V(\tilde{\pi}))$ is contained in the subspace of automorphic forms on $\widetilde{Sp}(Z_a)_{\mathbb{A}}$ generated by the automorphic functions

$$\tilde{g}_a \mapsto \int_{C_2Sp(Z)\backslash\widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g}) \text{Res}_{s=(n+a+1-m)/2} E(\tilde{j}(\tilde{g}_a, \tilde{g}'), f_s) d\tilde{g},$$

where $E(\tilde{x}, f_s)$ is the Eisenstein series corresponding to a $\tilde{K}_{Z_a \oplus Z}$ -finite section f_s in $I_{2n+a}(\chi_{V,\psi}, s) = \text{Ind}_{\tilde{P}_a}^{\widetilde{Sp}(Z_a \oplus Z)_{\mathbb{A}}}(\chi_{V,\psi} |\det|^s)$.

Proof. We repeat [Moe97a, § 2.1]. By definition, the space $\theta_{\psi,V}^{Z_a}(\theta_{\psi^{-1},Z}^V(\tilde{\pi}))$ is spanned by

$$\int_{O(V)\backslash O(V)_{\mathbb{A}}} \theta_{\psi}^{\phi'}(\tilde{g}_a, h) \int_{C_2Sp(Z)\backslash\widetilde{Sp}(Z)_{\mathbb{A}}} \theta_{\psi^{-1}}^{\phi}(\tilde{g}, h) \varphi_{\tilde{\pi}}(\tilde{g}) d\tilde{g} dh, \tag{3.2}$$

where $\varphi_{\tilde{\pi}}$ is a cusp form in the space of $\tilde{\pi}$, $\phi \in \mathcal{S}(V_{\mathbb{A}}^n)$, $\phi' \in \mathcal{S}(V_{\mathbb{A}}^{n+a})$ and $\tilde{g}_a \in \widetilde{Sp}(Z_a)_{\mathbb{A}}$. We have (see [MVW87, p. 36]) that for $\tilde{g} \in \widetilde{Sp}(Z)_{\mathbb{A}}$,

$$\theta_{\psi^{-1}}^{\phi}(\tilde{g}, h) = \theta_{\psi}^{\phi}(\tilde{g}', h).$$

Viewing $\phi' \otimes \phi$ as an element of $\mathcal{S}(V_{\mathbb{A}}^{2n+a})$, we have (see [MVW87, p. 37]) that

$$\theta_{\psi}^{\phi'}(\tilde{g}_a, h) \theta_{\psi}^{\phi}(\tilde{g}', h) = \theta^{\phi' \otimes \phi}(\tilde{j}(\tilde{g}_a, \tilde{g}'), h).$$

Hence, the integral (3.2) becomes

$$\int_{O(V)\backslash O(V)_{\mathbb{A}}} \int_{C_2Sp(Z)\backslash\widetilde{Sp}(Z)_{\mathbb{A}}} \theta_{\psi}^{\phi' \otimes \phi}(\tilde{j}(\tilde{g}_a, \tilde{g}'), h) \varphi_{\tilde{\pi}}(\tilde{g}) d\tilde{g} dh. \tag{3.3}$$

In order to interchange the order of the integrations with respect to \tilde{g} and h , we have to regularize the integration with respect to h . To this end, we choose a finite place ν_0 , which satisfies all of the requirements as given in § 2.3. Take an element α_{ν_0} in the spherical Hecke algebra of $\widetilde{Sp}((Z_a \oplus Z)_{\nu_0})$ as in § 2.3. Then $\theta_{\psi}^{\omega_{\psi\nu_0}(\alpha_{\nu_0}, 1)(\phi' \otimes \phi)}(\tilde{j}(\tilde{g}_a, \tilde{g}'), h)$ is rapidly decreasing in h , where $\omega_{\psi\nu_0}$ denotes, for brevity, the Weil representation of

$$\widetilde{Sp}((Z_a \oplus Z)_{\nu_0}) \times O(V)_{\nu_0}.$$

Hence, we can interchange the order of integrations in (3.3) and obtain, by using (2.25), that (3.3) equals

$$c_{\nu_0}^{-1} \int_{C_2Sp(Z)\backslash\widetilde{Sp}(Z)_{\mathbb{A}}} \int_{O(V)\backslash O(V)_{\mathbb{A}}} \theta_{\psi}^{\omega_{\psi\nu_0}(\alpha_{\nu_0}, 1)(\phi' \otimes \phi)}(\tilde{j}(\tilde{g}_a, \tilde{g}'), h) dh \varphi_{\tilde{\pi}}(\tilde{g}) d\tilde{g},$$

which can be written as

$$\int_{C_2Sp(Z)\backslash\widetilde{Sp}(Z)_{\mathbb{A}}} I_{\text{reg}, V, \psi}(\tilde{j}(\tilde{g}_a, \tilde{g}'), \phi' \otimes \phi) \varphi_{\tilde{\pi}}(\tilde{g}) d\tilde{g}. \tag{3.4}$$

Finally the proposition follows from Corollary 2.2. □

In order to further describe the structure of the subspace $\theta_{\psi,V}^{Z_a}(\theta_{\psi^{-1},Z}^V(\tilde{\pi}))$ as in Proposition 3.1, we have to investigate the following integral

$$\int_{C_2Sp(Z)\backslash\widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g}) E(\tilde{j}(\tilde{g}_a, \tilde{g}'), f_s) d\tilde{g}. \tag{3.5}$$

Consider the polarization

$$Z_a \oplus Z = Z_{a,\Delta}^+ + Z_{a,\Delta}^-,$$

where $Z_{a,\Delta}^\pm = i_1(l_a^\pm) + \{(v, \pm vc) \mid v \in Z\}$. Note that for $g \in \text{Sp}(Z)$, $j(g, g^c)$ acts as identity on $i_1(l_a^\pm)$, and $(v, \pm vc)j(g, g^c) = (vg, \pm(vg)c)$. Here, we use notation as in properties (b) and (e) of §2.1. Let $P_{a,\Delta}$ be the Siegel parabolic subgroup of $\text{Sp}(Z_a \oplus Z)$, which preserves the maximal totally isotropic subspace $Z_{a,\Delta}^-$. Write its Levi decomposition $P_{a,\Delta} = M_{a,\Delta}U_{a,\Delta}$, where $M_{a,\Delta}$ is the Levi part, which is isomorphic to $\text{GL}(2n + a)$. Thus, $\beta(g) = j(g, g^c)$ defines an embedding of $\text{Sp}(Z)$ into $M_{a,\Delta}$. Let $\delta_0 \in \text{Sp}(Z_a \oplus Z)$ be such that it acts as the identity on $i_1(l_a)$ and $\delta_0 P_{a,\Delta} \delta_0^{-1} = P_a$, the standard Siegel parabolic subgroup of $\text{Sp}(Z_a \oplus Z)$ defined before Proposition 3.1. For a standard section f_s in $I_{2n+a}(\chi_{V,\psi}, s)$, we define $f'_s(\tilde{b}) = f_s((\delta_0, 1)\tilde{b})$. Then

$$f'_s \in I'_{2n+a}(\chi'_{V,\psi}, s) = \text{Ind}_{\widetilde{P}_{a,\Delta}(\mathbb{A})}^{\widetilde{\text{Sp}}(Z_a \oplus Z)_{\mathbb{A}}}(\chi'_{V,\psi} \cdot |\det_{Z_{a,\Delta}^-}|^s),$$

where $\chi'_{V,\psi}(\tilde{p}) = \chi_{V,\psi}((\delta_0, 1)\tilde{p}(\delta_0, 1)^{-1})$. We have

$$E(\tilde{x}, f_s) = E((\delta_0, 1)\tilde{x}, f_s) = E'(\tilde{x}, f'_s), \tag{3.6}$$

where E' denotes the Eisenstein series corresponding to $I'_{2n+a}(\chi'_{V,\psi}, s)$. It is easy to check that at each place ν , we have, for $p_\nu \in P_{a,\Delta}(k_\nu)$,

$$(\delta_0, 1)(p_\nu, \epsilon)(\delta_0, 1)^{-1} = (\delta_0 p_\nu \delta_0^{-1}, \epsilon_\nu c_\nu(\delta_0, p_\nu)(\det_{Z_{a,\Delta}^-}(p_\nu), x_\nu(\delta_0))_\nu), \tag{3.7}$$

where c_ν is the Rao normalized cocycle on $\widetilde{\text{Sp}}((Z_a \times Z)_\nu)$ with respect to the basis above. Note the following properties of f'_s .

LEMMA 3.2. *Let $R_a \subset \text{Sp}(Z_a)$ be the parabolic subgroup which preserves l_a^- , and its Levi decomposition $R_a = M_{R_a}U_{R_a}$. We identify M_{R_a} with $\text{GL}(l_a^+) \times \text{Sp}(Z)$. In the following, $\epsilon = \pm 1$ and $\tilde{b} \in \widetilde{\text{Sp}}(Z_a)_{\mathbb{A}}$.*

(1) *Let $u \in U_{R_a}(\mathbb{A})$. Then*

$$f'_s(\tilde{j}_1(u, \epsilon)\tilde{b}) = \epsilon f'_s(\tilde{b}).$$

(2) *Let $d \in M_{R_a}(\mathbb{A})$ correspond to an element of $\text{GL}(l_a^+)_{\mathbb{A}}$. Then*

$$f'_s(\tilde{j}_1(d, \epsilon)\tilde{b}) = \epsilon \cdot \chi_{V,\psi}(\det_{l_a^+} d) |\det_{l_a^+} d|^{s+n+(a+1)/2} f'_s(\tilde{b}).$$

(3) *Let $g \in M_{R_a}(\mathbb{A})$ correspond to an element in $\text{Sp}(Z)_{\mathbb{A}}$ and let \tilde{g} be an inverse image of g in $\widetilde{\text{Sp}}(Z)_{\mathbb{A}}$. Then*

$$f'_s(\tilde{j}_1(\tilde{g})\tilde{b}) = f'_s(\tilde{j}_2(\tilde{g}')^{-1}\tilde{b}).$$

Proof. By (3.7) and the fact that $j_1(U_{R_a}) \subset U_{a,\Delta}$, we have

$$f'_s(\tilde{j}_1(u, \epsilon)\tilde{b}) = \epsilon \cdot c(\delta_0, j_1(u)) f'_s(\tilde{b}).$$

It is clear that $u \mapsto \tilde{j}_1(u, 1) = (j_1(u), 1)$ is an embedding of $U_{R_a}(\mathbb{A})$ inside $\widetilde{\text{Sp}}(Z_a \times Z)_{\mathbb{A}}$. This implies that $u \mapsto c(\delta_0, j_1(u))$ is a μ_2 -valued character of $U_{R_a}(\mathbb{A})$, and hence is trivial. This proves part (1).

For part (2), since δ_0 is the identity on $j_1(l_a)$, it follows that δ_0 and $j_1(d)$ commute. Hence $(\delta_0, 1)$ and $\tilde{j}_1(d, \epsilon)$ commute. It follows that

$$\chi'_{V,\psi}(\tilde{j}_1(d, \epsilon)) = \chi_{V,\psi}(\tilde{j}_1(d, \epsilon)) = \epsilon \cdot \chi_{V,\psi}(\det_{l_a^+} d).$$

Now the assertion follows.

Finally, we have $f'_s(\tilde{j}_1(\tilde{g})\tilde{b}) = f'_s(\tilde{j}(\tilde{g}, \tilde{g}')\tilde{j}_2(\tilde{g}')^{-1}\tilde{b})$. Note that $j(g, g^c)$ lies in $M_{R_a}(\mathbb{A})$ and its determinant on either of $Z_{a,\Delta}^\pm(\mathbb{A})$ is $\det g = 1$. It follows that $f'_s(\tilde{j}_1(\tilde{g})\tilde{b}) = \chi'_{V,\psi}(\tilde{j}(\tilde{g}, \tilde{g}')) f'_s(\tilde{j}_2(\tilde{g}')^{-1}\tilde{b})$.

Clearly, $\tilde{g} \mapsto \chi'_{V,\psi}(\tilde{j}(\tilde{g}, \tilde{g}'))$ is a μ_2 -valued character of $\widetilde{\text{Sp}}(Z)_{\mathbb{A}}$, trivial on $C_2(\mathbb{A})$, and hence, it is trivial. This proves part (3). □

Motivated by the global integral (3.5), we define, as in [Moe97a, p. 211],

$$f_{\varphi_{\tilde{\pi},s}}(\tilde{g}_a) = \int_{C_2 \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g}) f'_s(j(\tilde{g}_a, \tilde{g}')) d\tilde{g}. \tag{3.8}$$

As in [Moe97a, § 2.1], the previous lemma implies that, if the integrals (3.8) converge absolutely at a point s , then $f_{\varphi_{\tilde{\pi},s}}$ is a \tilde{K}_{Z_a} -finite element of the representation

$$J(\tilde{\pi}, s) = \text{Ind}_{\tilde{R}_a(\mathbb{A})}^{\widetilde{Sp}(Z_a)_{\mathbb{A}}} (\mu_{\psi} |\det_{l_a^+}|^s \otimes \tilde{\pi}),$$

where $\mu_{\psi} |\det_{l_a^+}|^s \otimes \tilde{\pi}$ is the representation of $\tilde{R}_a(\mathbb{A})$, which is trivial on $(U_{R_a}(\mathbb{A}), 1)$ and acts as

$$\epsilon \cdot \chi_{V,\psi}(\det_{l_a^+} d) |\det_{l_a^+} d|^s \tilde{\pi}(\tilde{g}) \tag{3.9}$$

for an element $(d, \epsilon)\tilde{g}$ in the Levi part $M_{R_a}(\mathbb{A})$ with $d \in GL(l_a^+)_{\mathbb{A}}$ and $\tilde{g} \in \widetilde{Sp}(Z)_{\mathbb{A}}$. Note that $|\det_{l_a^+} d|^{n+(a+1)/2} = \delta_{R_a}^{\frac{1}{2}}(d)$.

PROPOSITION 3.3. *The integral (3.8) converges absolutely for $\text{Re}(s) > n + (a + 1)/2$, and continues to a meromorphic function in the whole plane. Its poles are contained in the set of poles of an Eisenstein series corresponding to a certain fixed (independent of $\tilde{\pi}$) degenerate principal series for $\widetilde{Sp}(Z \oplus Z)_{\mathbb{A}}$, induced from a Siegel parabolic subgroup.*

Proof. As in [Moe97a, § 2, (2)], let us rewrite

$$\begin{aligned} f_{\varphi_{\tilde{\pi},s}}(\tilde{g}_a) &= \int_{C_2 \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g}) f'_s(j_2(\tilde{g}') \tilde{j}_1(\tilde{g}_a)) d\tilde{g} \\ &= \int_{C_2 Sp(Z) \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g}) \sum_{\gamma \in Sp(Z)} f'_s((j_2(\gamma), 1) \tilde{j}_2(\tilde{g}') \tilde{j}_1(\tilde{g}_a)) d\tilde{g}. \end{aligned} \tag{3.10}$$

Note that $(\gamma, 1)' = (\gamma^c, 1)$, for $\gamma \in Sp(Z)$. Since $\gamma \mapsto j_2(\gamma)$ is an embedding of $Sp(Z)$ inside $P_{a,\Delta} \backslash Sp(Z_a \times Z)$, the inner sum in the integral (3.10) is a sub-series of the series defining the Eisenstein series $E'(j(\tilde{g}_a, \tilde{g}'), f'_s)$ as in (3.4), and hence it converges absolutely for $\text{Re}(s) > n + (a + 1)/2$.

In order to obtain the analytic continuation, we may assume that $\tilde{g}_a = \tilde{h}$ lies in $\widetilde{Sp}(Z)_{\mathbb{A}} \subset \tilde{M}_{R_a}(\mathbb{A})$. Then we saw that $\tilde{h} \mapsto f_{\varphi_{\tilde{\pi},s}}(\tilde{h})$ lies in the space of $\tilde{\pi}$. Thus, it is enough to show the analytic continuation of

$$\int_{C_2 Sp(Z) \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} f_{\varphi_{\tilde{\pi},s}}(\tilde{h}) \bar{\xi}_{\tilde{\pi}}(\tilde{h}) d\tilde{h}, \tag{3.11}$$

for every cusp form $\xi_{\tilde{\pi}}$. By using part (3) of Lemma 3.2, the integral (3.11), in its convergence domain, is equal to

$$\begin{aligned} &\int_{C_2 Sp(Z) \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \int_{C_2 \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g}) \bar{\xi}_{\tilde{\pi}}(\tilde{h}) f'_s(j_1(\tilde{h}) \tilde{j}_2(\tilde{g}')) d\tilde{g} d\tilde{h} \\ &= \int_{C_2 Sp(Z) \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \int_{C_2 \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g}) \bar{\xi}_{\tilde{\pi}}(\tilde{h}) f'_s(j_2(\tilde{h}^{-1} \tilde{g}')) d\tilde{g} d\tilde{h} \\ &= \int_{C_2 Sp(Z) \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \int_{C_2 \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{h}\tilde{g}) \bar{\xi}_{\tilde{\pi}}(\tilde{h}) f'_s(j_2(\tilde{g}')) d\tilde{g} d\tilde{h} \\ &= \int_{C_2 \backslash \widetilde{Sp}(Z)_{\mathbb{A}}} f'_s(j_2(\tilde{g}')) \langle \tilde{\pi}(\tilde{g}) \varphi_{\tilde{\pi}}, \xi_{\tilde{\pi}} \rangle_{L^2} d\tilde{g}, \end{aligned}$$

where $\langle \cdot, \cdot \rangle_{L^2}$ is the standard L^2 -product of cuspidal automorphic forms on $C_2\mathrm{Sp}(Z)\backslash\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$. Note that the switch of order of integrations $d\tilde{g}$ and $d\tilde{h}$ is justified, since $\varphi_{\tilde{\pi}}$ and $\xi_{\tilde{\pi}}$ are rapidly decreasing. Thus, the integral (3.11), using (3.8), is equal to

$$\int_{C_2\backslash\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}} f'_s(\tilde{j}_1(\tilde{g}))\langle\varphi_{\tilde{\pi}}, \tilde{\pi}(\tilde{g})\xi_{\tilde{\pi}}\rangle_{L^2} d\tilde{g}. \tag{3.12}$$

We recognize (3.12) as the result of unfolding the Rankin–Selberg integrals of the doubling method applied to metaplectic groups. This unfolding is quite formal, and is obtained exactly as in the linear case. See [PR86] and [LR05]. Thus, let f_s^* be the restriction of f'_s to $\widetilde{\mathrm{Sp}}(Z \oplus Z)_{\mathbb{A}}$. Then f_s^* is a holomorphic $\tilde{K}_{Z \times Z}$ -finite section in

$$\mathrm{Ind}_{\tilde{P}_{\Delta}(\mathbb{A})}^{\widetilde{\mathrm{Sp}}(Z \oplus Z)_{\mathbb{A}}} (\chi'_{V,\psi} |\det|^{s+a/2}).$$

Here $P_{\Delta} = P_{0,\Delta}$, and we think of $\mathrm{Sp}(Z \oplus Z)$ as $\mathrm{Sp}(Z_0 \oplus Z)$. Let $E^*(\tilde{h}, f_s^*)$ be the corresponding Eisenstein series on $\widetilde{\mathrm{Sp}}(Z \oplus Z)_{\mathbb{A}}$. Then (3.12) equals

$$\int_{C_2\mathrm{Sp}(Z) \times C_2\mathrm{Sp}(Z) \backslash \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}} \times \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}} \bar{\xi}_{\tilde{\pi}}(\tilde{g})\varphi_{\tilde{\pi}}(\tilde{h})E^*(\tilde{j}(\tilde{g}, \tilde{h}), f_s^*) d\tilde{g} d\tilde{h}. \tag{3.13}$$

This integral is meromorphic in the whole plane, and its poles are included in the set of poles of $E^*(\tilde{x}, f_s^*)$. □

Remark 3.4. Note that the integral (3.12) is Eulerian, and its computation at unramified places was carried out by Li in [Li92]. Also, it is easy to see that integral (3.5) is (for $a = 0$) an inner integral of (3.13).

Let $E^{Ra}(\tilde{h}, f_{\varphi_{\tilde{\pi},s}})$ denote the Eisenstein series on $\widetilde{\mathrm{Sp}}(Z_a)_{\mathbb{A}}$ corresponding to the section $f_{\varphi_{\tilde{\pi},s}}$ given by (3.8). This Eisenstein series will give another description of the subspace $\theta_{\psi,V}^{Z_a}(\theta_{\psi^{-1},Z}^V(\tilde{\pi}))$.

THEOREM 3.5. *In the notation above, we have*

$$\int_{C_2\mathrm{Sp}(Z) \backslash \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g})E'(\tilde{j}(\tilde{g}_a, \tilde{g}'), f'_s) d\tilde{g} = E^{Ra}(\tilde{g}, f_{\varphi_{\tilde{\pi},s}}). \tag{3.14}$$

Proof. The proof is exactly as in [Moe97a, § 2], which is a generalization of the doubling method. The reason for the exact similarity is the fact that $\mathrm{Sp}(Z)$ (respectively, $\mathrm{Sp}(Z_a \oplus Z)$) is a subgroup of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$ (respectively, $\widetilde{\mathrm{Sp}}(Z_a \oplus Z)_{\mathbb{A}}$), and similarly for unipotent radicals (over \mathbb{A}). Thus, when we unwind the left-hand side of (3.14), we have to examine the double coset space

$$P_{a,\Delta}\backslash\mathrm{Sp}(Z_a \oplus Z)/j(\mathrm{Sp}(Z_a) \times \mathrm{Sp}(Z)).$$

This is described in [PR86] and used in [Moe97a, § 2], and exactly as in [Moe97a, § 2] only one double coset has a non-trivial contribution to the left-hand side of (3.14), and it is the double coset of the identity element: $P_{a,\Delta}j(\mathrm{Sp}(Z_a) \times \mathrm{Sp}(Z))$. Again, the reason is that for the other double cosets, we get a stabilizer in $\mathrm{Sp}(Z_a) \times \mathrm{Sp}(Z)$, whose projection on $\mathrm{Sp}(Z)$ contains a unipotent radical, which contributes an inner integration of a cusp form of $\tilde{\pi}$ along this radical, and hence results in zero. Thus, we get, as in [Moe97a, § 2], that the left-hand side of (3.14) equals

$$\sum_{\gamma \in R_a(k) \backslash \mathrm{Sp}(Z_a)} \int_{C_2 \backslash \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}} \varphi_{\tilde{\pi}}(\tilde{g})f'_s(\tilde{j}((\gamma, 1)\tilde{g}_a, \tilde{g}')) d\tilde{g} = \sum_{\gamma \in R_a(k) \backslash \mathrm{Sp}(Z_a)} f_{\varphi_{\tilde{\pi},s}}((\gamma, 1)\tilde{g}_a) = E^{Ra}(\tilde{g}_a, f_{\varphi_{\tilde{\pi},s}}).$$

This proves the theorem. □

Using Proposition 3.1 and (3.6), we conclude as follows.

THEOREM 3.6. *Let $\tilde{\pi}$ be an irreducible genuine cuspidal automorphic representation of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$. Assume that $\theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ is cuspidal. Let a be a positive integer such that $m < n + a + 1$. Then the space $\theta_{\psi, V}^{Z_a}(\theta_{\psi^{-1}, Z}^V(\tilde{\pi}))$ is contained in the space of automorphic forms on $\widetilde{\mathrm{Sp}}(Z_a)_{\mathbb{A}}$ generated by the residual automorphic forms*

$$\tilde{g}_a \mapsto \mathrm{Res}_{s=(n+a+1-m)/2} E^{R_a}(\tilde{g}_a, f_{\varphi_{\tilde{\pi}}, s}). \tag{3.15}$$

3.2 Proof of Theorem 1.1

We are now ready to complete the proof of Theorem 1.1 stated in the introduction. The details are exactly as in [Moe97a, Moe97b]. We indicate them briefly. Let us apply the constant term along U_{R_a} (as an operator) to the inclusion asserted in Theorem 3.6. For an automorphic form ξ , on $\widetilde{\mathrm{Sp}}(Z_a)_{\mathbb{A}}$, denote by $\xi_{U_{R_a}}$ its constant term along U_{R_a} , restricted to the Levi part $\widetilde{M}_{R_a}(\mathbb{A})$. For an automorphic representation τ of $\widetilde{\mathrm{Sp}}(Z_a)_{\mathbb{A}}$, we denote by $\tau_{U_{R_a}}$ the automorphic representation of $\widetilde{M}_{R_a}(\mathbb{A})$ whose space consists all of the $\xi_{U_{R_a}}$, as ξ varies in the space of τ . Recall that M_{R_a} is isomorphic to $\mathrm{GL}(l_a^+) \times \mathrm{Sp}(Z)$. By Rallis’ tower property [Ral84], for any cuspidal automorphic representation σ of $\mathrm{O}(V)_{\mathbb{A}}$,

$$\mathrm{Res}_{\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}} [\theta_{\psi, V}^{Z_a}(\sigma)]_{U_{R_a}} = \theta_{\psi, V}^Z(\sigma). \tag{3.16}$$

Here we view $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$ as a subgroup of $\widetilde{M}_{R_a}(\mathbb{A})$. Moreover, $\widetilde{\mathrm{GL}}(l_a^+)_{\mathbb{A}}$ acts on $[\theta_{\psi, V}^{Z_a}(\sigma)]_{U_{R_a}}$ by the character

$$(d, \epsilon) \mapsto \epsilon \chi_{V, \psi}(\det_{l_a^+} d) |\det_{l_a^+} d|^{m/2} = \mu_{\psi}(d) |\det_{l_a^+} d|^{m/2}. \tag{3.17}$$

The main formula which explains (3.16) and (3.17) is that, for $\phi \in S(V_{\mathbb{A}}^{n+a})$, of the form $\phi_1 \otimes \phi_2$, where $\phi_1 \in S(V_{\mathbb{A}}^a)$ and $\phi_2 \in S(V_{\mathbb{A}}^n)$ (in the notation of (2.25)),

$$\theta_{\psi}^{\phi}(\varphi_{\sigma})_{U_{R_a}}((d, \epsilon)\tilde{g}) = \epsilon \phi_1(0) \mu_{\psi}(d) |\det(d)|^{m/2} \theta_{\psi}^{\phi_2}(\varphi_{\sigma})(\tilde{g}), \tag{3.18}$$

for $\tilde{g} \in \widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, $d \in \mathrm{GL}(l_a^+)_{\mathbb{A}}$, $\epsilon = \pm 1$. From Theorem 3.6, we conclude that, over $\widetilde{M}_{R_a}(\mathbb{A})$,

$$\mu_{\psi} |\det_{l_a^+}|^{m/2} \otimes \theta_{\psi, V}^Z(\theta_{\psi^{-1}, Z}^V(\tilde{\pi})) \subset \{[\mathrm{Res}_{s=(n+a+1-m)/2} E^{R_a}(\cdot, f_{\varphi_{\tilde{\pi}}, s})]_{U_{R_a}} \mid \varphi_{\tilde{\pi}} \in V_{\tilde{\pi}}\}. \tag{3.19}$$

Exactly as in [Moe97a, § 2], by expressing the constant term along U_{R_a} , $[E^{R_a}(\cdot, f_{\varphi_{\tilde{\pi}}, s})]_{U_{R_a}}$, of the Eisenstein series $E^{R_a}(\cdot, f_{\varphi_{\tilde{\pi}}, s})$ in terms of intertwining operators, we can replace the right-hand side of (3.19) by a certain sum of residues of certain intertwining operators, which take values in the space of $\tilde{\pi}$. The details are exactly as in [Moe97a, § 2], only that we replace the characters

$$\mathrm{diag}(t_1, \dots, t_a) \mapsto |t_1|^{s_1} \cdots |t_a|^{s_a}$$

there by (our) characters,

$$(\mathrm{diag}(t_1, \dots, t_a), \epsilon) \mapsto \epsilon \chi_{V, \psi}(t_1 \cdots t_a) |t_1|^{s_1} \cdots |t_a|^{s_a}.$$

We then conclude from (3.19), by considering only the action of $\widetilde{\mathrm{Sp}}(Z)_{\mathbb{A}}$, that

$$\theta_{\psi, V}^Z(\theta_{\psi^{-1}, Z}^V(\tilde{\pi})) = \tilde{\pi}.$$

This concludes the proof of part (1) of Theorem 1.1.

The proof of part (2) of Theorem 1.1 now follows, as in [Moe97a, § 2], the only interesting case being $b \leq m - n - 1$. Otherwise, we just apply Theorem 3.6 for $a = b$. For a fixed integer $a > m - n - 1$, we follow the same process as in the proof of part (1) of Theorem 1.1 above. This time we apply the constant term along $U_{R_{a-b}}$ (as an operator) to the inclusion asserted in Theorem 3.6, viewing Z_a as $(Z_{a-b})_b$. This will prove part (2) of Theorem 1.1.

Finally, part (3) of Theorem 1.1 follows immediately from part (1) of Theorem 1.1 and the Rallis tower property [Ral84]. Indeed, since $\theta_{\psi^{-1}, Z}^V(\tilde{\pi})$ is assumed to be cuspidal, then any irreducible

sub-representation of it, σ , is cuspidal. Then $\theta_{\psi,V}^Z(\sigma)$ is non-trivial and is a sub-representation of $\theta_{\psi,V}^Z(\theta_{\psi^{-1},Z}^V(\tilde{\pi})) = \tilde{\pi}$, and hence equals $\tilde{\pi}$ and, in particular, is cuspidal. By the Rallis tower property [Ral84], $\theta_{\psi,V}^{Z-b}(\sigma) = 0$ for all positive integers b . This proves part (3) of Theorem 1.1 and hence completes the proof of the whole theorem.

3.3 Proof of Theorems 1.2 and 1.3

The proofs of these two theorems can be directly inferred from [Moe97b]. For this, let us explain how to proceed until we reach the point where we can directly use the results of [Moe97b].

Let σ be an irreducible cuspidal automorphic representation of $O(V)_{\mathbb{A}}$. Assume that $\theta_{\psi^{-1},V}^Z(\sigma)$ is cuspidal, as an automorphic representation of $\widetilde{Sp}(Z)_{\mathbb{A}}$. Let V_b be the orthogonal direct sum of the quadratic space V (equipped with its symmetric form Q) and b hyperbolic planes $ku_i + ku_{-i}$, $i = 1, \dots, b$. As usual, u_i and u_{-i} are isotropic, and the Gram matrix of the pair (u_i, u_{-i}) is $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

Put $\ell_b = \bigoplus_{i=1}^b(ku_i + ku_{-i})$ and $\ell_b^{\pm} = \bigoplus_{i=1}^b ku_{\pm i}$. Denote by Q_b the symmetric bilinear form on V_b . Consider the theta lift $\theta_{\psi,Z}^{V_b}(\theta_{\psi^{-1},V}^Z(\sigma))$. Its space is generated by the following automorphic forms on $O(V_b)_{\mathbb{A}}$

$$\int_{C_2Sp(Z)\backslash\widetilde{Sp}(Z)_{\mathbb{A}}} \theta_{\psi}^{\phi_2}(\tilde{g}, h_b) \int_{O(V)\backslash O(V)_{\mathbb{A}}} \theta_{\psi^{-1}}^{\phi_1}(\tilde{g}, h) \varphi_{\sigma}(h) dh d\tilde{g}. \tag{3.20}$$

Here $h_b \in O(V_b)_{\mathbb{A}}$, $\phi_1 \in \mathcal{S}(V_{\mathbb{A}}^n)$, $\phi_2 \in \mathcal{S}(V_b(\mathbb{A})^n)$, and φ_{σ} is a cusp form in the space of σ . Now, let us view $\omega_{\psi^{-1},Z \otimes V}$ as $\omega_{\psi,Z \otimes V'}$, where V' denotes the space V , equipped with the symmetric form $-Q$. (See [MVW87, p. 36].) We have a natural embedding over k

$$O(V_b) \times O(V') \rightarrow O(V_b \oplus V'),$$

where the symmetric form of $V_b \oplus V'$ is $Q_b \oplus (-Q)$. Note that $V_b \oplus V'$ is a totally split, even-dimensional quadratic space over k . We have [MVW87, p. 37]

$$\theta_{\psi,Z \otimes V_b}^{\phi_2}(\tilde{g}, h_b) \theta_{\psi,Z \otimes V'}^{\phi_1}(\tilde{g}, h) = \theta_{\psi,Z \otimes (V_b \oplus V')}^{\phi_2 \otimes \phi_1}(g, (h_b, h)),$$

where $g \in Sp(Z)_{\mathbb{A}}$ is the projection of \tilde{g} , and we think of $\phi_2 \otimes \phi_1$ as an element of $\mathcal{S}((V_b \oplus V')_{\mathbb{A}}^n)$. Thus, (3.20) becomes

$$\int_{Sp(Z)\backslash Sp(Z)_{\mathbb{A}}} \int_{O(V')\backslash O(V')_{\mathbb{A}}} \theta_{\psi,Z \otimes (V_b \oplus V')}^{\phi_2 \otimes \phi_1}(g, (h_b, h)) \varphi_{\sigma}(h) dh dg. \tag{3.21}$$

Now, we are at the situation of [Moe97a, § 1]. Indeed, we assume that $2n < m + b - 1$. By applying Theorem 2.4 with regularization in terms of a p -adic Hecke element α , which is the same as in the proof in [Moe97a], the integral (3.21) has the form

$$\int_{O(V')\backslash O(V')_{\mathbb{A}}} \varphi_{\sigma}(h) \text{Res}_{s=(m+b-1)/2-n} E((h_b, h), \xi_s) dh. \tag{3.22}$$

Here ξ_s is a $K_{V_b \oplus V'}$ -finite section of $\text{Ind}_{L_{b,\Delta}(\mathbb{A})}^{O(V_b \oplus V')_{\mathbb{A}}} |\det_{V_b,\Delta}|^s$, and $E(\cdot, \xi_s)$ denotes the corresponding Eisenstein series. We denote by $L_{b,\Delta}$ the maximal parabolic subgroup of $O(V_b \oplus V')$, which preserves the isotropic subspace

$$V_{b,\Delta} = \bigoplus_{i=1}^b k(u_i, 0) + \{(v, v) \in V \times V'\}.$$

The proof proceeds from this point on, word by word, as in [Moe97a, § 1]. Note that the parity of m is no longer relevant to the rest of the proof. First, we conclude, as in the previous section, as follows.

THEOREM 3.7. *Let σ be an irreducible cuspidal automorphic representation of $\mathrm{O}(V)_{\mathbb{A}}$. Assume that $\theta_{\psi^{-1}, V}^Z(\sigma)$ is cuspidal. Then $\theta_{\psi, Z}^{V_b} \theta_{\psi^{-1}, V}^Z(\sigma)$ is contained in the space of residues at $s = (m+b-1)/2-n$ of the Eisenstein series corresponding to $\mathrm{Ind}_{D_b(\mathbb{A})}^{\mathrm{O}(V_b)_{\mathbb{A}}} |\det_{\ell_b^+}|^s$, where D_b is the parabolic subgroup of $\mathrm{O}(V_b)$, which preserves ℓ_b^+ .*

Now Theorem 1.2 in the introduction follows in exactly the same way as the proof of Theorem 1.1 in the previous section. Theorem 1.3 in the introduction follows from Theorems 1.1 and 1.2, exactly as in [Moe97a, §2]. We omit the details.

4. Applications

In this section, we let $H_n = \mathrm{SO}_{2n+1} = \mathrm{SO}(V)$, where (V, Q) is the column space of dimension $2n + 1$ over k , equipped with the non-degenerate k -split symmetric bilinear form in $2n + 1$ variables associated to the symmetric matrix defined inductively by

$$J_{2n+1} = \begin{pmatrix} 0 & & 1 \\ & J_{2n-1} & \\ 1 & & 0 \end{pmatrix}.$$

We denote by $\tilde{G}_n(k_\nu)$ and $\tilde{G}_n(\mathbb{A})$ the local and the global metaplectic groups, respectively, corresponding to the symplectic group $G_n = \mathrm{Sp}_{2n}$ over k , with respect to the standard symplectic form on the $2n$ -dimensional row space. For an irreducible cuspidal automorphic representation σ of $H_n(\mathbb{A})$, we denote by $\tilde{\theta}_{\psi, n}^m(\sigma)_+$ the ψ -theta lift of σ to $\tilde{G}_m(\mathbb{A})$; it is generated by the automorphic forms $\theta_{\psi}^{\phi}(\varphi_{\sigma})_+$, which are defined as in (2.18), only that the integration in (2.18) is replaced by an integration over $H_n(k) \backslash H_n(\mathbb{A})$. Similarly, for an irreducible genuine cuspidal automorphic representation $\tilde{\pi}$ of $\tilde{G}_m(\mathbb{A})$, we denote by $\theta_{\psi, m}^n(\tilde{\pi})$ the ψ -theta lift of $\tilde{\pi}$ to $\mathrm{O}_{2n+1}(\mathbb{A})$. We denote by $\theta_{\psi, m}^n(\tilde{\pi})_+$ the automorphic representation of $H_n(\mathbb{A})$ obtained by restricting the functions of $\theta_{\psi, m}^n(\tilde{\pi})$ to $H_n(\mathbb{A})$.

Our first application is the proof of Theorem 1.4 in the introduction, which is part (4) of the main theorem in [Fur95] without the assumptions made there.

THEOREM 4.1. *Let σ be an irreducible cuspidal automorphic representation of $H_n(\mathbb{A})$. Assume that σ is generic, i.e. has a nonzero Whittaker Fourier coefficient. Then the ψ -theta lift of σ to $\tilde{G}_n(\mathbb{A})$, $\tilde{\theta}_{\psi, n}^n(\sigma)_+$, is non-trivial if and only if $L(\sigma, \frac{1}{2}) \neq 0$, where $L(\sigma, s)$ is the standard L -function attached to σ .*

Proof. Since σ is cuspidal and generic, the local components of σ are all unitary and locally generic, and hence, by the exponent structure of the generic unitary dual [LMT04], $L(\sigma_{\nu}, s)$ is holomorphic at $s = \frac{1}{2}$, for all places ν . Thus, $L(\sigma, \frac{1}{2}) \neq 0$, if and only if $L^S(\sigma, \frac{1}{2}) \neq 0$, for any finite set S of places of k , which includes the set of all archimedean places of k . This can also be deduced from the explicit Langlands functorial transfer from $\mathrm{SO}_{2n+1}(\mathbb{A})$ to $\mathrm{GL}_{2n}(\mathbb{A})$ for irreducible generic cuspidal automorphic representations σ of $\mathrm{SO}_{2n+1}(\mathbb{A})$, which has been established through [CKPS04, GRS01, JS03, JS04].

Choose S , such that σ and ψ are unramified outside S . If $L^S(\sigma, \frac{1}{2}) \neq 0$, then by [JS, Remark 2.3], σ has a Bessel model of special type with respect to $\mathrm{SO}_{1,1}$. Then by [Fur95, Proposition 1] (with $\lambda = 1$), $\tilde{\theta}_{\psi, n}^n(\sigma)_+$ has a non-trivial ψ -Whittaker Fourier coefficient, and hence $\tilde{\theta}_{\psi, n}^n(\sigma)_+$ is nonzero. Note that, for this direction, we did not use any of the results proved in the previous sections.

Conversely, assume that $\tilde{\theta}_{\psi, n}^n(\sigma)_+ \neq 0$. Then we claim that $\tilde{\theta}_{\psi, n}^n(\sigma)_+$ is cuspidal. Indeed, if this is not so, then the Rallis tower property [Ral84] (which is clearly valid for special orthogonal groups as well) implies that, for the first integer $m > 0$ such that $\tilde{\theta}_{\psi, n}^m(\sigma)_+ \neq 0$, the

representation $\tilde{\pi}' = \tilde{\theta}_{\psi,n}^m(\sigma)_+$ is cuspidal. Choose an irreducible summand $\tilde{\pi}$ of $\tilde{\pi}'$. In particular, at a given finite place ν of k , we get that $\tilde{\pi}_\nu$ is a local ψ_ν -Howe lift of the generic σ_ν , but this is impossible by [JS03, Proposition 2.1], since $m < n$. This proves the claim.

Let us denote by $Z_2 = \{\pm I_{2n+1}\}$ the center of O_{2n+1} . Note that $O_{2n+1} = Z_2 \times H_n$. Let μ be a character of $Z_2 \backslash Z_2(\mathbb{A})$, and denote by σ_μ the extension of σ to $O_{2n+1}(\mathbb{A})$ by the character μ on the center. This also allows us to extend cusp forms of σ from $H_n(\mathbb{A})$ to $O_{2n+1}(\mathbb{A})$ in the same way. The space of $\tilde{\theta}_{\psi,n}^n(\sigma)_+$ is generated by the integrals $\theta_\psi^\phi(\varphi_\sigma)(\tilde{g})_+$. It is clear that

$$\theta_\psi^\phi(\varphi_\sigma)(\tilde{g})_+ = \sum_\mu \int_{O_{2n+1}(F) \backslash O_{2n+1}(\mathbb{A})} \theta_\psi^\phi(\tilde{g}, h) \varphi_{\sigma_\mu}(h) dh.$$

Thus, it follows that there is μ such that $\tilde{\pi} = \tilde{\theta}_{\psi,n}^n(\sigma_\mu) \neq 0$ and is, of course, cuspidal. By Theorem 1.2, we know that $\tilde{\pi}$ is irreducible and $\sigma_\mu = \theta_{\psi^{-1},n}^n(\tilde{\pi})$. By [Fur95, Proposition 3], we get that $\tilde{\pi}$ is ψ -generic since σ is generic. Note that by restricting to $H_n(\mathbb{A})$, we get that $\sigma = \theta_{\psi^{-1},n}^n(\tilde{\pi})_+$. By [CKPS04], σ has a functorial lift to $GL_{2n}(\mathbb{A})$ and by [GRS01, Sou05], this lift has the form $\tau = \tau_1 \boxplus \dots \boxplus \tau_l$, which is an isobaric sum of l pairwise non-equivalent, irreducible, cuspidal, automorphic representations τ_i of $GL_{2n_i}(\mathbb{A})$, with the property that $L^S(\tau_i, \Lambda^2, s)$ has a pole at $s = 1$ for each i . It follows that with respect to ψ , $\tilde{\pi}$ lifts at almost all places to τ , and by [Sou05, Theorem 13], we conclude also that $L^S(\tau_i, \frac{1}{2}) \neq 0$ for all i . This implies that $L^S(\sigma, \frac{1}{2}) \neq 0$. \square

Next, we are going to consider the cases when the central value of the standard L -function $L^S(\sigma, \frac{1}{2})$ is zero.

Let σ be an irreducible generic cuspidal automorphic representation of $H_n(\mathbb{A})$. By [HP83, Theorem 5.7], the ψ -theta lift of σ to $\tilde{G}_{n+1}(\mathbb{A})$, $\tilde{\theta}_{\psi,n}^{n+1}(\sigma)_+$, is nonzero and admits non-trivial ψ -Whittaker Fourier coefficients. The calculation of this ψ -Whittaker Fourier coefficient can also be found in [Fur95, § 4]; its precise local variant appears in [JS03, Corollary 2.2]. The precise formula for the ψ -Whittaker Fourier coefficient of automorphic functions in $\tilde{\theta}_{\psi,n}^{n+1}(\sigma)_+$ is as follows.

Let U be the standard maximal unipotent subgroup of G_n . Since the Rao cocycle is trivial on U at all places, we may regard $U_{\mathbb{A}}$ as a subgroup of $\tilde{G}_n(\mathbb{A})$. Denote by ψ_U the standard non-degenerate (or Whittaker) character of $U_{\mathbb{A}}$ given by evaluating ψ at the sum of all simple root coordinates of elements of $U_{\mathbb{A}}$. Let φ_σ be a cusp form in the space of σ . Let $\omega_{\psi,V}^{n+1}$ denote the restriction of the ψ -Weil representation to the subgroup $\tilde{G}_{n+1}(\mathbb{A}) \cdot H_n(\mathbb{A})$, realized in the Schrödinger model $\mathcal{S}(V_{\mathbb{A}}^{n+1})$, where V is the $(2n + 1)$ -dimensional column space, over k , on which H_n acts. Let $\{v_1, \dots, v_n, v_0, v_{-n}, \dots, v_{-1}\}$ be the standard basis of V over k , i.e. the spans of the first n vectors and of the last n vectors are maximal totally isotropic subspaces, both orthogonal to the vector v_0 , $Q(v_i, v_{-j}) = \delta_{i,j}$ for $i, j \leq n$, and $Q(v_0, v_0) = 1$. For $\phi \in \mathcal{S}(V_{\mathbb{A}}^{n+1})$, we denote, as before but with more precision, by $\tilde{\theta}_{\psi,n}^{\phi,n+1}(\varphi_\sigma)_+$ the ψ -theta lift of the cusp form φ_σ in the space of σ to $\tilde{G}_{n+1}(\mathbb{A})$, which is defined as in (2.18), but with the integration taking place over $H_n(k) \backslash H_n(\mathbb{A})$. Consider the ψ -Whittaker Fourier coefficient of $\tilde{\theta}_{\psi,n}^{\phi,n+1}(\varphi_\sigma)_+$

$$\mathcal{W}_{\tilde{\theta}_{\psi,n}^{\phi,n+1}(\varphi_\sigma)_+}^\psi(\tilde{g}) = \int_{U \backslash U_{\mathbb{A}}} \tilde{\theta}_{\psi,n}^{\phi,n+1}(\varphi_\sigma)_+(u\tilde{g}) \psi^{-1}(u) du.$$

Then we have the following expression

$$\mathcal{W}_{\tilde{\theta}_{\psi,n}^{\phi,n+1}(\varphi_\sigma)_+}^\psi(\tilde{g}) = \int_{C_n(\mathbb{A}) \backslash H_n(\mathbb{A})} \omega_{\psi,V}^{n+1}(\tilde{g}, h) \phi(v_1, \dots, v_n, v_0) \mathcal{W}_{\varphi_\sigma}^\psi(h) dh. \tag{4.1}$$

Here $\mathcal{W}_{\varphi_\sigma}^\psi$ is the standard ψ -Whittaker Fourier coefficient of φ_σ , and C_n is the pointwise stabilizer in H_n of v_1, \dots, v_n, v_0 . Clearly, the right-hand side of (4.1) is not identically zero, since σ is generic.

Thus, if $L(\sigma, \frac{1}{2}) = 0$, then by Theorem 4.1, the ψ -theta lift to $\tilde{G}_n(\mathbb{A})$, $\tilde{\theta}_{\psi,n}^n(\sigma)_+$, vanishes. Then by the Rallis tower property [Ral84] the ψ -theta lift to $\tilde{G}_{n+1}(\mathbb{A})$, $\tilde{\theta}_{\psi,n}^{n+1}(\sigma)_+$ is cuspidal. As we did before, there is a character μ of $Z_2 \backslash Z_2(\mathbb{A})$ such that $\tilde{\theta}_{\psi,n}^{n+1}(\sigma_\mu) = \tilde{\pi}$ is irreducible, cuspidal, and ψ -generic.

Conversely, let $\tilde{\pi}$ be an irreducible genuine cuspidal automorphic representation of $\tilde{G}_{n+1}(\mathbb{A})$. Assume that $\tilde{\pi}$ is ψ -generic. If the ψ^{-1} -theta lift of $\tilde{\pi}$ to $O_{2n+1}(\mathbb{A})$, $\sigma' = \theta_{\psi^{-1},n+1}^n(\tilde{\pi})$, does not vanish, then σ' is cuspidal, according to [JS03, Corollary 2.2(2)]. By Theorems 1.1 and 1.3, we get that σ' is irreducible, and hence it is of the form σ_μ with notation as above. We conclude that $\tilde{\pi} = \tilde{\theta}_{\psi,n}^{n+1}(\sigma_\mu)$. Since $\tilde{\pi}$ is ψ -generic, then we can compute the ψ -Whittaker Fourier coefficient on $\tilde{\pi}$ as we did in (4.1), and obtain the same formula, except that, in the right-hand side of (4.1), we integrate over $C_n(\mathbb{A}) \backslash O_{2n+1}(\mathbb{A})$. In particular, $\mathcal{W}_{\phi_\sigma}^\psi$ is not identically zero, and this means that $\theta_{\psi,n+1}^n(\tilde{\pi})_+$ is generic. From Theorem 4.1, we also conclude that $L(\sigma, \frac{1}{2}) = 0$. Let us summarize the above discussion as follows.

THEOREM 4.2. *With notation above, the following hold.*

- (1) *Let σ be an irreducible generic cuspidal automorphic representation of $H_n(\mathbb{A})$. Assume that $L(\sigma, \frac{1}{2}) = 0$. Then $\tilde{\theta}_{\psi,n}^{n+1}(\sigma)_+$ is cuspidal and ψ -generic.*
- (2) *Let $\tilde{\pi}$ be an irreducible genuine ψ -generic cuspidal automorphic representation of \tilde{G}_{n+1} . Assume that $\sigma = \theta_{\psi^{-1},n+1}^n(\tilde{\pi})_+$ is non-trivial. Then σ is irreducible, cuspidal, and generic, such that $L(\sigma, \frac{1}{2}) = 0$.*

Finally, we are ready to complete our proof of Theorem 1.5. We are going to use the notation introduced in [JS] freely.

Let σ be an irreducible cuspidal automorphic representation of $H_n(\mathbb{A})$, which has a nonzero Bessel model of special type, i.e. of type $(D_\lambda, 1, \psi_{n,n-1;\lambda})$, with respect to the quadratic extension of k generated by the square root of $\lambda \in k^\times \setminus (k^\times)^2$. See [JS, § 2.2] for the definition of this notion. By [JS, Theorem 4.1(1)], we know that $\tilde{\theta}_{\psi,n}^k(\sigma)_+ = 0$, for $k < n - 1$, and by [Fur95, Proposition 1], the last space is non-trivial for $k = n$.

Assume now that $\tilde{\theta}_{\psi,n}^{n-1}(\sigma)_+ \neq 0$. Then, as before, we can find a character μ of $Z_2 \backslash Z_2(\mathbb{A})$ such that $\tilde{\tau} = \tilde{\theta}_{\psi,n}^{n-1}(\sigma_\mu)$ is cuspidal and irreducible. Again it follows from Theorems 1.2 and 1.3 that $\sigma_\mu = \theta_{\psi^{-1},n-1}^n(\tilde{\tau})$, and then that $\sigma = \theta_{\psi^{-1},n-1}^n(\tilde{\tau})_+$. Now, we can compute explicitly the Bessel model of special type of $\sigma = \theta_{\psi^{-1},n-1}^n(\tilde{\tau})_+$. It is defined by the integral in [JS, (2.11)], taking there $r = n - 1$ and the automorphic form ϕ to be the constant function 1. Then the calculation is exactly as in the p -adic case in [JS, Proof of Theorem 4.3]. We get that the Bessel model of special type (given by the integral) of $\sigma = \theta_{\psi^{-1},n-1}^n(\tilde{\tau})_+$ can be expressed in terms of the ψ_λ -Whittaker Fourier coefficient of $\tilde{\tau}$. Hence, $\tilde{\tau}$ must be ψ_λ -generic. (This calculation shows that starting with an irreducible, automorphic, cuspidal representation $\tilde{\tau}$ of $\tilde{G}_{n-1}(\mathbb{A})$, which is ψ_λ -generic, $\theta_{\psi^{-1},n-1}^n(\tilde{\tau})_+$ has a non-trivial Bessel model of special type, with respect to λ , as above, and, in particular, this theta-lift is nontrivial.) This proves the following theorem.

THEOREM 4.3. *Let σ be an irreducible cuspidal automorphic representation of $H_n(\mathbb{A})$. Assume that σ has a nonzero Bessel model of special type, i.e. of type $(D_\lambda, 1, \psi_{n,n-1;\lambda})$, with a nonsquare $\lambda \in k^\times$. If the ψ -theta lift to $\tilde{G}_{n-1}(\mathbb{A})$, $\tilde{\theta}_{\psi,n}^{n-1}(\sigma)_+$, does not vanish, then the automorphic representation $\tilde{\theta}_{\psi,n}^{n-1}(\sigma)_+$ of $\tilde{G}_{n-1}(\mathbb{A})$ is cuspidal and ψ_λ -generic.*

We briefly recall, from [JS], what we have proved for irreducible cuspidal automorphic representations σ of $H_n(\mathbb{A})$, which have a nonzero Bessel model of special type, i.e. of type $(D_\lambda, 1, \psi_{n,n-1;\lambda})$, with $\lambda \in k^\times$. We proved in [JS, § 2.3] that if this special Bessel model is k -split, i.e. $\lambda \in (k^\times)^2$, then the automorphic representation σ is generic [JS, Proposition 2.2].

If the special Bessel model of type $(D_\lambda, 1, \psi_{n,n-1;\lambda})$ is not k -split, i.e. λ is not a square, then we proved in [JS, Theorem 4.1] that the first occurrence of the ψ -theta lift of σ to the $\tilde{G}_m(\mathbb{A})$ -tower, i.e. the first index $m = m_\psi(\sigma)$, such that $\tilde{\theta}_{\psi,n}^m(\sigma)_+$ is nonzero, satisfies the following inequalities

$$n - 1 \leq m_\psi(\sigma) \leq n.$$

Then we proved that if $m_\psi(\sigma) = n$, then there are two possibilities. The first is that σ is nearly equivalent to an irreducible generic cuspidal automorphic representation σ'_n of $H_n(\mathbb{A})$, such that $\sigma'_n \otimes \chi_\lambda$ is in the image of the ψ^{-1} -theta lift, from irreducible, genuine, automorphic, cuspidal, ψ -generic representations of $\tilde{G}_n(\mathbb{A})$. In this case, we conclude, by Theorem 4.1, that $L(\sigma' \otimes \chi_\lambda, \frac{1}{2}) \neq 0$. Here χ_λ is the composition of α_λ with the spinor norm, and α_λ is the character given by the Hilbert symbol $(\cdot, \lambda)_k$.

The second possibility is that σ is a CAP representation with respect to the cuspidal data

$$(P_1; \alpha_\lambda | \cdot |^{\frac{1}{2}} \otimes \sigma_{n-1}),$$

where P_1 is the standard parabolic subgroup of H_n , which preserves an isotropic line, and σ_{n-1} is an irreducible cuspidal automorphic representation of $H_{n-1}(\mathbb{A})$, such that $\sigma_{n-1} \otimes \chi_\lambda$ is in the image of the ψ^{-1} -theta lift from an irreducible, genuine, automorphic, cuspidal, ψ -generic representation of $\tilde{G}_n(\mathbb{A})$. Note that in [JS, Theorem 4.1(2)], we were unable to determine further explicit properties of the cuspidal data, in particular, of σ_{n-1} , so that we were unable to complete the proof of the CAP conjecture [JS, Conjecture 1.1] for this case. Now by Theorem 4.2, we conclude that σ_{n-1} is generic and that the central value of the standard L -function twisted by χ_λ , $L(\sigma_{n-1} \otimes \chi_\lambda, \frac{1}{2})$, is zero. This proves a little more than what the CAP conjecture asserts in this case.

Now, let us analyze the remaining case, where the first occurrence $m_\psi(\sigma)$ is $n-1$. By Theorem 4.3, we conclude that the cuspidal automorphic representation $\tilde{\theta}_{\psi,n}^{n-1}(\sigma)_+$ is ψ_λ -generic. Thus, there is an irreducible ψ_λ -generic cuspidal automorphic representation $\tilde{\tau}$ of $\tilde{G}_{n-1}(\mathbb{A})$ such that $\sigma = \theta_{\psi^{-1},n-1}^n(\tilde{\tau})_+$ (by Theorems 1.2 and 1.3). In this case, we proved in [JS, Theorem 4.6] that either σ is CAP with respect to the cuspidal data

$$(P_1; | \cdot |^{\frac{1}{2}} \otimes \sigma_{n-1}),$$

or σ is CAP with respect to the cuspidal data

$$(P_{1,1}; | \cdot |^{\frac{1}{2}} \otimes \alpha_\lambda | \cdot |^{\frac{1}{2}} \otimes \sigma_{n-2}).$$

In the first case σ_{n-1} is cuspidal and generic on $H_{n-1}(\mathbb{A})$. The proof there shows, in this case, that $\sigma_{n-1} \otimes \chi_\lambda$ is the image under the ψ^{-1} -theta lift of the ψ -generic representation $\tilde{\tau}^\lambda$, which is an outer conjugation of $\tilde{\tau}$ by the similitude element $\text{diag}(I_n, \lambda I_n)$. By Theorem 4.1, we get that the central value of the standard L -function twisted by χ_λ , $L(\sigma_{n-1} \otimes \chi_\lambda, \frac{1}{2})$ is nonzero.

In the second case, $P_{1,1}$ is the standard parabolic subgroup whose Levi part is isomorphic to $GL_1 \times GL_1 \times H_{n-2}$, and σ_{n-2} is an irreducible cuspidal automorphic representation of $H_{n-2}(\mathbb{A})$. By construction,

$$\sigma_{n-2} \otimes \chi_\lambda = \tilde{\theta}_{\psi,n-1}^{n-2}(\tilde{\tau}^\lambda)_+.$$

Again, in [JS] we were unable to prove further properties for σ_{n-2} , so that the CAP conjecture can be completely verified for this case. Now, by Theorem 4.2, σ_{n-2} is also generic and the central value of the standard L -function twisted by χ_λ , $L(\sigma_{n-2} \otimes \chi_\lambda, \frac{1}{2})$ must be zero.

The proof of Theorem 1.5 is finally completed.

ACKNOWLEDGEMENTS

Some of the work related to this paper was carried out during our participation to the special program at ESI, 2006. We would like to thank the organizers S. Kudla, M. Rapoport and J. Schwermer for their warm invitation and thank ESI for the productive research environment and hospitality. Finally, we would like to thank the referee for very helpful comments.

REFERENCES

- CKPS04 J. Cogdell, H. Kim, I. Piatetski-Shapiro and F. Shahidi, *Functoriality for classical groups*, Publ. Math. Inst. Hautes Études Sci. **99** (2004), 163–233.
- Fur95 M. Furusawa, *On the theta lift from SO_{2n+1} to \widetilde{Sp}_n* , J. reine angew. Math. **466** (1995), 87–110.
- GRS01 D. Ginzburg, S. Rallis and D. Soudry, *Generic automorphic forms on $SO(2n + 1)$: functorial lift to $GL(2n)$, endoscopy, and base change*, Internat. Math. Res. Notices **14** (2001), 729–764.
- HP83 H. Howe and I. Piatetski-Shapiro, *Some examples of automorphic forms on Sp_4* , Duke Math. J. **50** (1983), 55–106.
- How79 R. Howe, *θ -series and invariant theory*, in *Automorphic forms, representations, and L-functions*, Proceedings of Symposia in Pure Mathematics Vol. 33, Part I (American Mathematical Society, Providence, RI, 1979), 275–285.
- Ioh01 A. Ichino, *On a regularized Siegel–Weil formula*, J. reine angew. Math. **539** (2001), 201–234.
- JS03 D. Jiang and D. Soudry, *The local converse theorem for $SO(2n + 1)$ and applications*, Ann. of Math. (2) **157** (2003), 743–806.
- JS04 D. Jiang and D. Soudry, *Generic representations and local Langlands reciprocity law for p -adic SO_{2n+1}* , in *Contributions to Automorphic Forms, Geometry and Number Theory, A volume in honor of Joseph S. Shalika*, eds H. Hida, D. Ramakrishnan and F. Shahidi (The Johns Hopkins University Press, Baltimore, MD, 2004), 457–519.
- JS D. Jiang and D. Soudry, *On the genericity of cuspidal automorphic forms on SO_{2n+1}* , J. reine angew. Math., to appear.
- Kud94 S. Kudla, *Splitting metaplectic covers of dual reductive pairs*, Israel J. Math. **87** (1994), 361–401.
- Kud96 S. Kudla, *Notes on the local theta correspondence*, in *Lecture series in the European School in Group Theory*, Schloss Hirschberg, September 1996, <http://www.math.umd.edu/~ssk/ssk.research.html>.
- KR90a S. Kudla and S. Rallis, *Degenerate principal series and invariant distributions*, Israel J. Math. **69** (1990), 25–45.
- KR90b S. Kudla and S. Rallis, *Poles of Eisenstein series and L-functions*, Israel Math. Conf. Proc. **3** (1990), 81–110.
- KR94 S. Kudla and S. Rallis, *A regularized Siegel–Weil formula: the first term identity*, Ann. of Math. (2) **140** (1994), 1–80.
- LMT04 E. Lapid, G. Muić and M. Tadić, *On the generic unitary dual of quasisplit classical groups*, Internat. Math. Res. Notices **26** (2004), 1335–1354.
- LR05 E. Lapid and S. Rallis, *On the local factors of representations classical groups*, in *Automorphic representations, L-functions and applications: Progress and Prospects. Proc. conf. honoring Steve Rallis*, Ohio State University Mathematical Research Institute Publications, vol. 11, eds J. Cogdell, D. Jiang, S. Kudla, D. Soudry and R. Stanton (de Gruyter, Berlin, 2005), 309–360.
- Li92 J.-S. Li, *Non-vanishing theorems for the cohomology of certain arithmetic quotients*, J. reine angew. Math. **428** (1992), 177–217.
- Moe97a C. Moeglin, *Non nullité de certains relèvements par séries théta*, J. Lie Theory **7** (1997), 201–229.
- Moe97b C. Moeglin, *Quelques propriétés de base des séries théta*, J. Lie Theory **7** (1997), 231–238.
- MVW87 C. Moeglin, M.-F. Vignéras and J.-L. Waldspurger, *Correspondances de Howe sur un corps p -adique*, Lecture Notes in Mathematics, vol. 1291 (Springer, Berlin, 1987).

- PR86 I. Piatetski-Shapiro and S. Rallis, ϵ factors for representations of classical groups, Proc. Natl. Acad. Sci. USA **83** (1986), 4589–4593.
- Ral82 S. Rallis, Langlands functoriality and the Weil representation, Amer. J. Math. **104** (1982), 469–517.
- Ral84 S. Rallis, On the Howe duality conjecture, Compositio Math. **51** (1984), 333–399.
- Ran93 R. Ranga Rao, On some explicit formulas in the theory of Weil representation, Pacific J. Math. **157** (1993), 335–371.
- Sou05 D. Soudry, On Langlands functoriality from classical groups to GL_n , in *Formes Automorphes (I). Actes du semestre du CEB, printemps 2000*, eds J. Tilouine, H. Carayol, M. Harris and M.-F. Vigneras, Astérisque **298** (2005), 335–390.
- Szp06 D. Szpruch, Uniqueness of Whittaker model for the metaplectic group, Preprint (2006).
- Zhu07 S. T. Lee and C.-B. Zhu, Degenerate principal series and local theta correspondence III: The case of complex groups, Preprint (2007).

Dihua Jiang dhjiang@math.umn.edu

School of Mathematics, University of Minnesota, 206 Church Street SE, Minneapolis, MN 55455, USA

David Soudry soundry@post.tau.ac.il

School of Mathematical Sciences, Tel Aviv University, Tel Aviv, 69978, Israel