

Review article

A review of the environmental safety of the PAT protein

Center for Environmental Risk Assessment, ILSI Research foundation

1156 Fifteenth Street N.W., Washington DC 20005-1743, USA

Keywords: PAT / *bar* / phosphinothricin / glufosinate / glufosinate ammonium / herbicide tolerant / genetically engineered / environmental risk assessment

INTRODUCTION

This document provides a comprehensive review of information and data relevant to the environmental risk assessment of the protein phosphinothricin-N-acetyl transferase (PAT) produced in genetically engineered (GE) plants by genes isolated from *Streptomyces viridochromogenes* (*pat* gene) or *Streptomyces hygroscopicus* (*bar* gene) and presents a summary statement about the environmental safety of this protein. All sources of information reviewed herein were publicly available and include: dossiers presented to regulatory authorities; decision summaries prepared by regulatory authorities; peer-reviewed literature; and product summaries prepared by product developers. Many GE plants contain the *pat* gene for use as a selectable marker during development. In those cases, there are one or more additional transgenes contained in the plant and the final product is not necessarily glufosinate tolerant. Although this document will not address these additional genes and phenotypes, their presence should be noted when looking at data on the GE plants that express PAT.

Environmental risk assessments related to the introduction of GE plants are conducted on a case-by-case basis taking into account the biology of the plant, the nature of the transgene and the protein or gene product it produces, the phenotype conferred by the transgene, as well as the intended use of the plant and the environment where it will be introduced (i.e. the receiving environment). These assessments typically involve comparisons of the transgenic event to an untransformed parent line and/or closely related isolate, and also use baseline knowledge of the relevant plant species (CBD, 2000b; Codex, 2003a, 2003b; EFSA, 2006a; NRC, 1989; OECD, 1992;

OECD, 2006). The objective of these comparisons is to identify potential risks that the GE plant might present beyond what is already accepted for similar plants in the environment by identifying meaningful differences between the GE crop and its conventional counterpart. Any identified differences that have the potential to cause relevant adverse effects can subsequently be evaluated for likelihood and consequence.

To date, regulatory authorities in 11 different countries have issued approvals for the environmental release of GE plants expressing the PAT protein, either by itself or in combination with other GE traits (see Tab. 1). This represents approximately 38 transformation events and includes 8 species of plant: *Beta vulgaris* L. (sugarbeet), *Brassica napus* L. and *Brassica rapa* L. (oilseed rape and turnip rape, respectively, although both can be referred to as canola), *Cichorium intybus* (chicory), *Glycine max* L. (soybean), *Gossypium hirsutum* L. (cotton), *Oryza sativa* L. (rice) and *Zea mays* L. (maize). These regulatory analyses have generally considered three categories of potential harm: (1) the PAT protein may have an adverse impact on non-target organisms; (2) transformation of the host plant and subsequent expression of the PAT protein may alter the characteristics of the plant, resulting in adverse environmental impacts (e.g. increased weediness); and (3) introgression of the gene encoding the PAT protein into a sexually compatible plant species may alter that species resulting in adverse environmental impacts (e.g. establishment of new weedy populations) (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1998, 1999a, 1999b,

Table 1. Regulatory approvals for the environmental release of GE plants containing PAT protein.

Species	Events/Crosses	Alternate Designations	Source of PAT Gene	United States	Canada	Japan	Australia	European Union	Brazil	Argentina	Colombia	Philippines	South Africa	Uruguay
<i>Beta Vulgaris</i> (sugarbeet)	ACS-BV001-3	T120-7	<i>Streptomyces viridochromogenes</i>	X	X									
<i>Brassica napus</i> (oilseed rape/canola)	HCN10		<i>Streptomyces viridochromogenes</i>	X	X	X								
	ACS-BN007-1	HCN92	<i>Streptomyces viridochromogenes</i>	X	X	X	X							
	ACS-BN004-7 x ACS-BN001-4	MS1, RF1 ; PGS1	<i>Streptomyces hygrosopicus</i>	X	X	X	X							
	ACS-BN004-7 x ACS-BN002-5	MS1, RF2 ; PGS2	<i>Streptomyces hygrosopicus</i>	X	X	X	X							
	ACS-BN005-8 x ACS-BN003-6	MS8 x RF3	<i>Streptomyces hygrosopicus</i>	X	X	X	X							
	PHY14, PHY35		<i>Streptomyces hygrosopicus</i>			X								
	PHY36		<i>Streptomyces hygrosopicus</i>			X								
	ACS-BN008-2	T45, HCN28	<i>Streptomyces viridochromogenes</i>	X	X	X	X							
<i>Brassica rapa</i> (bird rape/canola)	HCR-1		<i>Streptomyces viridochromogenes</i>		X									
<i>Cichorium intybus</i> (chicory)	RM3-3, RM3-4, RM3-6		<i>Streptomyces hygrosopicus</i>	X	X	X		X						
<i>Glycine max</i> (soybean)	ACS-GM005-3	A2704-12, A2704-21, A5547-35	<i>Streptomyces viridochromogenes</i>	X	X	X			X					
	ACS-GM006-4	A5547-127	<i>Streptomyces viridochromogenes</i>	X	X	X			X					
	ACS-GM003-1	GU262	<i>Streptomyces viridochromogenes</i>	X										
	ACS-GM001-8, ACS-GM002-9	W62, W98	<i>Streptomyces hygrosopicus</i>	X										
<i>Gossypium hirsutum</i> (cotton)	DAS-24236-5	281-24-236	<i>Streptomyces viridochromogenes</i>	X										
	DAS 21023-5	3006-210-23	<i>Streptomyces viridochromogenes</i>	X										
	DAS 21023-5 x DAS-24236-5		<i>Streptomyces viridochromogenes</i>	X										
	DAS 21023-5 x DAS-24236-5 x MON-01445-2		<i>Streptomyces viridochromogenes</i>	*										
	DAS 21023-5 x DAS-24236-5 x MON-88913-8		<i>Streptomyces viridochromogenes</i>	*										
	ACS-GH001-3	LLCotton25	<i>Streptomyces viridochromogenes</i>	X	X	X	X		X					
	ACS-GH001-3 x MON-15985-7	LLCotton25 x MON15985	<i>Streptomyces hygrosopicus</i>	*		X								
<i>Oryza sativa</i> (rice)	ACS-O5001-4, ACS-O5002-5	LLRice06, LLRice62	<i>Streptomyces hygrosopicus</i>	X										
	BCS-O5003-7	LLRice601	<i>Streptomyces hygrosopicus</i>	X										
<i>Zea mays</i> (maize/corn)	SYN-EV176-9	176	<i>Streptomyces hygrosopicus</i>	X	X	X	X							
	PH-000676-7, PH-000678-9, PH-000680-2	676, 678, 680	<i>Streptomyces viridochromogenes</i>	X	X	X								
	DKB-89790-5	B16, DLL25	<i>Streptomyces hygrosopicus</i>	X	X	X								

Table 1 (continued). Regulatory approvals for the environmental release of GE plants containing PAT protein.

Species	Events/Crosses	Alternate Designations	Source of PAT Gene	United States	Canada	Japan	Australia	European Union	Brazil	Argentina	Colombia	Philippines	South Africa	Uruguay
	SYN-BT011-1	BT11 (X4334CBR, X4734CBR)	<i>Streptomyces viridochromogenes</i>	X	X	X			X	X	X	X	X	X
	SYN-BT011-1 x MON-00021-9	BT11 x GA21	<i>Streptomyces viridochromogenes</i>	*	X	X			X					
	SYN-BT011-1 x SYN-IR162-4	BT11 x MIR162	<i>Streptomyces viridochromogenes</i>	X										
	SYN-BT011-1 x SYN-IR162-4 x SYN-IR604-5	BT11 x MIR162 x MIR604	<i>Streptomyces viridochromogenes</i>	X										
	SYN-BT011-1 x SYN-IR604-5	BT11 x MIR604	<i>Streptomyces viridochromogenes</i>	*	X									
	SYN-BT011-1 x SYN-IR604-5 x MON0021-9	BT11 x MIR604 x GA21	<i>Streptomyces viridochromogenes</i>	*										
	ACS-ZM004-3	CBH-351	<i>Streptomyces hygrosopicus</i>	X										
	DAS-06275-8		<i>Streptomyces hygrosopicus</i>	X	X	X								
	DAS-59122-7		<i>Streptomyces viridochromogenes</i>	X	X	X								
	DAS-59122-7, MON-00603-6	DAS-59122-7 x NK603	<i>Streptomyces viridochromogenes</i>	*	X	X								
	DAS-59122-7 x DAS-01507-1 x MON-00603-6	DAS-59122-7 x TC1507 x NK603	<i>Streptomyces viridochromogenes</i>	*	X	X								
	DKB-89614-9	DBT418	<i>Streptomyces hygrosopicus</i>	X	X	X				X				
	MON-89034-3 x DAS-01507-1 x MON-88017-3 x DAS-59122-7	MON89034 x TC1507 x MON88017 x DAS-59122-7	<i>Streptomyces viridochromogenes</i>	X	X	X								
	ACS-ZM001-9	MS3	<i>Streptomyces hygrosopicus</i>	X	X									
	ACS-ZM005-4	MS6	<i>Streptomyces hygrosopicus</i>	X										
	MON-00603-6 x ACS-ZM003-2	NK603 x T25	<i>Streptomyces viridochromogenes</i>	*		X								
	ACS-ZM002-1, ACS-ZM003-2	T14, T25	<i>Streptomyces viridochromogenes</i>	X	X	X		X	X	X				
	ACS-ZM003-2, MON-00810-6	T25 x MON810	<i>Streptomyces viridochromogenes</i>	*		X								
	DAS-01507-1	TC1507	<i>Streptomyces viridochromogenes</i>	X	X	X								
	DAS-01507-1, DAS-59122-7	TC1507 x DAS-59122-7	<i>Streptomyces viridochromogenes</i>	*	X	X								
	DAS-01507-1 x MON-00603-6	TC1507 x NK603	<i>Streptomyces viridochromogenes</i>	*	X	X								

x = Approved for environmental (commercial) release.

* These are stacked events that may be considered approved in the country indicated.

1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996a, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b).

Note that environmental effects that may be associated with the use of the herbicide glufosinate in association with GE plants producing PAT are outside the purview of this review.

ORIGIN AND FUNCTION OF PAT

Phosphinothricin, bialaphos, and glufosinate ammonium

In the early 1970's a previously unknown amino acid was isolated independently from two species of *Streptomyces* by laboratories working in Germany (from *Streptomyces viridochromogenes*) and Japan (from *Streptomyces hygroscopicus*) (Bayer et al., 1972; Kondo et al., 1973; OECD, 1999). Originally seen in a tripeptide with two alanine residues (see Fig. 1), the new amino acid (*L*-2-amino-4-[hydroxyl(methyl)phosphinyl] butyric acid) was given the name phosphinothricin (PT) and the tripeptide called phosphinothricin tripeptide (PTT) or bialaphos¹ (Bayer et al., 1972; Hoerlein, 1994; Kondo et al., 1973; OECD, 1999). In Germany, racemic mixtures were produced (*D,L*-phosphinothricin or *D,L*-PPT) and determined to have herbicidal activity. *D,L*-PPT-ammonium, referred to by the common name glufosinate ammonium (GLA) is the active ingredient in herbicide formulations marketed worldwide. In Japan, the bialaphos tripeptide was observed to have herbicidal activity and this has been commercialized as well (Hoerlein, 1994).




Figure 1. The structure of phosphinothricin, PTT and glutamic acid.

¹ Also sometimes “bilanafos” or “bilanaphos”.

Phosphinothricin inhibits the activity of the glutamine synthetase enzyme (GS) by competitively binding in place of the normal substrate, glutamate (glutamic acid). This prevents the synthesis of *L*-glutamine, which is not only an important chemical precursor for the synthesis of nucleic acids and proteins, but serves as the mechanism of ammonia (NH₃) incorporation for plants (Hoerlein, 1994; OECD, 1999, 2002). Treatment with phosphinothricin causes accumulation of ammonia and cessation of photosynthesis, probably due to the lack of glutamine (Hoerlein, 1994; OECD, 1999, 2002).

ISOLATION AND FUNCTION OF PHOSPHINOTHTRICIN ACETYL TRANSFERASE (PAT)

The identification of the GS inhibitor phosphinothricin from *Streptomyces* suggested that these bacteria employ a biochemical mechanism to preserve endogenous GS activity. In the late 1980s, two genes were identified independently based on their ability to confer resistance to phosphinothricin inhibition of GS, both of which encode a phosphinothricin acetyl transferase protein (PAT). The bialaphos resistance gene, *bar*, was isolated from *S. hygroscopicus* while the homologous gene from *S. viridochromogenes* was termed *pat* after the function of the enzyme (OECD, 1999a; Thompson et al., 1987; Wohlleben et al., 1998). Both proteins have been used extensively in genetic engineering of crop plants. They both consist of 183 amino acids, with a sequence identity of 85% (OECD, 1999a; Wehrmann et al., 1996; Wohlleben et al., 1998). Importantly, both proteins acetylate phosphinothricin but show no activity with glutamate, which is structurally similar, or with any other amino acids tested, indicating a high specificity (OECD, 1999a; Thompson et al., 1987; Wehrmann et al., 1996). The only recorded differences in activity between the two proteins are minor differences in the optimal pH, and a significantly different affinity for acetyl-coA (a co-substrate); these differences are not expected to be meaningful *in planta* (OECD, 1999a; Wehrmann et al., 1996). Because the PAT proteins encoded by *bar* and *pat* are structurally and functionally equivalent, with similar molecular weights, immuno-cross-reactivity, substrate affinity and specificity, they are considered together in this document and will both be referred to as PAT protein.

The PAT enzyme acetylates phosphinothricin at the N-terminus. N-acetyl phosphinothricin has no herbicidal activity, and resistance is therefore conferred through modification of the herbicide rather than the target of its activity (OECD, 1999a; Thompson et al., 1987; Wehrmann et al., 1996; Wohlleben et al., 1998).

EXPRESSION OF PAT IN PHOSPHINOTHRICIN-TOLERANT GE PLANTS

Data for the level of expression of PAT in phosphinothricin-tolerant GE plants that have obtained regulatory approvals are available in publicly accessible regulatory documents (ANZFA, 2000, 2001a, 2001b, 2002; CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2004, 2005, 2006a, 2006b; EC, 1996, 1997, 1998, 2001; EFSA, 2005, 2006, 2008a, 2008b, 2009a, 2009b; FSANZ, 2003, 2004a, 2004b, 2005a, 2005b, 2008; Japan BCH, 1996a, 1996b, 1996c; Health Canada, 2006a, 2006b; Japan BCH, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2003, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2003, 2006; Philippines, 2005; USDA APHIS, 1994a, 1994b, 1995b, 1995d, 1995e, 1996a, 1996d, 1997a, 1997b, 1997d, 1997e, 1998b, 1998d, 1998f, 1998h, 1998k, 1998l, 2000, 2001a, 2002a, 2002c, 2003a, 2003b, 2004b, 2004d, 2006a; USEPA, 2001, 2005, 2009a, 2009b). Tissue types tested and sampling methodologies vary greatly. The most common method uses enzyme-linked immunosorbent assay (ELISA) to quantify the amount of protein present in a given sample, but other methods include an assay for enzymatic activity and the use of Northern blots to quantify mRNA. Normally, one or more samples are collected from plants in field trials or greenhouse experiments and the amount of protein is given as a mean accompanied by either a standard deviation or a range of observed values to show variability. The result is often quantified as a ratio to the dry weight of the sample (e.g. $\mu\text{g PAT/g dry weight}$), but some reports calculate the ratio to the fresh weight of the sample or to the total extractable protein from the sample (e.g. $\mu\text{g PAT/g total protein}$).

Variations in methodology for both sample collection and subsequent analysis make direct statistical comparisons of the data inappropriate. However, the weight of evidence suggests PAT protein is expressed at low levels (see Annex I and associated references). The highest reported levels of expression observed in each species using ELISA are reported in Table 2.

ESTABLISHMENT AND PERSISTENCE OF PAT-EXPRESSING PLANTS IN THE ENVIRONMENT

Familiarity with the biology of the non-transformed or host plant species in the receiving environment is typically the starting point for a comparative environmental risk assessment of a GE plant (CBD, 2000b; Codex, 2003a, 2003b; EFSA, 2006a; NRC, 1989; OECD, 1992, 2006).

Table 2. Highest reported expression level of PAT protein using ELISA¹.

Species	Event GE Plant	Expression	Tissue	Reference
<i>Beta vulgaris</i>	T120-7	966 ng/g	Top ²	USDA APHIS, 1998b
<i>Brassica napus</i>	Topas19/2 x T45	944 ng/g	Leaf	USDA APHIS, 2002a
<i>Glycine max</i>	A5547-127	20202 \pm 359 ³ ng/g	Seed	Japan BCH, 2005f
<i>Gossypium hirsutum</i>	LLCOTTON25	127000 \pm 18000 ³ ng/g ⁴	Cleaned Seed	USDA APHIS, 2002c
<i>Oryza sativa</i>	LLRICE62	84700 ng/g ⁴	Leaf	CFIA, 2006b
<i>Zea mays</i>	DAS-06275-8	935000 ng/g ^{4,5}	Leaf	USDA APHIS, 2004b

¹ These values are not cross-comparable due to differences in sample collection and preparation methodology.

² Top refers to all above-ground tissue (i.e. leaves and stems).

³ Reported as mean \pm standard deviation.

⁴ Reported as ng/g fresh weight.

⁵ Represents the highest value in a reported range.

Information about the biology of the host plant can be used to identify species-specific characteristics that may be affected by the novel trait so as to permit the transgenic plant to become “weedy,” invasive of natural habitats, or to be otherwise harmful to the environment. It can also provide details on significant interactions between the plant and other organisms that may be important when considering potential harms. By considering the biology of the host plant, a risk assessor can identify potential hazards that may be associated with the expression of the novel protein (e.g. PAT) and then be able to assess the likelihood of these hazards being realized. For example, if the plant species is highly domesticated and requires significant human intervention to grow or reproduce, the assessor can take that into account when assessing the likelihood of the GE plant establishing outside of cultivation.

PHENOTYPIC DATA

In order to determine if GE plants expressing PAT are phenotypically different than their non-transformed counterparts, a variety of data have been collected and are presented with varying degree of detail in regulatory submissions related to the environmental release of these plants (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b,

2004, 2005, 2006a, 2006b; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c; Health Canada, 2006a, 2006b; Japan BCH, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2003, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2003, 2006; Philippines, 2005; USDA APHIS, 1994a, 1994b, 1995b, 1995d, 1995e, 1996a, 1996d, 1997a, 1997b, 1997d, 1997e, 1998b, 1998d, 1998f, 1998h, 1998k, 1998l, 2000, 2001a, 2002a, 2002c, 2003a, 2003b, 2004b, 2004d, 2006a; USEPA, 2001, 2005, 2009a, 2009b). The data that are reported are dependent on the species of plant, but in general include information on gross morphology (*e.g.* height, number of leaves, number of branches or nodes, etc.), reproductive characteristics including seed production, survival and germination, as well as seedling vigor, overwintering ability, susceptibility to disease and pest pressure, and frequently the potential to volunteer following harvest. Phenotypic analyses may also include agronomic characteristics such as yield and performance in the field (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2004, 2005, 2006a, 2006b; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c; Health Canada, 2006a, 2006b; Japan BCH, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2003, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2003, 2006; Philippines, 2005; USDA APHIS, 1994a, 1994b, 1995b, 1995d, 1995e, 1996a, 1996d, 1997a, 1997b, 1997d, 1997e, 1998b, 1998d, 1998f, 1998h, 1998k, 1998l, 2000, 2001a, 2002a, 2002c, 2003a, 2003b, 2004b, 2004d, 2006a; USEPA, 2001, 2005, 2009a, 2009b). Frequently, statistically significant differences in a handful of phenotypic characteristics are reported between GE plants and controls in a given experiment (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2004, 2005, 2006a, 2006b; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c; Health Canada, 2006a, 2006b; Philippines, 2005; USDA APHIS, 1994a, 1994b, 1995b, 1995d, 1995e, 1996a, 1996d, 1997a, 1997b, 1997d, 1997e, 1998b, 1998d, 1998f, 1998h, 1998k, 1998l, 2000, 2001a, 2002a, 2002c, 2003a, 2003b, 2004b, 2004d, 2006a; USEPA, 2001, 2005, 2009a, 2009b). However, these differences usually are not repeated in multiple experiments and regulatory decisions have concluded that any such differences are likely not due to the expression of the PAT protein and do not represent meaningful differences with respect to the potential for adverse impact to the environment (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006a; EC, 1996, 1997,

1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b).

WEEDINESS IN AGRICULTURAL ENVIRONMENTS

All of the plant species that have been engineered to express PAT have some potential to “volunteer” as weeds in subsequent growing seasons and demonstrate varying degrees of ability to persist in an agricultural environment (OECD, 1997, 2000, 2001, 2003a, 2008; OGTR, 2008). The characteristics that influence the ability of a plant to volunteer are largely the same as those for weediness in general, such as seed dormancy, shattering, and competitiveness (Baker, 1974). The data available indicate there is no linkage between PAT protein expression and any increased survival or over-wintering capacity that would alter the prevalence of volunteer plants in the subsequent growing season (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1998, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b). Following-season volunteers expressing PAT may complicate volunteer management programs, particularly if different crop species expressing glufosinate tolerance are planted in consecutive rotations. Alternative options are available for managing glufosinate-tolerant volunteers, including the use of other herbicides and mechanical weed control (Beckie et al., 2004; OECD, 1997, 2000, 2001, 2003a, 2008; OGTR, 2008).

WEEDINESS IN NON-AGRICULTURAL ENVIRONMENTS

The primary mechanisms by which PAT may be introduced into a non-agricultural environment are: (1) seed or propagule movement (which may include incidental release during transportation of commodities)

and establishment of the GE plant outside of cultivated areas, and; (2) gene flow from the GE plant to a naturalized (or feral) population of the same crop species or other sexually compatible relatives (Mallory-Smith and Zapiola, 2008). Risk assessments for GE plants expressing PAT have considered the potential impacts associated with both types of introduction (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1998, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b).

While all plants can be considered weeds in certain contexts, none of the crops for which glufosinate-tolerant GE lines are available are considered to be invasive or problematic weeds outside of agricultural systems. Most can persist under favorable conditions and they may at times require management, particularly when they volunteer in subsequent crops (OECD, 1997, 2000, 2001, 2003a, 2008; OGTR, 2008; USDA APHIS, 2004d). Based on agronomic and compositional data showing that PAT does not have a significant impact on agronomic or compositional traits (including those that are related to weediness), the evidence to date shows that expression of the PAT protein has not resulted in any altered potential for weediness for those GE plant events subjected to environmental risk assessment. PAT expression only affects the ability of the plant to survive if treated with glufosinate. Just as in agricultural environments, other management options to control glufosinate-tolerant plants in non-agricultural environments are available (Beckie et al., 2004; OECD, 1997, 2000, 2001, 2003a, 2008; OGTR, 2008).

MOVEMENT OF THE TRANSGENE TO WILD RELATIVES

The movement of transgenes to wild relatives is pollen-mediated and the production of reproductively viable hybrids depends on the physical proximity and flowering synchrony of the GE plants to sexually compatible species. The evidence shows that expression of the PAT protein in a range of plant species has not resulted in any

alteration to anticipated gene flow. However introgression of glufosinate tolerance into sexually compatible, weedy populations in agricultural or peri-agricultural ecosystems is possible and has the potential to raise management issues (Mallory-Smith and Zapiola, 2008; Warwick et al., 2007). In at least one instance, a regulatory decision has geographically limited the release of a herbicide-tolerant GE plant: the environmental approval of *B. rapa* event ZSR500/502 (glyphosate resistance) was limited to the western region of Canada due to the presence of feral populations of *B. rapa* in eastern Canada where it is considered a weed of agriculture (CFIA, 1998d). However, no such decisions have been made for plants expressing PAT that are glufosinate-resistant, and all of the publicly available regulatory decisions conclude that the movement of the *pat* gene to wild relatives is not a substantial risk for any of the GE plants that have been considered (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006a; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b).

ADVERSE IMPACTS ON OTHER ORGANISMS IN THE RECEIVING ENVIRONMENT

The potential for PAT protein expression in GE plants to have an adverse impact on organisms has been considered in regulatory risk assessments using a weight-of-evidence approach (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006a; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b). These risk assessments have generally considered the potential for the novel protein to be toxic to other organisms, as well as the history of prior

environmental exposure to the protein. Toxic proteins are known to act acutely (Sjoblad et al., 1992). Acute, intravenous toxicity experiments in mice show the PAT protein has no toxicity even at doses much higher than would be encountered due to environmental exposure to GE plants expressing the PAT protein (Herouet et al., 2005). In addition, the PAT protein shows no homology to known toxins or allergens and is rapidly digested in experiments simulating gastric environment (Herouet et al., 2005). The *Streptomyces* bacteria which are the source of PAT proteins are widespread in environments around the world, and additional species of *Streptomyces* are known to possess similar enzymatic activity, indicating that PAT protein homologs are likely ubiquitous in the environment and regulatory decisions have concluded that exposure to PAT proteins from GE plants does not represent a potential for adverse impacts on other organisms (Herouet et al., 2005; CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006a; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b).

Risk assessors have also considered whether the introduction of PAT proteins into GE plants might lead to changes in the plant that would adversely impact other organisms. Phenotypic characterization (see above) as well as compositional analyses (see below) and nutritional analyses show that the introduction of PAT proteins has not had any unanticipated effects on characteristics of GE plants that might impact other organisms (ANZFA, 2000, 2001a, 2001b, 2002; CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2004, 2005, 2006a, 2006b; EC, 1996, 1997, 1998, 2001; EFSA, 2005, 2006, 2008a, 2008b, 2009a, 2009b; FSANZ, 2003, 2004a, 2004b, 2005a, 2005b, 2008; Japan BCH, 1996a, 1996b, 1996c; Health Canada, 2006a, 2006b; Japan BCH, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2003, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010, OGTR, 2003, 2006; Philippines, 2005; USDA APHIS, 1994a, 1994b, 1995b, 1995d, 1995e, 1996a, 1996d, 1997a, 1997b, 1997d, 1997e, 1998b, 1998d, 1998f, 1998h, 1998k, 1998l, 2000,

2001a, 2002a, 2002c, 2003a, 2003b, 2004b, 2004d, 2006a; USEPA, 2001, 2005, 2009a, 2009b). Based on experimental evidence that PAT proteins are not toxic and the observation that exposure to PAT is widespread in the environment, regulatory authorities have concluded that the expression of PAT in GE plants does not have any meaningful potential to adversely impact other organisms (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006a; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b).

COMPOSITIONAL ANALYSIS OF PAT-EXPRESSING PLANTS

Detailed compositional analysis is a scientifically rigorous component of the characterization of GE plants and is a regulatory requirement for GE food and feed safety approvals (OECD, 1992; WHO, 1995; FAO/WHO, 1996; EFSA, 2006a; Codex, 2003a, 2003b). The choice of analyses conducted depends on the nature of the product and its intended uses. Although compositional analysis is not typically required for environmental risk assessments, it is often considered in the context of demonstrating whether or not there have been unanticipated changes to the GE plant (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006a; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b). GE plants expressing PAT have undergone a variety of compositional analyses, including for proximate (protein, fat, amino acid, fiber, ash) as well as for nutritional components and known toxicants or antinutrients (such as gossypol in cotton or glucosinolates in canola) (ANZFA, 2000, 2001a, 2001b, 2002; CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d,

1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2004, 2005, 2006a, 2006b; EC, 1996, 1997, 1998, 2001; EFSA, 2005, 2006, 2008a, 2008b, 2009a, 2009b; FSANZ, 2003, 2004a, 2004b, 2005a, 2005b, 2008; OGTR, 2003, 2006; Philippines, 2005; USDA APHIS, 1994a, 1994b, 1995b, 1995d, 1995e, 1996a, 1996d, 1997a, 1997b, 1997d, 1997e, 1998b, 1998d, 1998f, 1998h, 1998k, 1998l, 2000, 2001a, 2002a, 2002c, 2003a, 2003b, 2004b, 2004d, 2006a; USEPA, 2001, 2005, 2009a, 2009b). Although statistically significant differences between the composition of GE plants and their non-transformed counterparts have been reported, these differences have not been attributed to expression of the PAT protein, and subsequent regulatory decisions have concluded that the composition of GE plants expressing PAT is not meaningfully different with respect to potential impact on the environment (CFIA, 1995a, 1995b, 1996a, 1996b, 1996c, 1996d, 1996e, 1996f, 1998a, 1998b, 1998c, 1999, 2002a, 2002b, 2005, 2006a; EC, 1996, 1997, 1998, 2001; Japan BCH, 1996a, 1996b, 1996c, 1997a, 1997b, 1997c, 1997d, 1997e, 1999a, 1999b, 1999c, 1999d, 2002, 2005, 2006a, 2006b, 2006c, 2006d, 2006e, 2006f, 2007a, 2007b, 2008, 2009, 2010; OGTR, 2002, 2003, 2006; Philippines, 2005; USDA APHIS, 1995a, 1995c, 1995f, 1996b, 1996c, 1996e, 1997c, 1997f, 1998a, 1998c, 1998e, 1998g, 1998i, 1998j, 1999a, 1999b, 1999c, 2001b, 2001c, 2002b, 2003c, 2004a, 2004c, 2005, 2006b; USEPA, 2001, 2005, 2009a, 2009b).

CONCLUSION

The PAT protein expressed in GE plants is encoded by one of the homologous genes *pat* or *bar*, isolated from the related bacteria *Streptomyces viridochromogenes* or *Streptomyces hygroscopicus*, respectively. Environmental release approvals have been granted for 8 species of plants expressing PAT proteins in 11 different countries including at least 38 separate transformation events. Data from regulatory submissions and peer-reviewed literature show that the PAT protein expressed in GE plants has negligible impact on the phenotype of those plants, beyond conferring tolerance to the herbicide glufosinate. Risk assessments associated with regulatory review of these plants for use in the environment show that expression of PAT does not alter the potential for persistence or spread of GE plants in the environment, does not alter the reproductive biology or potential for gene flow, and does not increase the risks for adverse effects to other organisms. Although the introduction of PAT to GE plants has the potential to complicate the management of herbicide-tolerant volunteers or weedy

relatives in agriculture, the evidence does not indicate that expression of PAT has impacted the effectiveness or availability of alternative control measures such as other herbicides or mechanical weed control. Taken together, these regulatory analyses support the conclusion that, for the species and environments that have been considered to date, the expression of the PAT protein in GE plants does not present any meaningful risk to the environment.

REFERENCES

Journal articles and books

- Baker H.G. (1974). The evolution of weeds. *Annual Review of ecology and Systematics* 5:1–24.
- Beckie H.J., Seguin-Swartz G., Nair H., Warwick S.I., and Johnson E. (2004). Multiple herbicide-resistant canola can be controlled by alternative herbicides. *Weed Science* 52 (1):152–157.
- De Block M., Botterman J., Vandewiele M., Dockx J., Thoen C., Gossele V., Movva N.R., Thompson C., van Montagu M., and Leemans J. (1987). Engineering herbicide resistance in plants by expression of a detoxifying enzyme. *The EMBO Journal* 6(9):2513–2518.
- Droge-Laser W., Siemeling U., Puhler A., and Broer I. (1994). The metabolites of the herbicide L-Phosphinothricin (Glufosinate): Identification, stability and mobility in transgenic, herbicide-resistant and untransformed plants. *Plant Physiology* 105:159–166.
- Herouet C., Esdaile D. J., Mallyon B. A., Debruyne E., Schulz A., Currier T., Hendrickx K., van der Klis R.-J., and Rouan D. (2005). Safety evaluation of the phosphinothricin acetyltransferase proteins encoded by the *pat* and *bar* sequences that confer tolerance to glufosinate-ammonium herbicide in transgenic plants. *Regulatory Toxicology and Pharmacology* 41:134–149.
- Hoerlein G. (1994). Glufosinate (Phosphinothricin), a Natural Amino Acid with Unexpected Herbicidal Properties. *Reviews of Environmental Contamination and Toxicology* 138:73–145.
- Mallory-Smith C. and Zapiola M. (2008). Gene flow from glyphosate-resistant crops. *Pest Management Science* 64:428–440.
- Schwarz D., Berger S., Heinzmann E., Muschko K., Welzel K., and Wohlleben W. (2004). Biosynthetic gene cluster of the herbicide phosphinothricin tripeptide from *Streptomyces viridochromogenes* Tu494. *Applied and Environmental Microbiology* 70(12):7093–7102.
- Sjoblad R.D., McClintock J.T. and Engler R. (1992). Toxicological considerations for protein components of biological pesticide products. *Regulatory Toxicology and Pharmacology* 15: 3–9.
- Thompson C. J., Movva N. R., Tizard R., Crameri R., Davies J. E., Lauwereys M., and Botterman J. (1987). Characterization

of the herbicide-resistance gene *bar* from *Streptomyces hygroscopicus*. The EMBO Journal 6(9):2519–2523.

Warwick, S.I., Legere, A., Simard, M.-J., and James, T. (2008). Do escaped transgenes persist in nature? The case of an herbicide resistance transgene in a weedy *Brassica rapa* population. Molecular ecology. 17:1387–1395.

Wehrmann A., van Vliet A., Opsomer C., Botterman J., and Schulz A. (1996). The similarities of *bar* and *pat* gene products make them equally applicable for plant engineers. Nature Biotechnology 14:1274–1278.

Wohlleben W., Arnold W., Broer I. Hillermann D., Strauch E., and Puhler A. (1988). Nucleotide sequence of the phosphinothricin *N*-acetyltransferase gene from *Streptomyces viridochromogenes* Tu494 and its expression in *Nicotiana tabacum*. Gene 70:25–37.

Regulatory citations

- ANZFA (2000). Draft Risk Analysis Report: Food derived from insect-protected Bt-176 corn. Australia New Zealand Food Authority, Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/A385%20FA.pdf
- ANZFA (2001a). Draft Risk Assessment Report: Food derived from insect-protected, herbicide-tolerant BT11 corn. Australia New Zealand Food Authority, Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/A386_IR.pdf
- ANZFA (2001b). Risk assessment for application A375: food derived from glufosinate tolerant maize line T25. Australia New Zealand Food Authority, Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/A375%20Final%20AR.pdf
- ANZFA (2002). Final assessment report (inquiry - section 17) application A380 food from insect-protected and glufosinate ammonium-tolerant DBT418 corn. Australia New Zealand Food Authority, Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/A380_Final_Assessment_Report.pdf
- CFIA (1995a). Decision document DD95-01: Determination of environmental safety of Agrevo Canada Inc.'s glufosinate ammonium-tolerant canola. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9501e.shtml>
- CFIA (1995b). Decision document DD95-04: Determination of environmental safety of Plant Genetic systems Inc. (PGS) novel hybridization system for canola (*Brassica napus* L.) (MS1, RF1). Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9504e.shtml>
- CFIA (1996a). Decision document DD96-09: Determination of environmental safety of event 176 Bt corn (*Zea mays* L.) developed by Ciba Seeds and Mycogen Corporation. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9609e.shtml>
- CFIA (1996b). Decision document DD96-11: Determination of environmental safety of Agrevo Canada Inc.'s glufosinate ammonium-tolerant canola line HCN28(T45). Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9611e.shtml>
- CFIA (1996c). Decision Document DD96-12: Determination of environmental safety of Northrup King Seeds' european corn borer (ecB) resistant corn (*Zea mays* L.). Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9612e.shtml>
- CFIA (1996d). Decision document 96–15: CFIA (1996c). Determination of environmental safety of Dekalb Canada Inc.'s glufosinate ammonium-tolerant corn line DLL25. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9615e.shtml>
- CFIA (1996e). Decision document 96–16: Determination of environmental safety of Plant Genetic Systems Inc.'s (PGS) male sterile corn (*Zea mays* L.) line MS3. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9616e.shtml>
- CFIA (1996f). Decision document 96–17: Determination of environmental safety of Plant Genetic Systems Inc.'s (PGS) novel hybridization system for rapeseed (*Brassica napus* L.) (MS8xRF3). Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9617e.shtml>
- CFIA (1998a). Decision Document 98–22: Determination of the safety of AgrEvo Canada Inc.'s glufosinate ammonium tolerant corn (*Zea mays*) lines, T14 and T25. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9822e.shtml>
- CFIA (1998b). Decision Document DD98-23: Decision document 98–23: determination of environmental safety of Dekalb Genetics Corporation's European corn borer (ECB) resistant corn (*Zea mays* L.) line DBT418. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9823e.shtml>
- CFIA (1998c). Decision Document DD98-28: Determination of the safety of AgrEvo Canada Inc.'s glufosinate ammonium herbicide-tolerant *Brassica rapa* canola line HCR-1. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9828e.shtml>
- CFIA (1998d). Determination of the safety of Monsanto Canada Inc.'s roundup herbicide-tolerant *Brassica rapa* canola lines ZSR500, ZSR502, and ZSR503. Canadian Food Inspection Agency (CFIA), Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9821e.shtml>
- CFIA (1999). Decision Document DD99-30: Determination of the environmental safety of AgrEvo Canada Inc.'s glufosinate ammonium tolerant soybean (*Glycine max*) A2704-12. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd9930e.shtml>
- CFIA (2002a). Decision document DD2002-39: Determination of the safety of Aventis CropsScience Canada Inc.'s glufosinate ammonium-tolerant sugar beet (*Beta vulgaris*) lines derived from event T120-7. Canadian Food Inspection Agency (CFIA)

- Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd0239e.shtml>
- CFIA (2002b) Decision Document DD2002-41: Determination of the Safety of Dow AgroSciences Canada Inc. and Pioneer Hi-Bred International's Insect Resistant and Glufosinate – Ammonium Tolerant Corn (*Zea mays* L.) Line 1507. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd0241e.shtml>
- CFIA (2004). Decision Document DD2004-49: Determination of the Safety of Bayer CropScience's Herbicide Tolerant LibertyLink® Cotton Event LLCotton25 (*Gossypium hirsutum* L.). Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd0449e.shtml>
- CFIA (2005). Decision Document DD2005-55: Determination of the safety of Dow AgroSciences Canada Inc. and Pioneer Hi-Bred Production Inc.'s insect resistant and glufosinate-ammonium herbicide tolerant corn (*Zea mays* L.) Line 59122. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd0555e.shtml>
- CFIA (2006a). Decision document DD2006-59: Determination of the safety of Dow AgroSciences Canada Inc.'s insect resistant and glufosinate – ammonium tolerant corn (*Zea mays* L.) Event DAS-06275-8. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd0659e.shtml>
- CFIA (2006b). Decision Document DD2006-58: Determination of the safety of Bayer CropScience's glufosinate ammonium tolerant rice (*Oryza sativa*) Event LLrice62. Canadian Food Inspection Agency (CFIA) Ottawa, Canada. <http://www.inspection.gc.ca/english/plaveg/bio/dd/dd0658e.shtml>
- European Commission (1996). 96/424/EC: Commission Decision of 20 May 1996 concerning the placing on the market of genetically modified male sterile chicory (*Cichorium intybus* L.) with partial tolerance to the herbicide glufosinate ammonium pursuant to Council Directive 90/220/EEC. Official journal of the European Union NO. L 175, 13/07/1996 P. 0025 – 0026. http://www.biosafety.be/GB/Dir.Eur.GB/Market/96_424/96_424.html
- European Commission (1997). 97/98/EC: Commission Decision of 23 January 1997 concerning the placing on the market of genetically modified maize (*Zea mays* L.) with the combined modification for insecticidal properties conferred by the Bt-endotoxin gene and increased tolerance to the herbicide glufosinate ammonium pursuant to Council Directive 90/220/EEC. Official journal of the European Union NO. L 031, 01/02/1997 P. 0069 – 0070. http://www.biosafety.be/PDF/97_98.pdf
- European Commission (1998). 98/293/EC: Commission Decision of 22 April 1998 concerning the placing on the market of genetically modified maize (*Zea mays* L. T25), pursuant to Council Directive 90/220/EEC (Text with EEA relevance). Official Journal L 131, 05/05/1998 p. 0030 – 0031. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:1998:131:0030:0031:EN:PDF>
- European Commission (2001). Opinion of the scientific committee on plants regarding “submission for placing on the market of glufosinate tolerant maize (*Zea mays*) transformation event T25” by the AgrEvo company now Aventis CropScience (Notification C/F/95/12/07). European Commission, Brussels, Belgium. http://ec.europa.eu/food/fs/sc/scp/out04_en.html
- EFSA (2005) For citation purposes: Opinion of the Scientific Panel on Genetically Modified Organisms on an application (reference EFSA,-GMONL-2004-02) for the placing on the market of insect-tolerant genetically modified maize 1507, for food use, under Regulation (EC) No 1829/2003 from Pioneer Hi-Bred International/Mycogen Seeds, The EFSA Journal (2005) 182, 1–22. http://www.europabio.org/InfoOperators/EFSA,%20Opinion1507_190105.pdf
- EFSA (2006). Opinion of the Scientific Panel on Genetically Modified Organisms on an application (Reference EFSA,-GMO-NL-2005-13) for the placing on the market of glufosinate-tolerant genetically modified LLCotton25, for food and feed uses, and import and processing under Regulation (EC) No 1829/2003 from Bayer CropScience, The EFSA Journal (2006) 429, 1–19. <http://www.efsa.europa.eu/en/scdocs/doc/429.pdf>
- EFSA (2008a). Opinion of the Scientific Panel on Genetically Modified Organisms on an application (Reference EFSA-GMO-UK-2005-25) for the placing on the market of glufosinate-tolerant oilseed rape T45 for food and feed uses, import and processing and renewal of the authorization of oilseed rape T45 as existing products, both under Regulation (EC) 1829/2003 from Bayer CropScience, The EFSA Journal (2008) 635, 1–22. <http://www.efsa.europa.eu/en/scdocs/doc/635.pdf>
- EFSA (2008b) Scientific Opinion of the Panel on Genetically Modified Organisms on a request from Pioneer Hi-Bred International on the authorisation for the placing on the market of the insect-resistant and herbicide-tolerant genetically modified maize 59122 x NK603, for food and feed uses, and import and processing under Regulation (EC) No 1829/2003. The EFSA Journal (2008) 874, 1–34. <http://www.efsa.europa.eu/en/scdocs/doc/874.pdf>
- EFSA (2009a). Scientific Opinion of the Panel on Genetically Modified Organisms on an application (Reference EFSA-GMO-UK-2005-21) for the placing on the market of insect-resistant and herbicide-tolerant genetically modified maize 59122 x 1507 x NK603 for food and feed uses, and import and processing under Regulation (EC) No 1829/2003 from Pioneer Hi-Bred International, Inc. The EFSA Journal (2009) 1050, 1–32. <http://cera-gmc.org/docs/decdocs/09-235-005.pdf>
- EFSA (2009b). Opinion of the Scientific Panel on Genetically Modified Organisms on an application (Reference EFSA-GMO-NL-2005-15) for the placing on the market of the insect-resistant and herbicide-tolerant genetically modified maize 1507 x 59122, for food and feed uses, import and processing under Regulation (EC) No 1829/2003 from Mycogen Seeds, c/o Dow AgroSciences LLC and Pioneer Hi-Bred International, Inc. as represented by Pioneer

- Overseas Corporation, The EFSA Journal 1074, 1–28. <http://www.efsa.europa.eu/en/scdocs/scdoc/1074.htm>
- FSANZ (2003). Final assessment report, application A446: food derived from insect-protected and glufosinate ammonium-tolerant corn line 1507. Food Standards Australia New Zealand (FSANZ), Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/ACF18.pdf
- FSANZ (2004a). Final Assessment Report: Application A481- Food derived from glufosinate ammonium tolerant soybean lines A2704-12 and 15547-127. Food Standards Australia New Zealand (FSANZ), Canberra, Australia. http://www.foodstandards.govt.nz/_srcfiles/A481_GM_soy_FAR.pdf
- FSANZ (2004b). Final Assessment Report: Application A518 – Food derived from insect-protected herbicide tolerant cotton line MXB-13. Food Standards Australia New Zealand (FSANZ), Canberra, Australia. http://www.foodstandards.govt.nz/_srcfiles/A518_GM_Cotton_FAR_FINAL.doc
- FSANZ (2005a). Final Assessment Report: Application A533 – Food Derived from Glufosinate Ammonium-Tolerant Cotton Line LL25. Food Standards Australia New Zealand (FSANZ), Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/FAR_A533%20GM%20cotton.pdf
- FSANZ (2005b). Final Assessment Report: Application A543 – Food Derived from Insect-Protected Glufosinate Ammonium-Tolerant Corn Line 59122-7. Food Standards Australia New Zealand (FSANZ), Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/A543_GM_Corn_DAR2.pdf
- FSANZ (2008). Final Assessment Report: Application A589 – Food derived from glufosinate ammonium-tolerant rice LLRICE62. Food Standards Australia New Zealand (FSANZ), Canberra, Australia. http://www.foodstandards.gov.au/_srcfiles/A589_GM_LLRIICE62_FAR.pdf
- Health Canada (2006a). Novel Food Information; Glufosinate Tolerant Rice Event LLRICE62. Health Canada, Ottawa, Canada. <http://www.hc-sc.gc.ca/fn-an/gmf-agm/appro/nf-an90decdoc-eng.php>
- Health Canada (2006b). Novel Food Information: *(mo)Cry1F* Insect Resistant, Glufosinate Tolerant Corn Event TC6275. Health Canada, Ottawa, Canada. <http://www.hc-sc.gc.ca/fn-an/gmf-agm/appro/nf-an111decdoc-eng.php>
- Japanese Biosafety Clearing House (1996a). Outline of the biological diversity risk assessment report: Type 1 use approval for Topas 19/2 (HCN92). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1996b). Outline of the biological diversity risk assessment report: Type 1 use approval for ACS-BN004-7×ACS-BN001-4 (Ms1, RF1). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1996c). Outline of the biological diversity risk assessment report: Type 1 use approval for lepidopteran resistant, herbicide tolerant maize SYN-BT011-1. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1997a). Outline of the biological diversity risk assessment report: Type 1 use approval for ACS-BN004-7×ACS-BN002-5 (MS1, RF2). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1997b). Outline of the biological diversity risk assessment report: Type 1 use approval for ACS-BN008-2 (HCN28). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1997c). Outline of the biological diversity risk assessment report: Type 1 use approval for T14. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1997d). Outline of the biological diversity risk assessment report: Type 1 use approval for T25. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1997e). Outline of the biological diversity risk assessment report: Type 1 use approval for SYN-EV176-9. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1999a). Outline of the biological diversity risk assessment report: Type 1 use approval for ACS-BN005-8×ACS-BN003-6 (MS8x RF3). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1999b). Outline of the biological diversity risk assessment report: Type 1 use approval for soybean A2704-12. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1999c). Outline of the biological diversity risk assessment report: Type 1 use approval for DLL25. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (1999d). Outline of the biological diversity risk assessment report: Type 1 use approval for maize DBT418 (DKB-89614-9). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2002). Outline of the biological diversity risk assessment report: Type 1 use approval for cotton DAS-01507-1. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2003). Outline of the biological diversity risk assessment report: Type 1 use approval for T25 x MON810 maize. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2005). Outline of the biological diversity risk assessment report: Type 1 use approval for DAS-01507-1 x MON-00603-6. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2006a). Outline of the biological diversity risk assessment report: Type 1 use approval for cotton LLCotton25 x MON15985 (ACS-GH001-3 x MON-15985-7). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2006b). Outline of the biological diversity risk assessment report: Type 1 use approval for DAS-59122-7. Japan Biosafety Clearing House (BCH). Tokyo, Japan.

- Japanese Biosafety Clearing House (2006c). Outline of the biological diversity risk assessment report: Type 1 use approval for DAS-59122-7 x NK603. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2006d). Outline of the biological diversity risk assessment report: Type 1 use approval for DAS-59122-7 x TC1507 x NK603. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2006e). Outline of the biological diversity risk assessment report: Type 1 use approval for TC1507 x DAS-59122-7. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2006f). Outline of the biological diversity risk assessment report: Type 1 use approval for soybean A5547-127. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2007a). Outline of the biological diversity risk assessment report: Type 1 use approval for soybean A5547-127. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2007b). Outline of the biological diversity risk assessment report: Type 1 use approval for SYN-BT011-1 x MON-00021-9. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2008). Outline of the biological diversity risk assessment report: Type 1 use approval for TC6275, (DAS-06275-8). Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2009). Outline of the biological diversity risk assessment report: Type 1 use approval for SmartStax maize. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- Japanese Biosafety Clearing House (2010). Outline of the biological diversity risk assessment report: Type 1 use approval for NK603 x T25 maize. Japan Biosafety Clearing House (BCH). Tokyo, Japan.
- OECD (1992). Recombinant DNA safety considerations. Organization for economic Cooperation and Development (OECD), Paris, France.
- OECD (1993). Safety considerations for biotechnology: scale-up of crop plants. Organization for economic Cooperation and Development (OECD), Paris, France.
- OECD (1997). Consensus document on the biology of *Brassica napus* L. (oilseed rape). Organization for economic Cooperation and Development (OECD), Paris, France.
- OECD (1999a). Consensus document on general information concerning the genes and their enzymes that confer tolerance to phosphinothricin herbicide. Organization for economic Cooperation and Development (OECD), Paris, France. [http://www.olis.OECD.org/olis/1999doc.nsf/LinkTo/env-jm-mono\(99\)13](http://www.olis.OECD.org/olis/1999doc.nsf/LinkTo/env-jm-mono(99)13)
- OECD (1999b) Consensus Document on the Biology of *Oryza sativa* (Rice) No. 14, 1999 Organization for economic Cooperation and Development (OECD), Paris, France. [http://www.olis.OECD.org/olis/1999doc.nsf/LinkTo/env-jm-mono\(99\)26](http://www.olis.OECD.org/olis/1999doc.nsf/LinkTo/env-jm-mono(99)26)
- OECD (2000) Consensus document on the biology of *Glycine max* (L.) Merr. Organization for economic Cooperation and Development (OECD), Paris, France.
- OECD (2001). Consensus document on the biology of *Beta vulgaris* L. Organization for economic Cooperation and Development (OECD), Paris, France.
- OECD (2002). Module II: Herbicide Biochemistry, Herbicide Metabolism and the Residues in Glufosinate-Ammonium (Phosphinothricin)-Tolerant Transgenic Plants. Organization for economic Cooperation and Development (OECD), Paris, France. <http://www.OECD.org/officialdocuments/displaydocumentpdf?cote=ENV/JM/MONO%282002%2914&doclanguage=en>
- OECD (2003a). Consensus document on the biology of *Zea mays* subsp. *Mays*. Organization for economic Cooperation and Development (OECD), Paris, France.
- OECD (2003b). Description of selected key generic terms used in chemical hazard/risk assessment. Organization for economic Cooperation and Development (OECD), Paris. [http://www.olis.OECD.org/olis/2003doc.nsf/LinkTo/NT0000472/\\$FILE/JT00152557.PDF](http://www.olis.OECD.org/olis/2003doc.nsf/LinkTo/NT0000472/$FILE/JT00152557.PDF)
- OECD (2006). Points to consider for consensus documents on the biology of cultivated plants. Organization for economic Cooperation and Development (OECD), Paris, France.
- OECD (2008). Consensus document on the biology of cotton (*Gossypium* spp.). Organization for economic Cooperation and Development (OECD), Paris, France.
- OGTR (2003). Risk assessment and risk management plan: DIR 021/2002 Commercial release of genetically modified (InVigor hybrid) canola (MS1, RF1, RF2, RF3, T45, Topas 19/2, HCN92, MS1, MS8). Office of the Gene Technology Regulator (OGTR), Canberra, Australia. <http://www.OGTR.gov.au/internet/OGTR/publishing.nsf/Content/dir021-2002>
- OGTR (2006). Risk assessment and risk management plan: DIR 062/2005 Commercial release of herbicide tolerant Liberty Link cotton for use in the Australian cropping system. Office of the Gene Technology Regulator (OGTR), Canberra, Australia. <http://www.OGTR.gov.au/internet/OGTR/publishing.nsf/Content/dir062-2005>
- OGTR (2008). The biology of *Gossypium hirsutum* L. and *Gossypium barbadense* L. Office of the gene technology regulatory (OGTR) Department of Health and Ageing, Canberra, Australia.
- Philippines (2005). Determination of the Safety of Syngenta's Corn BT11 (Insect-resistant and herbicide-tolerant Corn) for Direct use as Food, Feed, and for Processing and for Propagation. Philippines Department of Agriculture, Bureau of Plant Industries, Manila, Philippines.
- USDA APHIS (1994a). Petition for determination of nonregulated status of Ciba Seeds' corn genetically engineered to express the Cry1A(b) protein from *Bacillus thuringiensis* subspecies *kurstaki*. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/94_31901p.pdf
- USDA APHIS (1994b). Petition for determination of nonregulated status for glufosinate resistant corn

- transformation events T14 and T25. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/94_35701p.pdf
- USDA APHIS (1995a). Environmental assessment and finding of no significant impact: Petition 94-319-01 for determination of nonregulated status for event 176 corn. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/94_31901p_com.pdf
- USDA APHIS (1995b). Petition for determination of nonregulated status: Glufosinate resistant corn line B16. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/95_14501p.pdf
- USDA APHIS (1995c). Environmental assessment and finding of no significant impact: Petition 95-145-01 for determination of nonregulated status for glufosinate resistant corn transformation line B16. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/95_14501p_com.pdf
- USDA APHIS (1995d). Petition for determination of nonregulated status for: Insect protected corn (*Zea mays* L.) expressing the CryIA(b) gene from *Bacillus thuringiensis* var. *kurstaki*(Bt11). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/95_19501p.pdf
- USDA APHIS (1995e). Petition for determination of nonregulated status for male sterile, glufosinate tolerant corn transformation event MS3. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/95_22801p.pdf
- USDA APHIS (1995f). Environmental assessment and finding of no significant impact: Petition 94-357-01 for determination of nonregulated status for glufosinate resistant corn transformation events T14 and T25. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/94_35701p_com.pdf
- USDA APHIS (1996a). Petition for determination of nonregulated status for glufosinate resistant soybean transformation events (W62, W98, A2704-12, A2704-21 and A5547-35). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/96_06801p.pdf
- USDA APHIS (1996b). Environmental assessment and finding of no significant impact: AgroEvo USA company petition 96-068-01p for determination of nonregulated status for transgenic glufosinate resistant soybean (GRS) lines W62, W98, A2704-12, A2704-21 and A5547-35. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/96_06801p_com.pdf
- USDA APHIS (1996c). Environmental assessment and finding of no significant impact: Petition 95-195-01 for determination of nonregulated status for Bt11 corn. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/95_19501p_com.pdf
- USDA APHIS (1996d). Petition for determination of nonregulated status: insect protected corn (*Zea mays* L.) with cryIA(c) gene from *Bacillus thuringiensis* subsp. *kurstaki* (DBT418). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/96_29101p.pdf
- USDA APHIS (1996e). Environmental assessment and finding of no significant impact: Petition 95-228-01 for determination of nonregulated status for MS3 corn. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/95_22801p_com.pdf
- USDA APHIS (1997a). Petition for determination of nonregulated status: Glufosinate tolerant canola transformation event T45. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/97_20501p.pdf
- USDA APHIS (1997b). Petition for determination of nonregulated status for *Radicchio rosso* lines with male sterility. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/97_14801p.pdf
- USDA APHIS (1997c). Environmental assessment and finding of no significant impact: USDA/APHIS Petition 97-148-01p for determination of nonregulated status for *Radicchio rosso* lines designated as RM3-3, RM3-4, and RM3-6. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/97_14801p_com.pdf
- USDA APHIS (1997d). Petition for determination of nonregulated status for male sterile corn lines 676, 678, and 680 with resistance to glufosinate. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/97_34201p.pdf
- USDA APHIS (1997e). Petition for determination of nonregulated status for Bt Cry9C insect resistant, glufosinate tolerant corn transformation event CBH-351. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/97_26501p.pdf
- USDA APHIS (1997f). Environmental assessment and finding of no significant impact: USDA/APHIS Petition 96-291-01p for determination of nonregulated status for insect-protected corn line DBT418. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/96_29101p_com.pdf
- USDA APHIS (1998a). Environmental assessment and finding of no significant impact: AgroEvo USA Company Petition

- 97-205-01p for determination of nonregulated status for transgenic glufosinate tolerant canola transformation (event T45). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/97_20501p_com.pdf
- USDA APHIS (1998b). Petition for determination of nonregulated status: Glufosinate tolerant sugar beet, transformation event T120-7. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/97_33601p.pdf
- USDA APHIS (1998c). Environmental assessment and finding of no significant impact: AgrEvo USA Company Petition 97-336-01p for determination of nonregulated status for transgenic glufosinate tolerant sugar beet transformation event T120-7. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/97_33601p_com.pdf
- USDA APHIS (1998d). Application for an extension of determination of nonregulated status for glufosinate resistant soybean transformation events (A5547-127). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/98_01401p.pdf
- USDA APHIS (1998e). Environmental assessment and finding of no significant impact: Request for extension of determination of nonregulated status for glufosinate resistant soybean transformation events (A5547-127). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/98_01401p_com.pdf
- USDA APHIS (1998f). Application for an extension of determination of nonregulated status for glufosinate resistant soybean transformation events (GU262). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/98_23801p.pdf
- USDA APHIS (1998g). Environmental assessment and finding of no significant impact: application for an extension of determination of nonregulated status for glufosinate resistant soybean transformation events (GU262). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/98_23801p_com.pdf
- USDA APHIS (1998h). Petition for determination of nonregulated status for LibertyLink rice transformation events LLRice06 and LLRice62. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/98_32901p.pdf
- USDA APHIS (1998i). Environmental assessment and finding of no significant impact: USDA/APHIS Petition 97-342-01p for determination of nonregulated status for genetically engineered corn lines 676, 678, and 680. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/97_34201p_com.pdf
- USDA APHIS (1998j). Environmental assessment and finding of no significant impact: Petition 97-265-01p for determination of nonregulated status for Bt. Cry9C insect resistant, glufosinate tolerant corn transformation event CBH-351. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/97_26501p_com.pdf
- USDA APHIS (1998k). Request for extension of a determination of nonregulated status for a hybrid seed production system in corn based on male sterility and glufosinate tolerance as a marker (95-228-01p): event MS6. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/98_34901p.pdf
- USDA APHIS (1998l). Petition for determination of nonregulated status for InVigor hybrid canola transformation events MS8/RF3. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/98_27801p.pdf
- USDA APHIS (1999a). Environmental assessment and finding of no significant impact: AgrEvo USA Petition 98-278-01p for determination of nonregulated status for canola transformation events MS8 and RF3 genetically engineered for pollination control and tolerance to glufosinate herbicide. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/98_27801p_com.pdf
- USDA APHIS (1999b). Environmental assessment and finding of no significant impact: AgrEvo USA Petition 98-329-01p for determination of nonregulated status for glufosinate tolerant rice transformation events LLRice06 and LLRice62. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/98_32901p_com.pdf
- USDA APHIS (1999c). Extension of determination of nonregulated status for corn genetically engineered for male sterility and glufosinate herbicide tolerance as a marker (MS6). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/98_34901p_com.pdf
- USDA APHIS (2000). Petition for the determination of nonregulated status B.t. Cry1F insect resistant, glufosinate tolerant maize line. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/00_13601p.pdf
- USDA APHIS (2001a). Application for an extension of the determination of nonregulated status for glufosinate-tolerant canola transformation (98-278-01p): MS1/RF1/RF2(01-206-01). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/01_20601p.pdf
- USDA APHIS (2001b). Extension of determination of nonregulated status for canola genetically engineered for

- male sterility, fertility restoration, and glufosinate herbicide tolerance (MS1, RF1, RF2). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/01_20601p_com.pdf
- USDA APHIS (2001c). Environmental assessment and finding of no significant impact: Approval of Mycogen Seeds c/o Dow AgroSciences LLC and Pioneer Hi-Bred International, Inc. request (00-136-01p) seeking a determination of non-regulated status for Bt Cry1F insect resistant, glufosinate tolerant corn line 1507. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/00_13601p_com.pdf
- USDA APHIS (2002a). Application for an extension of the determination of nonregulated status for glufosinate-tolerant canola transformation (97-205-01p): Topas 19/2 (01-206-02) (HCN92). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/01_20602p.pdf
- USDA APHIS (2002b). Extension of determination of nonregulated status for canola genetically engineered for glufosinate herbicide tolerance (HCN92). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/01_20602p_com.pdf
- USDA APHIS (2002c). Aventis CropScience USA LP petition for the determination of nonregulated status for glufosinate-tolerant cotton transformation event, 02-042-01p (LLcotton25). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/02_04201p.pdf
- USDA APHIS (2003a). Petition for determination of nonregulated status: B.t. Cry1F insect-resistant cotton event 281-24-236. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/03_03601p.pdf
- USDA APHIS (2003b). Petition for determination of nonregulated status: B.t. Cry1Ac insect-resistant cotton event 3006-210-23. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/03_03602p.pdf
- USDA APHIS (2003c). Environmental assessment and finding of no significant impact: Aventis CropScience USA LP petition for the determination of nonregulated status for glufosinate-tolerant cotton transformation event, 02-042-01p(LLcotton25). United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/02_04201p_com.pdf
- USDA APHIS (2004a). Environmental assessment and finding of no significant impact: petition for determination of nonregulated status: B.t. Cry1F insect-resistant cotton event 281-24-236 and Cry1Ac insect-resistant cotton event 3006-210-23. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/03_03602p_com.pdf
- USDA APHIS A (2004b). Application for an extension of the determination of nonregulated status for B.t. Cry1F insect-resistant, glufosinate tolerant maize (00-136-01p): maize line 6275. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/03_18101p.pdf
- USDA APHIS A (2004c). Environmental assessment and finding of no significant impact: approval of Mycogen Seeds c/o Dow AgroSciences request (03-181-01p) seeking extension of the determination of nonregulated status for Bt Cry1F insect-resistant, glufosinate tolerant corn line 6275. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/03_18101p_com.pdf
- USDA APHIS A (2004d). Application for the determination of nonregulated status for B.t. Cry34/35Ab1 insect-resistant, glufosinate tolerant corn: corn line 59122. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/03_35301p.pdf
- USDA APHIS A (2005). Environmental assessment and finding of no significant impact: request 03-353-01p seeking a determination of non-regulated status for B.t. Cry34Ab1/35Ab1 insect-resistant, glufosinate tolerant corn line 59122-7. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/03_35301p_com.pdf
- USDA APHIS (2006a). Application for an Extension of the Determination of Nonregulated Status for Glufosinate-Tolerant Rice (98-329-01p): Transformation Event LLRICE601. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs/06_23401p.pdf
- USDA APHIS (2006b). Environmental assessment and finding of no significant impact: extension of nonregulated status to rice line LLRice601. United States Department of Agriculture, Animal and Plant Health Inspection Service, Washington D.C. http://www.aphis.usda.gov/brs/aphisdocs2/06_23401p_com.pdf
- USEPA (2001). Biopesticide registration action document: *Bacillus thuringiensis* Cry1F corn. United States Environmental Protection Agency, Washington D.C. http://www.epa.gov/pesticides/biopesticides/ingredients/tech_docs/brad_006481.pdf
- USEPA (2005). *Bacillus thuringiensis* Cry1F/Cry1Ac construct 281/3006 insecticidal crystal protein as expressed in cotton (006445 & 006481) fact sheet. United States Environmental Protection Agency, Washington D.C. http://www.epa.gov/oppbppd1/biopesticides/ingredients/factsheets/factsheet_006445-6481.htm
- USEPA (2009a). *Bacillus thuringiensis* Cry1Ab Delta-Endotoxin Protein and the Genetic Material Necessary for Its

Production (via Elements of Vector pZO1502) in Event Bt11 Corn (OECD Unique Identifier: SYN-BT011-1)(006444) & *Bacillus thuringiensis* Vip3Aa20 Insecticidal Protein and the Genetic Material Necessary for Its Production (via Elements of Vector pNOV1300) in Event MIR162 Maize (OECD Unique Identifier: SYN-IR162-4)(006599) Fact Sheet. United States Environmental Protection Agency, Washington D.C. http://www.epa.gov/opp00001/biopesticides/ingredients/factsheets/factsheet_006599-006444.html

USEPA (2009b). Pesticide Fact Sheet (MON 89034 x TC1507 x MON 88017 x DAS-59122-7). United States Environmental Protection Agency, Washington D.C. <http://www.epa.gov/oppbpd1/biopesticides/pips/smartstax-factsheet.pdf>

ANNEX I: SUMMARY OF PAT PROTEIN EXPRESSION DATA

The tables that follow present summary data from peer-reviewed publications and regulatory submissions. Additional information on collection and sampling methodologies can be found in the referenced sources.

Table I.1. Quantities of PAT in *Beta vulgaris* event T120-7 as detected by ELISA (USDA APHIS, 1998b).

Plant Matrix ¹	% Protein ²	ng PAT/g protein ^{3,4}
Roots	6.8	137
Tops (above ground)	15.0	966
Pulp (dried)	9.7	ND
Molasses	9.9	ND

¹ Values reported are mean values from all sites.

² Literature values (see USDA APHIS, 1998b for citation).

³ Two extracts from each sample (18 tops; 18 roots from 6 field sites) were analyzed in triplicate.

⁴ Limit of detection = 2 ng/g root; 1.6 ng/g sugar, pulp; 0.4 ng/g molasses.

⁵ ND = not detected.

Table I.2. PAT contents in seed samples for *Brassica napus* event Topas 19/2 (HCN10 and HCN92) as detected by ELISA (USDA APHIS, 2002a).

Sample ID	PAT / Sample (ng/g)
Excel 1996 (control)	ND ¹
HCN92	295
HCN92	295
HCN10	189
HCN10	202

¹ ND < Limit of quantification (0.40 ng/mL).

Table I.3. PAT expression in seed and leaf samples from *Brassica napus* lines Topas 19/2, T45, and Topas 19/2 x T45 as detected by ELISA (USDA APHIS, 2002a).

Sample ID	Line/Treatment	Event	PAT/Sample (ng/g)	Total Protein (mg/g)	PAT/Protein (%)
Control		Topas 19/2	ND	9.51	
Plot 2	SW9782179	Topas 19/2	248	2.14	0.012
Plot 7	SW9782179	Topas 19/2	263	3.04	0.009
Plot 8	SW9782179	Topas 19/2	309	2.24	0.014
Plot 10	SW9782179	Topas 19/2	379	2.61	0.015
Plot 3	SW9782180	T45	555	2.68	0.021
Plot 5	SW9782180	T45	743	2.41	0.031
Plot 6	SW9782180	T45	717	2.17	0.033
Plot 1	SW9782213	Topas 19/2//T45	754	2.01	0.038
Plot 4	SW9782213	Topas 19/2//T45	906	2.00	0.045
Plot 9	SW9782213	Topas 19/2//T45	932	3.37	0.028
Plot 11	SW9782213	Topas 19/2//T45	944	3.12	0.030
Seed - UN	Untreated	Topas 19/2//T45	563	54.3	0.00104
Seed-TR	Treated ¹	Topas 19/2//T45	669	59.4	0.00113
Tmeal-UN	Untreated	Topas 19/2//T45	ND	85.9	ND
Tmeal-TR	Treated ¹	Topas 19/2//T45	ND	76.1	ND
RBD oil - UN	Untreated	Topas 19/2//T45	ND	ND	ND
RBD oil - TR	Treated ¹	Topas 19/2//T45	ND	ND	ND

¹ Treated with glufosinate.

Table I.4. PAT expression in seeds and leaves of *Brassica napus* lines RF3 and MS8 determined by enzyme activity (OGTR, 2003).

GM Canola Line	Seed $\mu\text{g PAT/mg Total Protein}$	Leaf $\mu\text{g PAT/mg Total Protein}$
RF3	0.10	1.33
MS8	0.04	0.51
RF1	Not tested	1.45
RF2	Not tested	0.7
MS1	Not tested	0.9

Table I.5. PAT expression in seeds of GM *Brassica napus* lines determined by ELISA (OGTR, 2003).

GM Canola Line	$\mu\text{g PAT/g Seed}$	$\mu\text{g PAT/mg Total Protein}$
RF3	0.69	0.012
MS8	0.07	0.002
RF3xMS8	0.34	0.013
RF1	0.50	0.015
RF2	0.42	0.012
MS1	0.07	0.002
MS1xRF1	0.20	0.006
MS1xRF2	0.35	0.007

Table I.6. PAT expression in seeds of GM *Brassica napus* lines determined by ELISA (OGTR, 2003).

GM Canola Line	Seed $\mu\text{g PAT/g Total Seed}$	Leaf $\mu\text{g PAT/mg Fresh Weight}$
T45	0.561	0.348
Topas	0.47	0.0843

Table I.7. Summary of mRNA expression analysis for *bar* in *Brassica napus* RF2 (USDA APHIS, 2001a).

Total RNA	$\text{pg } bar \text{ mRNA}/\mu\text{g Total RNA}$ (Range of Detected Values)
Leaves	0.8–1.6
Flower Buds	0.1–0.2
Seed	ND ¹
Pollen	ND

¹ ND = not detected (<2pg/ μg total RNA).

Table I.8. Summary of mRNA expression analysis for *bar* in *Brassica napus* MS8 (USDA APHIS, 1998)¹.

Total RNA	Transgene Expression (pg/ μg Total RNA) ²
Leaf A	0.03
Leaf B	0.22
Flower buds 2mm A	0.14
Flower buds 2mm B	0.11
Flower buds 3mm A	0.19
Flower buds 3mm B	0.03
Dry seed	ND ³

¹ Data shown for two plants (A and B) with single samples of each tissue.

² pGembar/SP6 plasmid used for preparation of RNA probe.

³ ND = not detected. Limit of detection = 0.1pg/g total RNA.

Table I.9. Summary of mRNA expression analysis for *bar* in *Brassica napus* RF3 (USDA APHIS, 1998).

Total RNA	Transgene Expression (pg/ μg Total RNA) ¹
Leaf A	1.1
Leaf B	0.2
Flower buds 2mm A	0.46
Flower buds 2mm B	0.52
Flower buds 3mm A	0.38
Flower buds 3mm B	0.34
Dry seed	ND ²
Pollen	ND ²

¹ Data shown for two plants (A and B) with single samples of each tissue.

² ND = not detected. Limit of detection = 0.05pg/g total RNA.

Table I.10. Summary of PAT expression in seed of *Brassica napus* lines T45 (HCN28) and HCN 92 (method not reported) (CFIA, 1996b).

Sample	$\mu\text{g/mg Sample}$ (Reported Range)
T45 Seed	95–245
HCN 92 Seed	150–223

Table I.11. Summary of PAT expression in *Brassica napus* line T45 (HCN28) using Northern blot and ELISA (EFSA, 2008).

Sample	Northern Blot ¹	ELISA
Leaves	+	NA ²
Stems	+	NA
Roots	+	NA
Seeds	–	930 ng/g dry weight

¹ Results are reported as presence (+) or absence (–) of detectable mRNA.

² NA = not available.

A review of the environmental safety of the PAT protein

Table I.12. Summary of PAT content of seed from *Brassica rapa* line HCR-1 (method not reported) (CFIA, 1998c).

Tissue	Mean (ng/g)	Range (ng/g)
Seed	107	84–132

Table I.13. Summary of PAT protein in seeds of *Glycine max* lines AS2704-12 and A5547-127 as detected by ELISA (FSANZ, 2004a).

Sample	Year	Treatment ¹	PAT/Sample (ng/g)	Crude Protein (%)	PAT Protein as % of Crude Protein
AS2704-12	1997	NA ²	573	37–45	0.00016
A5547-127	1998	NA	10800	37–45	0.00292
AS2704-12	1999	sprayed	879 (264) ³	NA	0.000227
AS2704-12	1999	unsprayed	862 (272)	NA	0.000227
A5547-127	1999	sprayed	10100 (816)	NA	0.00285
A5547-127	1999	unsprayed	9971 (940)	NA	0.00283
AS2704-12	NA	sprayed	2183	38.9	0.00050
AS2704-12	NA	unsprayed	1948	38.5	0.00056
A5547-127	NA	sprayed	17471	36.5	0.0048
A5547-127	NA	unsprayed	20202	35.8	0.0056

¹ sprayed = treated with glufosinate; unsprayed = not treated with glufosinate.

² NA = not available.

³ Mean (Standard Deviation).

Table I.14. Summary of PAT protein detected in *Glycine max* line A2704-12 using ELISA (Japan BCH, 1999b).

Tissue	Mean PAT (µg/g Fresh Weight) ± Standard Deviation	Crude Protein (% of Fresh Weight)	PAT Protein (% of Crude Protein)
Root	2.23 ± 1.29	1.95	0.011
Stem	7.63 ± 2.20	3.58	0.021
Leaf	14.5 ± 2.4	5.96	0.024

Table I.16. Summary of PAT protein detected in *Glycine max* line A5547-127 using ELISA (Japan BCH, 2006f).

Tissue	Mean PAT (µg/g Fresh Weight) ± Standard Deviation	Crude Protein (% of Fresh Weight)	PAT Protein (% of Crude Protein)
Root	3.73 ± 0.98	2.15	0.017
Stem	11.5 ± 1.8	3.62	0.032
Leaf	19.0 ± 5.0	6.70	0.028

Table I.15. Summary of PAT protein detected in seeds of *Glycine max* line A2704-12 using ELISA (Japan BCH, 1999b).

Sample	PAT (ng/g Sample) Mean ± Standard Deviation	Crude Protein Content (%)	PAT/Crude Protein (%)
1	1057	NA ¹	NA
2	573	NA	NA
3	862 ± 268	38.03	0.000227
4	2138 ± 33	43.5	0.00049

¹ NA = not available.

Table I.17. Summary of PAT protein detected in seeds of *Glycine max* line A5547-127 using ELISA (Japan BCH, 2006f).

Sample	PAT (ng/g Sample) Mean ± Standard Deviation	Crude Protein Content (%)	PAT/Crude Protein (%)
1	6341	NA ¹	NA
2	10800 ± 1210	NA	NA
3	9971 ± 846	35.26	0.00282
4	20202 ± 359	40.4	0.0050

¹ NA = not available.

Table I.18. Summary of PAT content in leaves of *Glycine max* line A5547-127 as detected by ELISA (USDA APHIS, 1998d)¹.

Sample	mg TEP ² /g Sample	µg PAT/g Sample	% PAT/TEP	% PAT/Fresh Weight g/g
A5547-127	4.6	1.72	0.037	1.72 × 10 ⁻⁴

¹ Values are the average from two replicate extractions from two samples of 10 day old seedling leaves.

² TEP = Total Extractable Protein.

Table I.19. Summary of PAT content in leaves of *Glycine max* line GU262 as detected by ELISA (USDA APHIS, 1998f)¹.

Sample	mg TEP ² /g Sample	µg PAT/g Sample	% PAT/TEP	% PAT/Fresh Weight g/g
GU262	4.7	3.03	0.064	3.03 × 10 ⁻⁴

¹ Values are the average from two replicate extractions from two samples of 10 day old seedling leaves.

² TEP = Total Extractable Protein.

Table I.20. Summary of PAT protein in *Glycine max* lines W62 and W98 as detected by enzymatic activity (USDA APHIS, 1996a)¹.

Tissue	Site and Year	Plant	µg PAT/g Sample ²
Fodder (whole plant)	Arkansas 1993	W62	10.8 (6.3–15.3)
	Iowa 1993	W98	0.75 (0.65–1.0)
	Illinois 1993	W98	10.9 (9.1–12.7)
Seed	Arkansas 1993	W62	217.0 (147.1–267.3)
	Iowa 1993	W98	27.1 (15.0–39.2)
	Illinois 1993	W98	38.3 (23.5–60.9)

¹ Values are the average from two replicate extractions from two samples of 10 day old seedling leaves.

² Mean (range of observed values).

Table I.21. Summary of PAT protein expression in *Gossypium hirsutum* event 281-24-236 as determined by ELISA (USDA, 2003a).

Tissue	PAT (ng/mg Dry Weight)		
	Mean ¹	Standard Deviation	Minimum-Maximum Range
Young Leaf (3–6 week)	0.43	0.12	0.18–0.67
Terminal Leaf	0.21	0.12	ND ⁴ –0.38
Flower	0.29	0.11	0.07 ² –0.44
Square	0.51	0.15	0.06 ² –0.79
Boll (early)	0.22	0.09	0.08 ² –0.48
Whole Plant (seedling)	0.31	0.07	0.21–0.46
Whole Plant (pollination)	0.23	0.07	0.09 ² –0.33
Whole Plant (defoliation)	0.19	0.13	ND–0.46
Root (seedling)	0.07 ²	0.05	ND–0.12
Root (pollination)	ND	NA ³	ND–0.11
Root (defoliation)	ND	NA	ND–0.11
Pollen ⁵	0.09 ²	0.15	ND–0.45
Nectar ⁵	ND	NA	ND–ND
Seed ⁵	0.47	0.17	0.23 ² –1.02

¹ Calculated from samples at six locations.

² Value below the limit of quantification of the method.

³ NA = not applicable.

⁴ ND = absorbance of the sample was lower than the absorbance of the lowest standard.

⁵ Results are given relative to fresh weight.

Table I.22. Summary of PAT protein expression in *Gossypium hirsutum* even 3006-210-23 as determined by ELISA (USDA, 2004a).

Tissue	PAT (ng/mg Dry Weight)		
	Mean ¹	Standard Deviation	Minimum-Maximum Range
Young Leaf (3–6 week)	ND ⁴	NA ³	0.18–0.67
Terminal Leaf	ND	NA	ND-0.12
Flower	ND	NA	ND-ND
Square	ND	NA	ND-0.08
Boll (early)	ND	NA	ND-0.08
Whole Plant (seedling)	ND	NA	ND-0.09
Whole Plant (pollination)	ND	NA	ND-0.14
Whole Plant (defoliation)	0.11	0.05	ND-0.20
Root (seedling)	ND	NA	ND-0.07
Root (pollination)	ND	NA	ND-ND
Root (defoliation)	ND	NA	ND-ND
Pollen ⁵	ND	NA	ND-ND
Nectar ⁵	ND	NA	ND
Seed ⁵	0.06 ²	0.06	ND-0.23 ²

¹ Calculated from samples at six locations.

² Value below the limit of quantification of the method.

³ NA = not applicable.

⁴ ND = absorbance of the sample was lower than the absorbance of the lowest standard.

⁵ Results are given relative to fresh weight.

Table I.23. Summary of PAT protein expression in *Gossypium hirsutum* even 3006-210-23 as determined by ELISA (USDA, 20036).

Tissue	Mean Protein Expression (ng/mg dry weight)
Young Leaf (3–6 week)	0.43
Terminal Leaf	0.23
Flower	0.35
Square	0.52
Boll (early)	0.27
Whole Plant (seedling)	0.35
Whole Plant (pollination)	0.30
Whole Plant (defoliation)	0.34
Root (seedling)	0.06 ²
Root (pollination)	ND
Root (defoliation)	0.05 ²
Pollen ¹	0.05 ²
Nectar ¹	ND
Seed ¹	0.54

¹ Results reported as ng/mg fresh weight.

² Calculated concentration is less than the limit of quantification of the method.

Table I.24. Summary of PAT expression in *Gossypium hirsutum* line LLCotton25 measured by ELISA (CFIA, 2004).

Tissue	PAT Protein (µg/g Fresh Weight)
Root	7.97
Leaf	52.9
Stem	36.8
Fuzzy Seed	69.9
Cleaned Seed	127
Pollen	19.2

Table I.25. Summary of PAT protein content in *Gossypium hirsutum* line LLCotton25 as measured by ELISA (OGTR, 2006).

Tissue	PAT (µg/g Fresh Weight)		PAT as % of Crude Protein	Average TEP ² (mg/g Fresh Weight ± SD)	Average PAT Content as % of TEP
	Range	Average			
Roots	5.63–10.1	7.97 ± 1.86	0.08	2.26 ± 0.22	0.35
Stems	34.3–45.5	36.8 ± 6.7	0.23	4.99 ± 0.92	0.74
Leaves	45.1–57.3	52.9 ± 6.0	0.19	7.13 ± 0.79	0.74
Pollen (frozen)	4.44–13.0	8.23 ± 3.20	NA ¹	146 ± 8	0.006
Pollen (fresh)	0.11–170	19.3 ± 39.2	NA	107 ± 21	0.018

¹ Not applicable.

² TEP = Total Extractable Protein.

Table I.26. Summary of PAT protein content in leaves of *Gossypium hirsutum* line LLCotton25 as measured by ELISA (OGTR, 2006).

Sample	PAT Protein (µg/g Fresh Weight ± SD)			
	2–4 leaf	4–5 leaf	Early Bloom	Full Bloom
Non GM Control	ND ¹	ND	ND	ND
Sprayed Once ²	NA	85.0 ± 15.6	98.3 ± 16.8	92.6 ± 15.1
Sprayed Twice	NA	NA	NA	92.6 ± 20.3
Unsprayed	57.7 ± 5.3	74.0 ± 12.3	90.2 ± 14.4	75.1 ± 25.6

¹ ND = not detected; NA = not applicable.

² Sprayed with Liberty Link.

Table I.27. Summary of PAT protein content RACs of *Gossypium hirsutum* line LLCotton25 as measured by ELISA (OGTR, 2006).

Sample	Average PAT Protein (µg/mg Fresh Weight) ± SD		Average Protein Content (as % of Crude Protein)	
	LL Sprayed	Unsprayed	LL Sprayed	Unsprayed
Cleaned Seed	127 ± 18	113 ± 24	NA ¹	NA
Lint Coat	1.5 ± 0.45	0.92 ± 0.50	NA	NA
Fuzzy Seed	69.9 ± 6.0	63.0 ± 10.3	0.030	0.027
Lint	0.78 ± 0.63	0.50 ± 0.42	0.003	0.003

¹ NA = not applicable.

Table I.28. Summary of PAT protein in *Gossypium hirsutum* line LLCotton25 as measured by ELISA (USDA APHIS, 2002c).

Sample	PAT Protein (µg/g Fresh Weight) ± Standard Deviation		PAT Content (% of Crude Protein)	
	Liberty Herbicide	Conventional Herbicide	Liberty Herbicide	Conventional Herbicide
Cleaned Seed	127 ± 18	113 ± 24	NA	NA
Lint Coat	1.15 ± 0.45	0.92 ± 0.50	NA	NA
Fuzzy Seed	69.9 ± 6.0	63.0 ± 10.3	0.030	0.027
Lint	0.78 ± 0.63	0.50 ± 0.42	0.003	0.003

Table I.29. Summary of PAT protein content in leaves of *Gossypium hirsutum* line LLCotton25x15985 as measured by ELISA (Japan BCH, 2006a).

Sample	Protein Content (Mean ± Standard Deviation) (µg/g of Leaf)
LLCotton25 x 15985	60.9 ± 8.7 ¹
LLCotton25	65.9 ± 10.6 ¹

¹ The difference in expression between the stacked line and the single event is not significant.

Table I.30. Summary of PAT protein content in *Oryza sativa* line LLRice62 as measured by ELISA (CFIA, 2006b).

Tissue	Mean Protein Content µg/g Fresh Weight
Grain	12.1
Straw	75.3
Rice Hulls	1.56
Roots	12.7
Stems	30.9
Leaves	84.7

Table I.31. Summary of PAT protein content in *Oryza sativa* line LLRICE62 as measured by ELISA (FSANZ, 2008).

Tissue	Average PAT Content (µg/g ± SD)	Crude Protein (% w/w)	PAT Protein (% of Crude Protein)
Grain (Year 1)	12.1 ± 0.6	7.19	0.017
Straw (Year 1)	75.3 ± 4.4	2.38	0.316
Grain (Year 2)	10.6 ± 1.3	7.41	0.014

Table I.32. Summary of PAT protein content in grain of *Oryza sativa* lines LLRICE06 and LLRICE62 as detected by ELISA (USDA APHIS, 1998h).

Plant	mg TEP ¹ /g Sample	µgPAT/g Sample	% PAT/TEP	% PAT/Fresh Weight (g/g)
LLRICE06	1.89 ± 0.49	0.419 ± 0.04	0.02	0.00005
LLRICE62	2.54 ± 0.09	12.4 ± 2.4	0.63	0.00124

¹ TEP = Total Extractable Protein

Table I.33. Summary of PAT protein content in seed of *Oryza sativa* line LLRICE601 (method not reported) (USDA APHIS, 2006a).

Plant	Protein Content (ng/g Fresh Weight)	% of Crude Rice Protein
LLRICE601	120	0.00034

Table I.34. PAT protein expression in *Zea mays* line Bt-176 and hybrid plants as determined by ELISA (USDA APHIS, 1995a)¹.

		PAT Expression ng/g Fresh Weight			
		Seedling	Anthesis	Seed Maturity	Senescence
Leaves	Bt176	<2004	<200	<200	ND
	176 × 554 ²	<200	ND	ND	ND
	176 × 564 ³	<200	ND	<200	<200
Whole Plant	Bt176	ND	<200	<200	<200
	176 × 554	<200	ND	<200	<200
	176 × 564	<200	ND	ND	<200
Kernels	Bt176			ND	ND
	176 × 554			ND	ND
	176 × 564			ND	ND
Pollen	Bt176		ND		
	176 × 554		NA		
	176 × 564		ND		
Roots	Bt176	ND ⁴	<100	<100	NA
	176 × 554	ND	<100	<100	NA
	176 × 564	ND	ND	<100	NA
Pith	Bt176	NA ⁵	<200	<200	NA
	176 × 554	NA	ND	<200	NA
	176 × 564	NA	NA	ND	NA

¹ Blank cell indicate no developmental relevance.

² 176 × 554 = hybrid progeny of CG00526-176 and untransformed CG00554 and is hemizygous for the introduced genes.

³ 176 × 564 = hybrid progeny of CG00526-176 and untransformed CG00564 and is hemizygous for the introduced genes.

⁴ When trace amounts were detectable but not quantifiable, results are shown as < lower limit of quantification.

⁵ ND = not detected.

⁶ NA = not analyzed.

Table I.35. Expression of PAT protein in leaf tissue of male sterile *Zea mays* lines 676, 678, and 680 as determined with ELISA (USDA APHIS, 1997d).

Male Sterile Corn Line	PAT Concentration µg/g Total Protein
676	601–617
678	204–278
680	<20 ¹

¹ Below the limit of quantification (20µg/g).

Table I.36. Expression of PAT protein in *Zea mays* line B16 (method not reported) (CFIA, 1996d).

Tissue	PAT Protein Detected
Leaves	1.0–4.6 mg/g protein
Roots	+ ¹
Stalk	+
Tassel	+
Cob	+
Husk	+
Kernels	– ²
Silk	–
Pollen	–

¹ + = Protein detected but quantity not reported.

² – = Protein not detected.

Table I.37. Expression of PAT protein in *Zea mays* line B16 determined by Western blot (USDA APHIS, 1995b).

Tissue	PAT Concentration (ng/µg Total Protein)	PAT Concentration (ng/mg Fresh Weight)
Coleoptile (6 days)	1.8	13.8
Leaf (24 days)	1.0	55.6
Leaf (44 days)	2.8 ± 0.1	166.0 ± 24.2
Leaf (93 days)	2.1	106.1
F ₂ ovule (0 days pp)	0.8	5.5
Immature F ₂ seed (16 days pp)	0.3	3.4
Immature F ₂ seed (45 days pp)	0.3	5.7
Hybrid seed (F ₁)	0.2	1.9
Root (24 days)	0.06	4.2
Root (44days)	1.3	8.1
Prop root (49 days)	1.9 ± 0.3	19.8 ± 2.4
Cob (56 days)	2.2	67.2
Husk (56 days)	1.1	2.5
Silk	ND ¹	4.2
Stalk (24 days)	2.0	
Stalk (77 days)	4.6 ± 0.4	15.7
Immature tassel (49 days)	2.0	11.2 ± 2.7
Pollen	ND	30.8
Silage	ND	

¹ ND = not detected. <0.05 ng/mg in silk; <0.08 ng/mg in pollen and silage.

Table I.38. Mean level of expression of the PAT protein in *Zea mays* line Bt11 (X4334-CBR and X4743) using ELISA (ANZFA, 2001a).

	Mean (µg/g Fresh Weight)			
	Leaf	Kernel	Husk	Stalk
X4334-CBR	0.0386 ± 0.0029	lod ¹	lod	lod
X4734-CBR	0.0494 ± 0.005	lod	nd ²	nd
Control (NK4242)	lod	lod	lod	lod

¹ lod (limit of detection) for the procedure is 1ng PAT/mL extract. These values are considered not above background.

² nd = no data.

A review of the environmental safety of the PAT protein

Table I.39. Mean level of expression of the PAT protein in *Zea mays* line Bt11 (method not reported) (CFIA, 1996c).

	Mean ($\mu\text{g/g}$ fresh weight)					
	Leaf	Tassel	Silk	Roots	Kernel	Pollen
BT11	0.049	0.027	0.005	ND ¹	ND	ND

¹ ND = not detected.

Table I.40. Mean expression of PAT protein in *Zea mays* line CBH-351 using ELISA (USDA APHIS, 1997e)¹.

Tissue	Stage 1		Stage 2		Stage 3		Stage 4	
	Total Protein (mg/g)	PAT Protein ($\mu\text{g/g}$)	Total Protein (mg/g)	PAT Protein ($\mu\text{g/g}$)	Total Protein (mg/g)	PAT Protein ($\mu\text{g/g}$)	Total Protein (mg/g)	PAT Protein ($\mu\text{g/g}$)
Whole Plant	12.5 \pm 4.6	189.7 \pm 23.5	7.8 \pm 1.9	105.7 \pm 16.7	5.7 \pm 1.4	44.4 \pm 5.4	2.9 \pm 1.6	6.8 \pm 1.6
Leaf	3.1 \pm 1.3	45.4 \pm 9.7	0.6 \pm 0.1	14.0 \pm 0.1	1.5 \pm 0.1	2.3 \pm 0.2	1.8 \pm 0.2	0.0 \pm 0.0
Root	0.7 \pm 0.1	39.1 \pm 2.7	25.8 \pm 19.2	4.4 \pm 0.8	0.3 \pm 0.2	0.6 \pm 0.5	0.6	2.2 \pm 2.8
Stalk	NA ²	NA	2.8 \pm 1.2	0.5 \pm 0.5	0.0 \pm 0.0	0.5 \pm 0.2	\pm 0.2	0.0 \rightarrow 0.1
Tassel	NA	NA	175.0 \pm 100.4	4.2 \pm 1.2	NA	NA	NA	NA
Kernel	NA	NA	NA	NA	NA	NA	6.7 \pm 1.1	17.8 \pm 5.0

¹ Values are expressed as mean \pm standard deviation. All values are relative to dry weight of samples.

² NA = not applicable.

Table I.41. Mean level of PAT protein expression in hybrids derived from *Zea mays* line CBH-351 using ELISA (USDA APHIS, 1997e)¹.

Hybrid	Stage 1		Stage 2	
	Total Protein (mg/g)	PAT Protein ($\mu\text{g/g}$)	Total Protein (mg/g)	PAT Protein ($\mu\text{g/g}$)
Hybrid A	19.1 \pm 8.2	364.0 \pm 85.1	4.9 \pm 2.0	40.7 \pm 16.7
Hybrid B	18.3 \pm 1.1	291.0 \pm 4.2	7.2 \pm 2.8	77.1 \pm 12.9
Hybrid C	11.6 \pm 6.3	227.0 \pm 33.4	6.8 \pm 1.3	100.2 \pm 24.9
Hybrid D	9.1 \pm 3.3	176.3 \pm 63.0	4.1 \pm 1.4	88.3 \pm 23.4

¹ Values are expressed as mean \pm standard deviation. All values are relative to dry weight of samples.

Table I.42. Mean expression of PAT protein in *Zea mays* line DAS-06275-8 at six field trial sites in the U.S. and Canada as measured by ELISA (CFIA, 2006a).

	Leaf	Root	Stalk	Grain	Pollen	Forage
PAT Protein Expression (ng/mg Dry Weight)	129.2–224.21	61.1–70.0 ¹	103.3	8.94	0.73	106.9

¹ Range represents means measured across multiple growth stages.

Table I.43. Mean expression of PAT protein in *Zea mays* line DAS-06275-8 in grain (method not reported) (Health Canada, 2006).

	U.S.	Chile
PAT Protein Expression (ng/mg Dry Weight)	5.94	23

Table I.44. Summary of mean expression of BAR (PAT)protein in a *Zea mays* hybrid derived from DAS-06275-8 measured by ELISA (USDA APHIS, 2004b).

Growth Stage	Tissue	Mean ± Standard Deviation (ng/mg Dry Weight)	Minimum-Maximum Range (ng/mg Dry Weight)
V9	Leaf	323 ± 91.0	0–538
	Root	112 ± 35.3	0–170
	Whole plant	5 ± 3.50	1–11
R1	Leaf	674 ± 98.1	539–935
	Root	253 ± 162	61–673
	Whole plant	72 ± 32.9	35–108
R4	Pollen	0 ± 0.766	0–4.07
	Stalk	282 ± 68.5	177–475
	Leaf	682 ± 254	451–1584
Maturity	Root	223 ± 105	85–511
	Forage	7 ± 7.05	1–19
	Grain	23 ± 6.33	13–33
Senescence	Leaf	0 ± 0.461	0–1
	Root	41 ± 49.5	0–148
	Whole plant	18 ± 5.27	9–23

Table I.45. Summary of PAT protein expression in *Zea mays* line DAS-59122-7 at multiple sites in the U.S. and Canada measured by ELISA (CFIA, 2005).

	Leaf	Root	Stalk	Grain	Pollen	Forage
PAT Protein Expression (ng/mg Dry Weight)	0.25–11.4 ¹	0.18–0.42 ¹	0.38	0.1	LOD ²	2.4

¹ Range represents means measured across multiple growth stages.

² LOD = below the limit of detection (<0.30 ng/mg dry weight).

Table I.46. Summary of expression levels of PAT protein in *Zea mays* line DAS-59122-7 as measured by ELISA (USDA APHIS, 2004d).

Growth Stage	Tissue	Mean ± Standard Deviation (ng/mg Dry Weight)	Minimum-Maximum Range (ng/mg Dry Weight)
V9	Leaf	11.1 ± 3.68	5.61–19.2
	Root	0.47 ± 0.15	0.27–0.87
	Whole plant	0.18 ± 0.13	0–0.40
R1	Leaf	11.2 ± 3.49	6.36–18.2
	Root	0 ± 0	0–0
	Whole plant	0.13 ± 0.03	0.07–0.20
R4	Pollen	0.27 ± 0.12	0.11–0.62
	Stalk	0.13 ± 0.23	0–0.58
	Leaf	8.13 ± 3.02	0–14.2
Maturity	Root	0.09 ± 0.12	0–0.34
	Forage	0 ± 0	0–0
	Grain	0 ± 0	0–0
Senescence	Leaf	0.38 ± 0.46	0–1.33
	Root	0.08 ± 0.11	0–0.46
	Whole plant	0 ± 0	0–0

A review of the environmental safety of the PAT protein

Table I.47. Summary of expression levels of PAT protein in grain of *Zea mays* hybrids with line 59122 as detected by ELISA (EFSA, 2008b, 2009a).

	59122 x 1507 x NK603		59122 X NK603		59122	
	Mean	Range	Mean	Range	Mean	Range
PAT (ng/mg Dry Weight)	2.1	1.5–3.1	0.18	0–0.46	0.09	0–0.18

Table I.48. Summary of expression levels of PAT protein in *Zea mays* line DBT418 using Western blot (USDA, 1996d).

Tissue	Genotype	Mean Protein Level (µg/g Dry Weight)							
		V6-V7		Pollen Shed		Dough		Harvest	
		Mean	n ; SE	Mean	n ; SE	Mean	n ; SE	Mean	n ; SE
Leaf	Inbred ¹	351.1	7; 52.91	522.0	6; 59.04	NA	NA	60.8 ⁶	6; 12.46
	Hemizygous hybrid ²	276.3	8; 25.51	501.8	8; 34.75	NA	NA	180.5	8; 24.68
	Homozygous hybrid ³	554.9	2; 136.03	1099.4	3; 76.29	NA	NA	213.6	4; 61.92
Stalk	Inbred	NA	NA	75.8	8; 12.24	NA	NA	95.2	6; 16.86
	Hemizygous hybrid	NA	NA	60.0	8; 11.98	NA	NA	64.4	8; 8.23
	Homozygous hybrid	NA	NA	77.0	4; 11.66	NA	NA	136.3	2; 12.74
Root Ball	Inbred	95.1	7; 16.91	54.1	8; 9.15	NA	NA	24.5	7; 3.71
	Hemizygous hybrid	59.4	8; 3.53	27.5	8; 6.25	NA	NA	21.3	8; 2.23
	Homozygous hybrid	88.1	4; 21.45	69.5	4; 23.58	NA	NA	28.8	3; 7.37
Kernel	Inbred	NA	NA	NA	NA	NA	NA	6.0	6; 1.88
	Hemizygous hybrid	NA	NA	NA	NA	NA	NA	3.1	8; 0.35
	Homozygous hybrid	NA	NA	NA	NA	NA	NA	4.9	4; 0.63
Silk	Inbred	NA	NA	128.2	8; 17.21	NA	NA	NA	NA
	Hemizygous hybrid	NA	NA	29.1	8; 2.97	NA	NA	NA	NA
	Homozygous hybrid	NA	NA	133.3	2; 60.01	NA	NA	NA	NA
Pollen	Hemizygous hybrid ⁴	NA	NA	BLD ⁵	8; NA	NA	NA	NA	NA
	Hemizygous hybrid	NA	NA	BLD	8; NA	NA	NA	NA	NA
	Homozygous hybrid	NA	NA	BLD	4; NA	NA	NA	NA	NA
Whole Plant	Inbred	NA	NA	111.1	8; 16.50	190.5	8; 30.76	NA	NA
	Hemizygous hybrid	NA	NA	72.8	8; 5.88	39.5	7; 7.51	NA	NA
	Homozygous hybrid	NA	NA	119.5	4; 25.63	135.2	3; 10.42	NA	NA

¹ The S4 inbred line (AW/BC2/DBT418 S4) is an unfinished inbred.

² This hybrid (AW/BC2/DBT418.BS/BC1/DBT418) contains two integrated copies of DBT418 insertion.

³ This hybrid (DK.DL (DBT418)) is a “finished hybrid” with one copy of DBT418 coming from an inbred parent line.

⁴ An additional group of the Hemizygous hybrid (AW/BC2/DBT418.BS/BC1/DBT418) was substituted because insufficient pollen was available from the S4 hybrid.

⁵ Below the limit of detection (12.10 µg/g dry weight).

⁶ 2 of 8 samples were below the limit of detection and not used to calculate the mean or standard error.

Table I.49. Summary of the quantification of bar mRNA transcripts in *Zea mays* line MS3 (method not reported) (CFIA, 1996e).

Line	Approximate mRNA (pg/µg total RNA)				
	Leaves	Immature Kernels	Roots	Dry Seeds ²	Germinating Seeds ²
MS3	0.05	0.05	ND ¹	ND	ND

¹ ND = not detected.
² A PAT enzyme activity assay did not detect any PAT in MS3 seeds.

Table I.50. Summary of the PAT protein content *Zea mays* line MS6 as detected by ELISA (USDA APHIS, 1998k).

Tissue	Mg TEP ¹ /g Sample	µg PAT/g Sample	% PAT/TEP
Grain	8.73	3.54	0.04
Forage	1.31	2.01	0.15
Fodder	1.26	2.15	0.17

¹ TEP = Total Extractable Protein.

Table I.51. Summary of PAT protein levels in *Zea mays* hybrid and inbred lines derived from T25 as detected by ELISA (ANZFA, 2001b).

Plant	Mean Levels ± Standard Deviation (ng/mg Protein)			
	Kernel	Silage	Forage	Fodder
Hybrid T25-1	ND ²	14.82 ± 0.86	NA ³	NA
Hybrid T25-2	ND	12.51 ± 1.38	NA	NA
Hybrid T25-3 ¹	ND	14.81 ± 1.30	NA	NA
Inbred T25	4.02 ± 0.62	119.24 ± 13.36	62.70 ± 40.07	79.91 ± 5.23

¹ Plants were treated with phosphinothricin at the V8 stage.

² ND = not detected.

³ NA = data not available.

Table I.52. Summary of PAT protein quantities detected by ELISA in *Zea mays* lines T14 and T25 (USDA, 1994b)¹.

Matrix	% Protein	ng PAT/µg Protein	µg PAT/g Matrix	% PAT in Matrix
T14 silage	0.19	13.03	36.97	3.70
T25 silage	0.05	13.54	6.62	0.67
T14 grain	1.59	0.008	0.115	0.0115

¹ Two extracts from each sample (2 each for silage, 6 for grain) were analyzed in triplicate. Means are reported from all field sites combined.

Table I.53. Summary of PAT protein expression in *Zea mays* line T25 and hybrid crosses with MON810 as detected by ELISA.

Tissue	T25 X MON810		T25	
	Mean	Min-Max	Mean	Min-Max
Leaves	33.3	17.1-54.5	33.9	11.9-64.6

Table I.54. Summary of PAT protein expression as determined by ELISA in *Zea mays* line TC1507(CFIA, 2002b; EFSA, 2005)¹.

Location of Cultivation	Tissue					
	Leaf	Tassel	Silk	Roots	Kernel	Pollen
Canada	<LOD ²	<LOD	<LOD	<LOD	<LOD	NA
Chile	<LOD	<LOD	<LOD	<LOD	<LOD	NA
EU	42 pg/µg (<LOD-136.8 pg/µg) ³	<LOD	<LOD	<LOD	<LOD	NA
United States	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD (<LOD-38.0 pg/µg)

¹ Data presented are from descriptive paragraphs describing different aspects of the same data set. These have been combined for simplicity.

² LOD = limit of detection (7.5 pg/µg total protein for samples from Canada, 20 pg/µg for all other locations).

³ Mean (Range).

A review of the environmental safety of the PAT protein

Table I.55. Summary of PAT protein expression as determined by ELISA in *Zea mays* line TC1507 (USDA, 2001c)¹.

Leaf	Pollen	Silk	Stalk	Whole Plant	Grain	Senescent Whole Plant
<LOD2 (<LOD-40.8)	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD

¹ All values are pg/μg protein. Mean values are listed with observed ranges in parentheses.

² LOD = limit of detection (20 pg/μg total protein).

Table I.56. Summary of PAT protein expression in plants derived from *Zea mays* line TC1507 and DAS59122 as detected by ELISA (EFSA, 2009b)¹.

	1507 x5912		1507		59122	
Grain	Mean	Range	Mean	Range	Mean	Range
	0.15	0.09–0.27	<LOD ²	<LOD	0.09	0–0.18
Forage	2.53	1.01–3.97	NA	NA	NA	NA

¹ Values are ng/mg dry weight.

² LOD = limit of detection.