
INSTRUCTION TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
Keppel Street, London, WC1E 7HT,
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5×18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 13.5×18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and Physiology of parasites* (ed A. M. Fallis) pp. 213 – 222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49 – 59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

| | Pages |
|--|---------|
| The serodiagnosis of human hydatid disease: 1. The routine use of latex-agglutination and complement-fixation in diagnosis. M. W. DIGHERO and C. M. P. BRADSTREET | 283-286 |
| The serodiagnosis of human hydatid disease: 2. Additional studies on selected sera using indirect haemagglutination (IHA), enzyme linked immunosorbent assay (ELISA) and defined antigen substrate spheres (DASS). R. M. MATOSSIAN <i>et al.</i> | 287-291 |
| Antigenic components of <i>Taenia saginata</i> and their relevance to the diagnosis of bovine cysticercosis by immunoelectrophoresis. S. GEERTS, V. KUMAR and N. AERTS | 293-299 |
| <i>Moaciria chondropythonis</i> sp. n. (Heterakidae: Spinicaudinae), a nematode from a Papuan tree python, <i>Chondropython viridis</i> . L. M. GIBBONS | 301-306 |
| The morphology and biology of <i>Xenopodistomum xenopodis</i> from the gall bladder of the African clawed toad, <i>Xenopus laevis</i> . R. C. TINSLEY and R. W. OWEN | 307-316 |
| <i>Goezia lacerticola</i> sp. n. (Nematoda: Anisakidae) in <i>Alligator mississippiensis</i> from Florida. T. L. DEARDORFF and R. M. OVERSTREET | 317-320 |
| Studies of the relationship between <i>Schistosoma</i> and their intermediate hosts. III. The genus <i>Biomphalaria</i> and <i>Schistosoma mansoni</i> from Egypt, Kenya, Sudan, Uganda, West Indies (St. Lucia) and Zaire (two different strains: Katanga and Kinshasa). F. FRANSEN | 321-348 |
| Studies of the relationship between <i>Schistosoma</i> and their intermediate hosts. IV. The genus <i>Bulinus</i> and <i>Schistosoma bovis</i> from Morocco. F. FRANSEN | 349-355 |
| <i>Asymphylogora fishelsoni</i> sp. n. (Trematoda: Monorchidae) from a cyprinodontid fish, <i>Aphanius dispar richardsoni</i> (Boulenger), from Israel, and its allometric growth. J. M. FISCHTHAL | 357-362 |
| Species of <i>Cloacina</i> Linstow, 1898 (Nematoda: Strongyloidea) from the Blacktailed Wallaby, <i>Wallabia bicolor</i> Desmarest, 1804) from eastern Australia. I. BEVERIDGE | 363-378 |