

Supernova interaction with dense mass loss

Roger A. Chevalier

Dept. of Astronomy, University of Virginia
P.O. Box 400325, Charlottesville, VA 22903, USA
email: rac5x@virginia.edu

Abstract. Supernovae of Type II_n (narrow line) appear to be explosions that had strong mass loss before the event, so that the optical luminosity is powered by the circumstellar interaction. If the mass loss region has an optical depth $> c/v_s$, where v_s is the shock velocity, the shock breakout occurs in the mass loss region and a significant fraction of the explosion energy can be radiated. The emission from the superluminous SN 2006gy and the normal luminosity SN 2011ht can plausibly be attributed to shock breakout in a wind, with SN 2011ht being a low energy event. Superluminous supernovae of Type I may derive their luminosity from interaction with a mass loss region of limited extent. However, the distinctive temperature increase to maximum luminosity has not been clearly observed in Type I events. Suggested mechanisms for the strong mass loss include pulsational pair instability, gravity-waves generated by instabilities in late burning phases, and binary effects.

Keywords. (stars:) supernovae: general

1. Introduction

Once the radiation from the supernova shock wave emerges from the stellar surface, the interaction with the surrounding medium commences. In the case of core collapse supernovae, the initial interaction is with mass loss from the progenitor star. At low circumstellar densities, the emission from the interaction is a small part of the supernova power and typically involves X-ray and radio radiation. This is the case for Type IIP supernovae, where the inferred circumstellar density is roughly consistent with that expected from the red supergiant progenitors of these events (Chevalier *et al.* 2006). However, there are supernovae, of Type II_n, that are thought to have most of their peak optical power come from interaction with a circumstellar medium (e.g., Chugai & Danziger 1994). Here I discuss cases where the circumstellar interaction is an important part of the optical emission. The shock breakout phase is treated in Section 2 and the initial viscous shock propagation in Section 3. Speculations on the formation of the dense circumstellar medium are discussed in Section 4 and the final comments in Section 5.

2. Shock Breakout

A useful way of considering the enhanced supernova luminosity that can result from very dense circumstellar interaction is in terms shock breakout in the dense gas (Chevalier & Irwin 2011, Balberg & Loeb 2011). While a shock front is inside the supernova progenitor, the shock front is mediated by radiation for standard supernova conditions. The optical depth across the shock transition for a radiation dominated shock is $\sim c/v_s$, where v_s is the shock velocity and c is the speed of light. This result is determined by the condition that the hydrodynamic time for the shock wave to cross the shock thickness is approximately equal to the diffusion time for the radiation. When the shock front reaches a position that is at an optical depth of $\sim c/v_s$ from the surface, radiation can diffuse

out to optically thin layers and shock breakout occurs. For low circumstellar densities, the timescale for the breakout event is typically quite brief and depends on the radius of the progenitor star, R . For a red supergiant progenitor, the most extended normal progenitor, the supernova shock takes about a day to traverse the star and the breakout phase lasts about half an hour, determined by the light travel time across the star, R/c .

A low density, optically thin circumstellar medium is not expected to have much influence on the optical light near maximum, although it can be very important for radio and X-ray emission. At higher circumstellar densities, the medium becomes optically thick and the breakout radiation must diffuse out through the circumstellar medium. The result is that the radiated energy in the breakout is not changed, but the radiation is emitted over a longer period of time, lowering the luminosity.

When the circumstellar optical depth, τ_w , becomes of order c/v_s or more, the radiation dominated shock can propagate into the circumstellar region. In this case, substantial supernova energy can be present as internal energy at a larger radius than would otherwise be the case, so that the effects of adiabatic expansion are lessened and the luminosity rises. Chevalier & Irwin (2011) suggested that this model can describe the initial rise of 10's of days of superluminous supernovae like SN 2006gy. When the circumstellar interaction occurs at a large radius where diffusion of radiation can occur rapidly and the circumstellar medium is sufficiently massive to thermalize the supernova kinetic energy, a large fraction of the supernova energy can be radiated. The physical parameters, a circumstellar medium of $\sim 10 M_\odot$ extending out to $\sim 10^{16}$ cm, are similar to those in the diffusion model of Smith & McCray (2007). Chevalier & Irwin (2011) used an analytic model for the emission, which has limitations; in particular, the initial interaction was described by a self-similar solution, but that solution breaks down as the reverse shock wave propagates into the supernova ejecta. Ginzburg & Balberg (2012) undertook numerical simulations that did not have this problem and obtained comparable parameters for SN 2006gy. In their simulations, Moriya *et al.* (2013) found that a density law $\rho \propto r^{-2}$ did not give a good fit to light curve; their favored explosion parameters are mass loss as high as $15 M_\odot$ and energy no more than 4×10^{51} ergs.

A general expectation of the shock breakout model is that the initial rise to maximum luminosity should be primarily due to heating of the photosphere as the shock wave begins to affect the stellar emission. This effect is not clearly observed in the data on SN 2006gy, although there is some sign of an increasing temperature in the spectra of Smith *et al.* (2010).

A Type II_n event that apparently does show the increasing temperature in the rise to maximum is SN 2011ht. Early Swift observations show a clear rise in temperature to maximum (Roming *et al.* 2012), which is characteristic of shock breakout in the circumstellar gas. The rise time to maximum, 40 days, is an indicator of the density of the circumstellar gas. The supernova radiation is expected to be efficiently radiated, so that the radiated energy is a significant fraction of the supernova energy if the energy is thermalized by the interaction. Interestingly, the light curve is similar in shape to that of SN 2006gy, but the absolute magnitude is fainter by 5 magnitudes, with a radiated energy $\sim 2.5 \times 10^{49}$ ergs. The implication is that the explosion energy is smaller by a factor up to $\sim 10^2$. The possibility that the event was not a supernova was noted by Roming *et al.* (2012). However, there are supernovae that occur in this low energy range and Mauerhan *et al.* (2013) suggest that the late emission from SN 2011ht is due to a small amount of ^{56}Ni . In the end, it is not possible to definitively answer the question of whether SN 2011ht was a supernova.

For luminous supernovae that are of Type II_n, it is very plausible that the high optical luminosity is produced by circumstellar interaction (see next section). There are also

luminous supernovae that are not of the “n” type. SN 2008es is a luminous Type II (Gezari *et al.* 2009, Miller *et al.* 2009) and SN 2010gx is an example of a luminous Type Ib/c (Pastorello *et al.* 2010). The Type Ib/c objects outnumber those of Type II. Chevalier & Irwin (2011) suggested that the optical light from these objects is powered by circumstellar interaction in which the mass loss region ends before the diffusion radius is reached by the shock front. Ginzburg & Balberg (2012) carried out numerical simulations of this situation and successfully modeled the optical light curve from SN 2010gx. As described above, an expectation in this model is that the temperature should increase in the rise to maximum luminosity. In the case of SN 2006oz, there are observations on the rise to maximum and there is no evidence for an increasing temperature; the temperature remains roughly constant (Leloudas *et al.* 2012). Another mechanism, such as magnetar power (Kasen & Bildsten 2010), may be indicated.

Another superluminous event of interest is PS1-10afx (Chornock *et al.* 2013). In this case, the luminosity and temperature at maximum implied a photospheric radius of 5×10^{15} cm, while the observed lines indicated an expansion velocity of 11,000 km s⁻¹. The implied expansion timescale is 50 days, but the rise time for the light curve is $\sim 10 - 15$ days. This evolution is naturally explained in the shock breakout view (Chevalier & Irwin 2011); the shock front must traverse the optically thick star before shock breakout occurs. However, the rise to luminosity maximum did not show the increasing temperature that might be expected in the breakout scenario and there is no spectral evidence for circumstellar interaction, so the nature of the event is still in doubt.

Overall, the shock breakout scenario is attractive for the events that show narrow line spectra because there is evidence for slowly expanding circumstellar matter ahead of the shock front. When these lines are not present, there is no clear evidence for dense circumstellar matter. There may be more than one mechanism that gives rise to very luminous supernovae.

3. Viscous Shock Wave

Once the forward shock wave gets into a regime where the optical depth to the surface $\tau < c/v_s$, a radiation dominated shock can no longer be maintained and there is a transition to a viscous shock front. There may initially be radiative acceleration of the unshocked gas, so that a viscous shock does not initially form, but, if the circumstellar medium is extended, the formation of a viscous shock is inevitable. The maximum optical depth at which it can form is somewhat less than c/v_s .

At low circumstellar densities the emission from the reverse shock dominates that from the forward shock because the density is higher and the shock velocity is lower, ~ 1000 km s⁻¹ or less. However, as the circumstellar density rises, the reverse shock becomes radiative first, which leads to a lower increase in luminosity with increasing density and to a dense shell that can absorb radiation from the reverse shock, thus affecting the X-ray luminosity.

The forward shock, with a velocity of 1000's of km s⁻¹ or more, can heat gas to a temperature of $100v_4^2$ KeV, where v_4^2 is the shock velocity in units of 10^4 km s⁻¹. Katz *et al.* (2011) found that inverse Compton cooling can balance the shock heating at a temperature of 60 keV if the optical depth is c/v_s (breakout conditions) and the shock velocity is 10^4 km s⁻¹. This estimate assumed that the electrons are heated only by Coulomb collisions with the ions. If there is additional collisionless heating, the gas temperature is raised. Chevalier & Irwin (2012) estimated regions of the shock velocity vs circumstellar density plane where the cooling time is short compared to the age and where inverse Compton cooling is larger than bremsstrahlung cooling. The regime where

the cooling is rapid roughly overlaps the regime where the electron scattering optical depth is >1 . If inverse Compton losses are larger than bremsstrahlung, the X-ray emission from the hot shocked gas is reduced as a fraction of the shock power.

The X-rays that are emitted by the hot gas have to propagate through the preshock circumstellar medium. One effect of the surrounding medium is Compton recoil (Chevalier & Irwin 2012, Svirski *et al.* 2012), which reduces the energies of the high energy photons to a maximum of $\sim 511/\tau_{es}^2$ keV, where τ_{es} is the electron scattering optical depth and is assumed >1 . It can be seen that $\tau_{es} > 8$ is necessary to have an effect in the energy range around 10 keV that is accessible to X-ray telescopes like *Chandra*. *NuSTAR*, with its sensitivity to 80 keV, has a better chance of detecting this effect.

Another effect is photoabsorption by the preshock medium (Chevalier & Irwin 2012), which has a larger effect at low X-ray energies. An important aspect of photoabsorption is that its occurrence requires that the absorbing medium not be completely ionized. For the X-ray emission that is expected from the hot gas, there is the possibility of complete ionization (Chevalier & Irwin 2012). Complete ionization is expected at higher shock velocities $\sim 10^4$ km s $^{-1}$, but not at lower velocities, <5000 km s $^{-1}$. The result also depends on clumping and asymmetries in the circumstellar gas. If the gas is not fully ionized, the column density corresponding to an electron scattering optical depth τ_{es} is $3 \times 10^{23} \tau_{es} (Z/Z_{\odot})^{-1}$ cm $^{-2}$, where Z is the supernova metallicity and Z_{\odot} is the solar metallicity. Even for $\tau_{es} = 1$, the column density is orders of magnitude larger than a typical interstellar column density and would have a dramatic effect on X-ray emission.

A supernova that provides some test of these models is the Type II_n (narrow line) SN 2010jl. The optical spectrum showed narrow, presumably circumstellar, emission lines on top of broad wings that could be attributed to electron scattering (Smith *et al.* 2010). The scattering requires an electron scattering optical depth ≥ 1 . An X-ray spectrum with the *Chandra* observatory at 2 months required a column density $\sim 10^{24}$ cm $^{-2}$, (Chandra *et al.* 2012). At an age of 12 months, the column dropped to $\sim 3 \times 10^{23}$ cm $^{-2}$, confirming that the absorption is connected to circumstellar gas. The observations with *Chandra* showed a hot thermal component with temperature ≥ 10 keV, corresponding to a shock velocity ≥ 3000 km s $^{-1}$. The emission is presumably from the forward shock front in SN 2010jl.

The presence of the large column density in SN 2010jl shows that it is possible for a large fraction of the X-ray emission to be extinguished. The absorption in SN 2010jl corresponds to $\tau_{es} \sim 1$; larger values of τ_{es} are expected close to the time of shock breakout in a wind. The expected absorption optical depth at 10 keV is $\sim 1.5\tau_{es} (Z/Z_{\odot})$ if the circumstellar region is not fully ionized. Chevalier & Irwin (2012) noted that the observed low X-ray luminosity of SN 2006gy, $\leq 10^{40}$ ergs s $^{-1}$ (Smith *et al.* 2007; Ofek *et al.* 2007) compared to an optical luminosity of 3×10^{44} ergs s $^{-1}$ at a time close to maximum light (Smith *et al.* 2010), is consistent with breakout in a wind. The case of SN 2011ht was also mentioned above as a likely case of a wind breakout. An X-ray luminosity of 10^{39} ergs s $^{-1}$ was initially reported for this object based on *Swift* observations (Roming *et al.* 2012), but higher spatial observations with *Chandra* showed that the source is not coincident with SN 2011ht (Pooley 2012). Taking a conservative upper limit to the X-ray luminosity of 10^{39} ergs s $^{-1}$, the X-ray emission is again a small fraction of the optical luminosity, which was 3×10^{42} ergs s $^{-1}$. The strong suppression of the X-ray emission in SN 2006gy and SN 2011ht is consistent with both of these objects being shock breakouts in a wind, although the explosion energy is probably much smaller in SN 2011ht.

Dwarkadas & Gruszko (2012) make the point that the evolution of X-ray emission from supernovae is typically not what is expected for a steady wind, so that estimates of mass loss rate can be misleading. However, the preshock density at any time can be

estimated from $\rho_0 = L/(2\pi\eta R^2 v_s^3)$, where η is the efficiency of conversion of the shock power into the luminosity L . If observations are made over a period of time, an estimate of the mass required for the luminosity can be made and if the velocity of the mass loss can be estimated from narrow lines, the timescale for the emission can be obtained. The mass and timescale estimates provide useful constraints on possible mechanisms for the mass loss.

An interesting aspect of the viscous shock front in dense mass loss is that the shock wave is expected to be collisionless, so that diffusive shock acceleration of particles to relativistic energies can occur (Katz *et al.* 2011, Murase *et al.* 2011). These studies show that an unusually nearby dense interaction supernova would have to occur to have a chance of detecting high energy gamma-ray emission from the supernova.

4. Origin of the Dense Mass Loss

Many of the characteristics of Type II_n supernovae require an optical depth to electron scattering ≥ 1 in the circumstellar medium, which can be expected to produce wings to narrow line profiles and obscure the inner high velocities related to the supernova (if present). Typical parameters for Type II_n events involve masses of $0.01 - 10 M_\odot$ on scales of $10^{15} - 10^{16}$ cm from the supernova. As noted above, a circumstellar mass as high as $10 - 20 M_\odot$ has been estimated for SN 2006gy. These events have been linked to luminous blue variables (LBVs) because they are some of the only objects known to have such extreme mass loss (Kiewe *et al.* 2012). In addition, the Type II_n SN 2005gl was found to have a probable LBV progenitor in pre-explosion images (Gal-Yam & Leonard 2009). Although this link has been made, there is still the issue that the reason for the extreme mass loss from LBVs is not understood. In addition, stellar evolution models do not predict supernova explosions at the time of the LBV phase. The timing of the explosion close in time to the mass loss is a remarkable feature of the Type II_n events.

One explanation for SN 2006gy was pulsational pair instability of massive stars (Woosley *et al.* 2007). In this model, there is no terminal explosion, but the emission results from the interaction of pulsationally driven shells. The shells can have a mass of several M_\odot and the radiated energy from their interaction can be 10^{50} ergs.

An interesting proposal by Quataert & Shiode (2012) is that super-Eddington fusion reactions at the end of the life of a massive star can generate gravity waves that deposit their energy/momentum in the outer parts of the star, driving strong mass loss. The prime burning phases for this are Ne and O burning, which occur in the last year of evolution before core collapse. However, only a fraction of massive stars show evidence for the late mass loss and, in this hypothesis, it is unclear what determines the fraction that show the mass loss. A variant of this hypothesis is that the burning instability leads to expansion of the star instead of mass loss, and the mass loss is driven by binary interaction (Soker 2013).

Ofek *et al.* (2013) found emission from SN 2010mc in the 40 days leading up to the explosion and suggested that the observations support the gravity wave driven mass loss hypothesis. The reasons are that the circumstellar velocities are relatively high, ~ 2000 km s⁻¹ and that the brightening was observed 40 days before explosion, when the late unstable burning phases are expected. However, Smith *et al.* (2013) show that the light curve of SN 2010mc is very much like that of SN 2009ip during the period June - Sept 2012, indicating that they involve a similar physical situation. SN 2009ip showed eruptive behavior over the period 2009 to 2012, somewhat longer than might be expected for the Ne and O burning phases. Also, high velocities observed in spectral lines are possibly the result of electron scattering. In any case, Fraser *et al.* (2013) note that there is some

question whether SN 2009ip was in fact a supernova in Sept 2012, a question that can also be raised for SN 2010mc.

Another possibility for the strong mass loss is that it is driven by close binary interaction. Common envelope evolution is expected in close binaries, accompanied by strong mass loss. This has the advantage that there can be considerable variety in the events. Also, recent observations have shown that massive stars are in close binaries more frequently than had been thought (Sana *et al.* 2012). The problem is to have an explosion close to the time that the mass loss occurs. Chevalier (2012) speculated that the event involves the spiral in of a neutron star due to common envelope evolution. In cases where the interaction occurs at a relatively small separation, the neutron star spirals into the core of its companion because of the steep density gradient. In cases where there is spiral in, but the separation is somewhat greater, the companion star can become a red supergiant with a flat density profile, which causes spiral in to stop. The result is a neutron star – helium star binary. When the He star evolves, there is again the possibility of mass loss accompanied by the spiral in of the neutron star to the core.

In one view, the spiral in of the neutron star to the core gives rise to a red supergiant with a neutron star core, a Thorne-Żytkow star (Thorne & Żytkow 1977). However, neutrino losses of the matter near the neutron star may lead to strong accretion onto the neutron star, especially when the neutron star is in the core of the companion star (Chevalier 1996). The rapid accretion may lead to black hole formation and an explosion, as in the model for gamma-ray bursts (GRBs) of Fryer & Woosley (1998). A variant of this model is that the rapid accretion is accompanied by strong magnetic field amplification of the neutron star, leading to an explosion (Barkov & Komissarov 2011).

In this scenario for explosions preceded by strong mass loss, the explosion mechanism is not the neutrino mechanism, which is generally favored for most core collapse supernovae, but is related to rapid rotation and accretion on a central compact object. Spiral in of a compact object is an efficient way of achieving rapid rotation in the central region. The mechanism is believed to operate for GRBs, which have been associated with energetic ($>10^{52}$ ergs) supernovae. Some Type IIn supernovae also appear to have had unusually large energies; for example, SN 2003ma had an integrated bolometric luminosity of 4×10^{51} ergs over 4.7 years and an estimated explosion energy $> 10^{52}$ ergs (Rest *et al.* 2011). Studies of the neutrino mechanism for core collapse supernova explosions indicate that the maximum energy that can be attained is $\sim 2 \times 10^{51}$ ergs (Janka 2012). Both GRBs and some Type IIn supernovae may require an explosion mechanism involving rotation and magnetic fields in order to produce high energy explosions.

5. Discussion

Although circumstellar interaction at low densities can be successfully described in terms of spherical models, there are signs that at the higher densities considered here, where the interaction contributes to the optical luminosity, the situation is more complex. Galactic LBVs show a complex circumstellar structure, binary evolution can lead to aspherical structure (as apparently occurred in SN 1987A), and Type IIn supernovae often show significant polarization in their optical light. In addition to these complexities, the Type IIn events appear to include a wide range of stellar masses and explosion energies, indicating a variety of evolutionary paths. We are still at the early stages of understanding these events.

Acknowledgement

I am grateful to my collaborators for their participation in this research, and to NASA grant NNX12AF90G and NSF grant AST-0807727 for support.

References

- Balberg, S. & Loeb, A. 2011, *MNRAS*, 414, 1715
- Barkov, M. V. & Komissarov, S. S. 2011, *MNRAS*, 415, 944
- Chandra, P., Chevalier, R. A., Irwin, C. M., *et al.* 2012, *ApJ(Letters)*, 750, L2
- Chevalier, R. A. 1996, *ApJ*, 459, 322
- Chevalier, R. A. 2012, *ApJ(Letters)*, 752, L2
- Chevalier, R. A. & Irwin, C. M. 2011, *ApJ(Letters)*, 729, L6
- Chevalier, R. A. & Irwin, C. M. 2012, *ApJ(Letters)*, 747, L17
- Chevalier, R. A., Fransson, C., & Nymark, T. K. 2006, *ApJ*, 641, 1029
- Chornock, R., Berger, E., Rest, A., *et al.* 2013, preprint, arXiv:1302.0009
- Chugai, N. N. & Danziger, I. J. 1994, *MNRAS*, 268, 173
- Dwarkadas, V. V. & Gruszko, J. 2012, *MNRAS*, 419, 1515
- Fraser, M., Inserra, C., Jerkstrand, A., *et al.* 2013, *MNRAS*, submitted, arXiv:1303.3453
- Fryer, C. L. & Woosley, S. E. 1998, *ApJ(Letters)*, 502, L9
- Gal-Yam, A. & Leonard, D. C. 2009, *Nature*, 458, 865
- Gezari, S., Halpern, J. P., Grupe, D., *et al.* 2009, *ApJ*, 690, 1313
- Ginzburg, S. & Balberg, S. 2012, *ApJ*, 757, 178
- Janka, H.-T. 2012, *ARNPS*, 62, 407
- Kasen, D. & Bildsten, L. 2010, *ApJ*, 717, 245
- Katz, B., Sapir, N., & Waxman, E. 2011, preprint, arXiv:1106.1898
- Kiewe, M., Gal-Yam, A., Arcavi, I., *et al.* 2012, *ApJ*, 744, 10
- Leloudas, G., Chatzopoulos, E., Dilday, B., *et al.* 2012, *A&A*, 541, A129
- Mauerhan, J. C., Smith, N., Silverman, J. M., *et al.* 2013, *MNRAS*, 1053
- Miller, A. A., Chornock, R., Perley, D. A., *et al.* 2009, *ApJ*, 690, 1303
- Moriya, T. J., Blinnikov, S. I., Tominaga, N., *et al.* 2013, *MNRAS*, 428, 1020
- Murase, K., Thompson, T. A., Lacki, B. C., & Beacom, J. F. 2011, *Phys. Rev. D*, 84, 043003
- Ofek, E. O., Cameron, P. B., Kasliwal, M. M., *et al.* 2013, *ApJ(Letters)*, 659, L13
- Ofek, E. O., Sullivan, M., Cenko, S. B., *et al.* 2013, *Nature*, 494, 65
- Pastorello, A., Smartt, S. J., Botticella, M. T., *et al.* 2010, *ApJ(Letters)*, 724, L16
- Pooley, D. 2012, *The Astro. Tel.*, 4062, 1
- Quataert, E. & Shiode, J. 2012, *MNRAS*, 423, L92
- Rest, A., Foley, R. J., Gezari, S., *et al.* 2011, *ApJ*, 729, 88
- Roming, P. W. A., Pritchard, T. A., Prieto, J. L., *et al.* 2012, *ApJ*, 751, 92
- Sana, H., de Mink, S. E., de Koter, A., *et al.* 2012, *Science*, 337, 444
- Smith, N. & McCray, R. 2007, *ApJ(Letters)*, 671, L17
- Smith, N., Li, W., Foley, R. J., *et al.* 2007, *ApJ*, 666, 1116
- Smith, N., Chornock, R., Silverman, J. M., Filippenko, A. V., & Foley, R. J. 2010, *ApJ*, 709, 856
- Smith, N., Mauerhan, J. C., Kasliwal, M. M., & Burgasser, A. J. 2013, preprint, arXiv:1303.0304
- Soker, N. 2013, preprint, arXiv:1302.5037
- Svirski, G., Nakar, E., & Sari, R. 2012, *ApJ*, 759, 108
- &Thorne, K. S., Żytkow, A. N. 1977, *ApJ*, 212, 832
- Woosley, S. E., Blinnikov, S., & Heger, A. 2007, *Nature*, 450, 390

Discussion

SUTARIA: Concerning SN2011ht there is a controversy about the nature of this event; is it a true SN II_n or is it a SN imposter. Could you please comment?

CHEVALIER: In the shock breakout view of this event, the circumstellar density is high, but the shock velocity, and thus the energy, are low. The model does not clearly answer the SN vs. imposter question.

NOMOTO: What is your model for Type I Superluminous Supernovae?

CHEVALIER: In the common envelope (CE) view, there is an initial CE episode with the neutron star spiraling in the 1f envelope. Depending on parameters, the outcome may be a neutron star He star binary. Eventually the He star expands and there is another spiraling phase for the compact object. This can lead to a Type I SLSN.

SURNIS: What are the chances of detecting the central compact object in X-rays and radio for Galactic SNRs having dense CSM? What if the optical density of the CSM is low?

CHEVALIER: The dense surroundings might make it more difficult to see a compact object because of obscuration, or continuing interaction could compete with central emission. The effect of the region immediately around the center depends on the supernova properties, which are not well understood.