

---

## INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology  
London School of Hygiene and Tropical Medicine  
Keppel Street, London WC1E 7HT  
England

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5 × 18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints, preferably of the same size as they are to appear in the Journal (maximum size 13.5 × 18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and physiology of parasites* (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

---

# Contents

	Pages
Studies on humoral immunity against <i>Taenia taeniaeformis</i> infection in rats. B. H. KWA and F. Y. LIEW	99-107
The distribution of <i>Ancylostoma caninum</i> larvae in the central nervous system of the mouse infected with single or repeated doses. M. K. BHOPALE and G. N. JOHRI	109-113
The influence of physical factors on the behaviour and infectivity of miracidia of <i>Schistosoma mansoni</i> and <i>S. haematobium</i> . 11. Effect of light and depth. S. K. PRAH and C. JAMES	115-120
Studies with <i>Brugia pahangi</i> . 15. Cobalt 60 irradiation of the worm. P. OOTHUMAN, D. A. DENHAM, P. B. MCGREEVY and G. S. NELSON	121-126
Studies with <i>Brugia pahangi</i> . 16. Precipitation antibody in infected cats detected by counterimmunoelectrophoresis. R. A. DESOWITZ, P. OOTHUMAN and D. A. DENHAM	127-130
Two new species of <i>Enenterum</i> Linton, 1910 (Digenea) in the marine fish <i>Neoscorpis lithophilus</i> (Kyphosidae) from the south-western Indian Ocean. R. A. BRAY	131-139
<i>Diocetowittus</i> spp. Chabaud and Le Van Hoa 1960 (Nematoda: Adenophorea) from Australian snakes. H. I. JONES	141-145
<i>Polyonchobothrium scleropagis</i> n. sp. (Cestoda: Pseudophyllidea) from the Australian osteoglossid fish <i>Scleropages leichardti</i> Günther. D. BLAIR	147-153
<i>Phaneropsolus aspinosus</i> sp. n. (Lecithodendriidae: Phaneropsolinae) from the leaf monkey <i>Macaca fascicularis</i> (Raffles). J. R. PALMIERI and M. KRISHNASAMY	155-158
Histochemical distribution of succinic dehydrogenase in the lymphatic system of a trematode <i>Ceylonocotyle scoliocoelium</i> . P. N. SHARMA	159-162
SUPPLEMENTARY REVIEW ARTICLE	
Ascaridoid nematodes of amphibia and reptiles: <i>Paraheterotyphlum</i> . J. F. A. SPRENT	163-170