

project, creating a community capable of acting at the intersection of policymaker, companies and citizens.

Conclusion: The case of NOFEAR demonstrates how, thanks to a European project, created a network of individuals and businesses that interact and or collaborate with each other can accelerate knowledge driven and sustainable growth of multidisciplinary ecosystems able to mitigate the fragmentation of the emergency medical systems.

Prehosp. Disaster Med. 2023;38(Suppl. S1):s192–s193

doi:10.1017/S1049023X2300496X

Implementing Crisis Standards of Care in the Intensive Care Unit: A Scoping Review

David Sellers^{1,2}, Julia Crilly PhD^{1,3}, Lynda Hughes PhD², Jamie Ranse PhD^{1,3}

1. Menzies Health Institute Griffith University, Gold Coast, Australia
2. School of Nursing and Midwifery, Griffith University, Gold Coast, Australia
3. Department of Emergency Medicine, Gold Coast Health, Gold Coast, Australia

Introduction: Disasters have the potential to cause a surge of patients, some of which may require admission to an intensive care unit (ICU). Due to the high resource requirements of ICUs, normal standards of care may need to be altered to treat more people with limited resources, a care model referred to as crisis standards of care (CSC). The pragmatic implementation of CSC in ICUs due to patient surges from disasters has not been well explored in the literature.

Method: This scoping review guided by the Joanna Briggs institute methodology for scoping reviews searched medical databases including CINAHL, PubMed, ProQuest and SCOPUS. Articles were included if they reflected on the actual implementation of CSC delivered in ICU as a result of a patient surge from a disaster. Quantitative data was extracted into tables and qualitative content was thematically analyzed.

Results: A total of 17 papers were included in the review. The disaster event that dominated the results was COVID-19. Most papers relayed subjective accounts of how care models were impacted by patient surges. Common themes included the repurposing of other clinical areas to accommodate ICU patients, resource shortages (particularly ventilators) and staff shortages. Moral strain was felt when processes such as palliation and treatment modality were altered due to resource restrictions.

Conclusion: This review highlights the dearth of high-quality research in implementing CSC in ICUs. Understanding the pragmatic experiences of CSC shows not only the logistical insufficiencies that have been experienced, but the moral and clinical repercussions that these insufficiencies have caused. Inadequate preparation for future disasters, particularly short notice disasters, may lead to further implementation of CSC resulting in poorer outcomes for patients and detrimental impacts on healthcare workers. More research into the practical

application of CSC in ICU may help mitigate the impact of patient surges from disasters.

Prehosp. Disaster Med. 2023;38(Suppl. S1):s193

doi:10.1017/S1049023X23004971

Women are More Infected and Seek Care Faster but are Less Severely Ill: Gender Gaps in Covid-19 Morbidity and Mortality During Two Years of a Pandemic in Israel

Arielle Kaim MPH^{1,2}, Mor Saban PhD^{1,3}

1. Gertner Institute, Tel Aviv, Israel
2. Tel Aviv University, Sackler School of Medicine, Department of Emergency and Disaster Management, Ramat Gan, Israel
3. Tel Aviv University, School of Health Professions, Ramat Gan, Israel

Introduction: In the context of COVID-19 outcomes, global data has deduced a gender bias towards severe disease among males. The aim of this study is to compare morbidity and mortality during two years of the COVID-19 pandemic in female and male patients with COVID-19, as well as to assess length of stay, health seeking behavior time after positive diagnosis, and vaccination differences.

Method: A retrospective-archive study was conducted in Israel from March 1st (patient zero cases) to March 1st, 2022 (two consecutive years). Data were obtained from the Israeli Ministry of Health's (MOH) open COVID-19 database.

Results: The findings indicate female infections are 1.12 times more likely, across almost all age groups, apart from the youngest (0-19) age groups. Despite this, the relative risk of severe illness, intubation and mortality is higher among men. In addition, our findings indicate that the mean number of days taken by unvaccinated men from positive diagnosis to hospital admission was greater than among unvaccinated women among the deceased population.

Conclusion: Targeted approaches including risk communication which take into consideration sex and gender and the intersecting factors are necessary to engage in the fight against COVID-19 for ensuring the most effective and equitable pandemic response.

Prehosp. Disaster Med. 2023;38(Suppl. S1):s193

doi:10.1017/S1049023X23004983

Emergency Healthcare Providers' Perceptions of Preparedness and Willingness to Work during Disasters and Public Health Emergencies

Mohammed Sultan PhD(c)^{1,2}, Jarle Lørve Sørensen PhD³, Eric Carlström PhD¹, Luc Mortelmans PhD⁴, Amir Kborram-Manesh PhD⁵

1. Institute of Health and Care Sciences, Sahlgrenska Academy, Gothenburg University, Gothenburg, Sweden
2. Healthcare Transformation, Model of Care, Regional Health Directorate, Najran, Saudi Arabia
3. USN School of Business, Campus Vestfold, University of South-Eastern Norway, Kongsberg, Norway