

## ON SOME DIMENSION FORMULA FOR AUTOMORPHIC FORMS OF WEIGHT ONE, II

TOYOKAZU HIRAMATSU

### §0. Introduction

Let  $\Gamma$  be a fuchsian group of the first kind and assume that  $\Gamma$  contains the element  $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$  ( $= -I$ ), and let  $\chi$  be a unitary representation of  $\Gamma$  of degree 1 such that  $\chi(-I) = -1$ . Let  $S_1(\Gamma, \chi)$  be the linear space of cusp forms of weight one on the group  $\Gamma$  with character  $\chi$ . We shall denote by  $d_1$  the dimension of the linear space  $S_1(\Gamma, \chi)$ . It is not effective to compute the number  $d_1$  by means of the Riemann-Roch theorem. Because of this reason, it is an interesting problem in its own right to determine the number  $d_1$  by some other method (for example, [5]).

When the group  $\Gamma$  has a compact fundamental domain in the upper half plane  $S^1$ , we have obtained the following dimension formula which is a slightly modified form of the previous result ([1]):

$$(1) \quad d_1 = \frac{1}{2} \sum_{\{M\}} \frac{\chi(M)}{[\Gamma(M): \pm I]} \frac{\bar{\zeta}}{1 - \bar{\zeta}^2} + \frac{1}{2} \operatorname{Res}_{s=0} \zeta^*(s),^{2)}$$

where the sum over  $\{M\}$  is taken over the distinct elliptic conjugacy classes of  $\Gamma/\{\pm I\}$ ,  $\Gamma(M)$  denotes the centralizer of  $M$  in  $\Gamma$ ,  $\bar{\zeta}$  is one of the eigenvalues of  $M$ , and  $\zeta^*(s)$  denotes the Selberg type zeta-function defined by

$$\zeta^*(s) = \sum_{\alpha=1}^{\infty} \sum_{k=1}^{\infty} \frac{\chi(P_\alpha)^k \log \lambda_{0,\alpha}}{\lambda_{0,\alpha}^k - \lambda_{0,\alpha}^{-k}} |\lambda_{0,\alpha}^k + \lambda_{0,\alpha}^{-k}|^{-s}.$$

Here,  $\lambda_{0,\alpha}$  denotes the eigenvalue ( $\lambda_{0,\alpha} > 1$ ) of a representative  $P_\alpha$  of the primitive hyperbolic conjugacy classes  $\{P_\alpha\}$  in  $\Gamma/\{\pm I\}$ .

---

Received November 9, 1985.

Revised September 12, 1986.

<sup>1)</sup> In this case,  $S_1(\Gamma, \chi)$  denotes simply the space of all holomorphic automorphic forms of weight one with  $\chi$ .

<sup>2)</sup> For this modified formula of  $d_1$ , refer to Hiramatsu ([6], Remark 1 in § 2).

The purpose of this paper is to give a similar formula of the number  $d_1$  when the group  $\Gamma$  is of finite type *reduced at infinity* and  $\chi^2 \neq 1$ , by using the method of Selberg ([3], [4]). In this case, the operator  $\omega_s$  in [1] is not generally completely continuous on the space  $L^2(\Gamma \backslash \tilde{S}, \chi)$  and a new term from cusp ought to have added to the formula (1). The notation used here will generally be those of [1].

It is a pleasure to record my thanks to Professor H. Shimizu for his advice and encouragement during the writing up.

**§ 1. The Selberg eigenspace  $\mathfrak{M}(1, -3/2, \chi)$  for a group  $\Gamma$  of finite type**

Let  $\Gamma$  be a fuchsian group of the first kind containing the element  $-I$ , and suppose that  $\Gamma$  has a non-compact fundamental domain in  $S$ . Let  $T$  be the real torus and put  $\tilde{S} = S \times T$ . Denote by  $L^2(\Gamma \backslash \tilde{S}, \chi)$  the following set

$$\left\{ f \in L^2(\Gamma \backslash \tilde{S}) : f(g(z, \phi)) = \chi(g)f(z, \phi) \text{ for all } g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma \right\},$$

where

$$f(g(z, \phi)) = f\left(\frac{az + b}{cz + d}, \phi + \arg(cz + d)\right).$$

Moreover we denote by  $\mathfrak{M}_r(k, \lambda, \chi) = \mathfrak{M}(k, \lambda, \chi)$  the set of all functions  $f(z, \phi)$  satisfying the following conditions:

- (i)  $f(z, \phi) \in L^2(\Gamma \backslash \tilde{S}, \chi)$ ,
- (ii)  $\bar{\Delta}f(z, \phi) = \lambda f(z, \phi), \frac{\partial}{\partial \phi} f(z, \phi) = -ikf(z, \phi)$ .

Then we have the following

LEMMA. *To each function  $f(z, \phi) \in \mathfrak{M}(1, \lambda, \chi)$ , we associate a function on  $S$  by letting*

$$F(z) = e^{i\phi}y^{-1/2}f(z, \phi).$$

*Then the function  $F(z)$  belongs to  $S_1(\Gamma, \chi)$  if and only if*

$$f(z, \phi) \in \mathfrak{M}(1, -3/2, \chi).$$

*Proof.* For each  $F(z) \in S_1(\Gamma, \chi)$  we define  $f(z, \phi)$  on  $\tilde{S}$  by

$$(1.1) \quad f(z, \phi) = e^{-i\phi}y^{1/2}F(z).$$

Then the function  $f(z, \phi)$  satisfies the conditions:

- (1.2)  $f(g(z, \phi)) = \chi(g)f(z, \phi)$  for all  $g \in \Gamma$ ;
- (1.3)  $(\partial/\partial\phi)f(z, \phi) = -if(z, \phi)$ ;
- (1.4)  $\bar{\Delta}f(z, \phi) = -(3/2)f(z, \phi)$  by regularity of  $F(z)$  on  $S$ ;
- (1.5) Since  $y^{1/2}|F(z)|$  is bounded on  $S$ ,

$$\begin{aligned} \|f\| &= \frac{1}{\pi} \int_{\Gamma \backslash \mathbb{S}} |e^{-i\phi} y^{1/2} F(z)|^2 \frac{dx dy d\phi}{y^2} \\ &= \int_{\Gamma \backslash S} |y^{1/2} F(z)|^2 \frac{dx dy}{y^2} < \infty. \end{aligned}$$

Therefore, by (1.2)~(1.5), the function  $f(z, \phi)$  belongs to  $\mathfrak{M}(1, -3/2, \chi)$ . We now prove conversely that any function in  $\mathfrak{M}(1, -3/2, \chi)$  must be of the form (1.1) with  $F(z) \in S_1(\Gamma, \chi)$ . Let  $f(z, \phi)$  be a function in  $\mathfrak{M}(1, -3/2, \chi)$ . Put

$$F(z) = e^{i\phi} y^{-1/2} f(z, \phi).$$

Then the  $\Gamma$ -invariance of  $f(z, \phi)$  is equivalent to a transformation law for  $F(z)$ :

$$F(g(z)) = \chi(g)(cz + d)F(z)$$

for all  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ . Therefore, it is sufficient for the proof of the latter half of our lemma to show that  $F(z)$  is holomorphic with respect to the complex variable  $z$  on  $S$ , and vanishes at every cusp of  $\Gamma$ .

Let  $\mathfrak{g}$  be the Lie algebra of  $SL_2(\mathbf{R}) (=G)$ . Then we can take the basis  $\alpha$  of  $\mathfrak{g}$  such that the Lie derivatives associated with the elements of  $\alpha$  are given by the following invariant differential operators:

$$\begin{aligned} X &= y \cos 2\phi \frac{\partial}{\partial x} - y \sin 2\phi \frac{\partial}{\partial y} + \frac{1}{2}(\cos 2\phi - 1) \frac{\partial}{\partial \phi}, \\ Y &= y \sin 2\phi \frac{\partial}{\partial x} + y \cos 2\phi \frac{\partial}{\partial y} + \frac{1}{2} \sin 2\phi \frac{\partial}{\partial \phi}, \\ \Phi &= \frac{\partial}{\partial \phi}. \end{aligned}$$

It is easy to see that

$$\bar{\Delta} = \left(X + \frac{1}{2}\Phi\right)^2 + Y^2 + \Phi^2.$$

Now we put

$$A^- = 2\left(X + \frac{1}{2}\Phi\right) + 2iY.$$

Then, the function  $F(z)$  is holomorphic on  $S$  if and only if

$$(1.6) \quad A^-f(z, \phi) = 0.$$

To prove (1.6), first note that the operation of  $A^-$  depends only on the representations of the Lie algebra  $\mathfrak{g}$ . Let  $L^2_a(\Gamma \backslash G, \chi)$  be the discrete part of the space  $L^2(\Gamma \backslash G, \chi)$ . Then  $f \in L^2_a(\Gamma \backslash G, \chi)$ . Let

$$L^2_a(\Gamma \backslash G, \chi) = \sum_i V_i$$

be the irreducible splitting of the space  $L^2_a(\Gamma \backslash G, \chi)$  and put

$$f = \sum_i f_i \quad (f_i \in V_i).$$

Then, if  $f_i \neq 0$ , we have

$$\tilde{\Delta}f_i = -\frac{3}{2}f_i, \quad \frac{\partial}{\partial \phi}f_i = -\sqrt{-1}f_i.$$

Therefore, each subspace  $V_i$  such that  $f_i \neq 0$  is isomorphic to the space  $H_1$  of the irreducible representation of the limit of discrete series. Hence it is sufficient for the proof of (1.6), to show that for any highest weight vector  $\varphi$  in  $H_1$ ,

$$(1.7) \quad A^-\varphi = 0.$$

For example, by Lemma 5.6 in [2], the formula (1.7) is well known.

Next we shall see the condition for  $F(z)$  at every cusp of  $\Gamma$ . Let  $s$  be a cusp of  $\Gamma \cap \text{Ker } \chi$ . We may assume that  $s = \infty$  and the intersection of a fundamental domain for  $\Gamma$  and a neighborhood of  $\infty$  is the following type

$$\{z = x + iy: 0 \leq x \leq 1, y \geq M\},$$

where  $M$  denotes a positive constant. Then, by the given condition  $f(z, \phi) \in L^2(\Gamma \backslash \tilde{S}, \chi)$ ,

$$\int_M^\infty \left\{ \int_0^1 y |F(z)|^2 dx \right\} \frac{dy}{y^2} < \infty.$$

Let

$$F(z) = \sum_{n=-\infty}^\infty a_n e^{2\pi i n z}$$

be the Fourier expansion of  $F$  at  $\infty$ . Then, we have

$$\begin{aligned} \int_0^1 |F(z)|^2 dx &= \int_0^1 (\sum_n a_n e^{2\pi i n z}) (\sum_m \bar{a}_m e^{-2\pi i m \bar{z}}) dx \\ &= \sum_{n,m} a_n \bar{a}_m \int_0^1 e^{2\pi i (n-m)x - 2\pi i (n+m)y} dx \\ &= \sum_n |a_n|^2 e^{-4\pi n y}. \end{aligned}$$

Therefore,

$$\int_M^\infty y (\sum_n |a_n|^2 e^{-4\pi n y}) \frac{dy}{y^2} = \sum_n |a_n|^2 \int_M^\infty y^{-1} e^{-4\pi n y} dy.$$

If  $n \leq 0$ , then

$$\int_M^\infty y^{-1} e^{-4\pi n y} dy = \infty,$$

so that  $a_n = 0$  for all  $n \leq 0$ .

Q.E.D.

**§ 2. Eisenstein series and continuous spectrum**

**2.1.** This section is essentially based upon the work of Selberg ([3], [4]). We shall review the definition and elementary properties of Eisenstein series for the cusp  $\infty$ , and the spectral decomposition of  $L^2(\Gamma \backslash \tilde{S}, \chi)$  (abbreviated hereafter as  $L^2(\Gamma \backslash \tilde{S})$ ). Let  $\Gamma$  be of finite type reduced at  $\infty$ , namely,  $\infty$  is a cusp of  $\Gamma$  and the stabilizer  $\Gamma_\infty$  of  $\infty$  in  $\Gamma$  is equal to  $\pm \Gamma_0$  with  $\Gamma_0 = \left\{ \begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix} : m \in \mathbf{Z} \right\}$ . The Eisenstein series  $E_\chi(z, \phi; s)$  attached to the cusp  $\infty$  and  $\chi$  is then defined by

$$(2.1) \quad E_\chi(z, \phi; s) = \sum_{\substack{M \in \Gamma_\infty \backslash \Gamma \\ M = \begin{pmatrix} * & * \\ c & d \end{pmatrix}}} \frac{\bar{\chi}(M) y^s}{|cz + d|^{2s}} e^{-i(\phi + \arg(cz + d))},$$

where  $s = \sigma + ir$  with  $\sigma > 1$ . It is easy to check that

- (i)  $E_\chi(M(z, \phi); s) = \chi(M) E_\chi(z, \phi; s)$  for  $M \in \Gamma$ ;
- (ii)  $\bar{\Delta} E_\chi(z, \phi; s) = \left\{ s(s-1) - \frac{5}{4} \right\} E_\chi(z, \phi; s)$ ;
- (iii)  $\frac{\partial}{\partial \phi} E_\chi(z, \phi; s) = -i E_\chi(z, \phi; s)$ .

Since  $E_\chi(z+1, \phi; s) = E_\chi(z, \phi; s)$ , we have a Fourier-Bessel expansion of the form

$$E_\chi(z, \phi; s) = \sum_{m=-\infty}^\infty a_{m,\chi}(y, \phi; s) e^{2\pi i m x},$$

where

$$a_{m,x}(y, \phi; s) = \int_0^1 E_x(z, \phi; s) e^{-2\pi i m x} dx .$$

Let us now try to find the constant term  $a_{0,x}(y, \phi; s)$  explicitly. Put

$$a_{0,x}(y, \phi; s) = e^{-i\phi} a_{0,x}(y; s) ,$$

and

$$E_x(z; s) = e^{i\phi} E_x(z, \phi; s) ;$$

then

$$\begin{aligned} a_{0,x}(y; s) &= \int_0^1 E_x(z; s) dx \\ &= \int_0^1 \sum_{\substack{M \in \Gamma_\infty \backslash \Gamma \\ M = \begin{pmatrix} * & * \\ c & d \end{pmatrix} \\ c > 0}} \frac{\bar{\chi}(M) y^s}{|cz + d|^{2s}} \lambda(cz + d)^{-1} dx \quad \left( \lambda(z) = \frac{z}{|z|}, z \neq 0 \right) \\ &= y^s + \int_0^1 \sum_{\substack{M \in \Gamma_\infty \backslash \Gamma \\ M = \begin{pmatrix} * & * \\ c & d \end{pmatrix} \\ c > 0}} \frac{\bar{\chi}(M) y^s}{|cz + d|^{2s}} \lambda(cz + d)^{-1} dx \\ &= y^s + \int_{-\infty}^{\infty} \sum_{\substack{M \in \Gamma_\infty \backslash \Gamma / \Gamma_\infty \\ M = \begin{pmatrix} * & * \\ c & d \end{pmatrix} \\ c > 0}} \frac{\bar{\chi}(M) y^s}{|cz + d|^{2s}} \lambda(cz + d)^{-1} dx \\ &= y^s + \sum_{\substack{c > 0 \\ d \bmod c \\ \begin{pmatrix} * & * \\ c & d \end{pmatrix} \in \Gamma}} \frac{\bar{\chi}(c, d)}{|c|^{2s}} \int_{-\infty}^{\infty} \frac{y^s}{\left| z + \frac{d}{c} \right|^{2s}} \lambda\left( z + \frac{d}{c} \right)^{-1} dx \\ &= y^s + y^s \varphi_{0,x}(s) \int_{-\infty}^{\infty} \frac{\lambda(z)^{-1}}{|z|^{2s}} dx \\ &= y^s + y^{1-s} \varphi_{0,x}(s) \int_{-\infty}^{\infty} \frac{\lambda(i+t)^{-1}}{(1+t^2)^s} dt , \end{aligned}$$

where

$$\varphi_{0,x}(s) = \sum_{\substack{c > 0 \\ d \bmod c \\ \begin{pmatrix} * & * \\ c & d \end{pmatrix} \in \Gamma}} \frac{\bar{\chi}(c, d)}{|c|^{2s}}$$

Furthermore,

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{\lambda(i+t)^{-1}}{(1+t^2)^s} dt &= \int_0^\pi \frac{(i + \cot \theta)^{-1}}{|\operatorname{cosec} \theta|^{2s-1}} \operatorname{cosec}^2 \theta d\theta \\ &= \int_0^\pi \sin^{2s-2} \theta (\cos \theta - i \sin \theta) d\theta \end{aligned}$$

$$\begin{aligned}
 &= -i \int_0^\pi \sin^{2s-1} \theta \, d\theta \\
 &= -i\sqrt{\pi} \frac{\Gamma(s)}{\Gamma\left(s + \frac{1}{2}\right)}.
 \end{aligned}$$

Therefore we have

$$a_{0,\chi}(y; s) = y^s - i\sqrt{\pi} \frac{\Gamma(s)}{\Gamma\left(s + \frac{1}{2}\right)} \varphi_{0,\chi}(s) y^{1-s}.$$

**2.2.** Since the group  $\Gamma$  is reduced at  $\infty$ , the integral operator  $\omega_\delta$  in [1] is not generally completely continuous on  $L^2(\Gamma \backslash \tilde{S})$  and beside the discrete spectrum in  $L^2(\Gamma \backslash \tilde{S})$ , the operator  $\omega_\delta$  has one or more continuous spectra in  $L^2(\Gamma \backslash \tilde{S})$ . The space  $L^2(\Gamma \backslash \tilde{S})$  has the following spectral decomposition

$$L^2(\Gamma \backslash \tilde{S}) = L_0^2(\Gamma \backslash \tilde{S}) \oplus L_{sp}^2(\Gamma \backslash \tilde{S}) \oplus L_{cont}^2(\Gamma \backslash \tilde{S}),$$

where  $L_0^2$  is the space of cusp forms and is discrete,  $L_{sp}^2$  is the discrete part of the orthogonal complement of  $L_0^2$ , and  $L_{cont}^2$  is continuous part of the spectrum, In the following we shall only consider the case

$$\chi\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right) = 1.$$

By using the analytic continuation of the Eisenstein series (2.1) as a function of  $s$  for  $s = 1/2 + ir$ , we put

$$\tilde{H}_\delta(z, \phi; z', \phi') = \frac{1}{4\pi^2} \int_{-\infty}^\infty h(r) E_\chi\left(z, \phi; \frac{1}{2} + ir\right) \overline{E_\chi\left(z', \phi'; \frac{1}{2} + ir\right)} \, dr,$$

where  $h(r)$  denotes the eigenvalue of  $\omega_\delta$  in  $\mathfrak{M}(1, \lambda, \chi)$  ([1], p. 217):

$$(2.2) \quad h(r) = 2^{2+\delta} \pi \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1+\delta}{2}\right)}{\Gamma(\delta) \Gamma\left(1 + \frac{\delta}{2}\right)} \Gamma\left(\frac{\delta}{2} + ir\right) \Gamma\left(\frac{\delta}{2} - ir\right)$$

with  $\lambda = s(s - 1) - 5/4$  and  $s = 1/2 + ir$ . Then the integral operator  $\tilde{K}_\delta = K_\delta - \tilde{H}_\delta$  is now completely continuous on  $L^2(\Gamma \backslash \tilde{S})$  and has all discrete spectra of  $K_\delta$ , where

$$K_\delta(z, \phi; z', \phi') = 2 \sum_{M \in \Gamma/\{\pm I\}} \chi(M) \omega_\delta(z, \phi; M(z', \phi')).$$

Furthermore, an eigenvalue of  $f(z, \phi)$  in  $L^2_0(\Gamma \backslash \tilde{S}) \oplus L^2_{sp}(\Gamma \backslash \tilde{S})$  for  $\tilde{K}_\delta$  is equal to that for  $K_\delta$  and the image of  $\tilde{K}_\delta$  on it is contained in  $L^2_0(\Gamma \backslash \tilde{S})$ . Considering the trace of  $\tilde{K}_\delta$  on  $L^2_0(\Gamma \backslash \tilde{S})$ , we now obtain the following modified trace formula

$$(2.3) \quad \begin{aligned} \sum_{n=1}^{\infty} h(\lambda^{(n)}) &= \int_{\tilde{D}} \tilde{K}_\delta(z, \phi; z, \phi) d(z, \phi) \\ &= \int_{\tilde{D}} \left\{ 2 \sum_{M \in \Gamma / \{\pm I\}} \chi(M) \omega_\delta(z, \phi; M(z, \phi)) - \tilde{H}_\delta(z, \phi; z, \phi) \right\} d(z, \phi), \end{aligned}$$

where  $\tilde{D}$  denotes a fundamental domain of  $\Gamma$  in  $\tilde{S}$ , and each of  $\lambda^{(n)}$  denotes an eigenvalue corresponding to an orthogonal basis  $\{f^{(n)}\}$  for  $L^2_0(\Gamma \backslash \tilde{S})$ .

**§ 3. A formula for the dimension  $d_1$**

The purpose of this section is to obtain an explicit formula for the dimension  $d_1$  by calculating the integrals in (2.3). We put

$$\begin{aligned} &\int_{\tilde{D}} \left\{ 2 \sum_{M \in \Gamma / \{\pm I\}} \chi(M) \omega_\delta(z, \phi; M(z, \phi)) - \tilde{H}_\delta(z, \phi; z, \phi) \right\} d(z, \phi) \\ &= J(I) + J(P) + J(R) + J(\infty), \end{aligned}$$

where  $J(I)$ ,  $J(P)$ ,  $J(R)$ , and  $J(\infty)$  denote respectively the identity component, the hyperbolic component, the elliptic component, and the parabolic component of the traces. Then the components  $J(I)$ ,  $J(P)$ , and  $J(R)$  were obtained already in [1] and [6]. So in the following we shall calculate the component  $J(\infty)$ . Since  $\Gamma$  is reduced at  $\infty$ , the set  $\Gamma / \{\pm I\} - \{I\}$  gives a complete system of representatives of the parabolic conjugacy classes in  $\Gamma / \{\pm I\}$  and for each  $\gamma \in \Gamma_\infty - \{\pm I\}$ , the stabilizer of  $\gamma$  in  $\Gamma$  is always equal to  $\Gamma_\infty$ . Put

$$P_\Gamma = \left\{ \gamma^{-1} \begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix} \gamma : m \in \mathbb{Z}, m \neq 0, \gamma \in \Gamma_\infty \setminus \Gamma \right\}.$$

Then  $P_\Gamma$  is the set consisting of all parabolic elements in  $\Gamma / \{\pm I\}$ . Hence

$$\begin{aligned} &\int_{\tilde{D}} \sum_{\substack{M \in \Gamma / \{\pm I\} \\ M: \text{parabolic}}} \chi(M) \omega_\delta(z, \phi; M(z, \phi)) d(z, \phi) = \sum_{\substack{M \in \Gamma / \{\pm I\} \\ M: \text{parabolic}}} \int_{\tilde{D}} \chi(M) \omega_\delta(z, \phi; M(z, \phi)) d(z, \phi) \\ &= \sum_{\substack{M \in \Gamma_0 \\ M \neq I}} \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \int_{\tilde{D}} \omega_\delta(\gamma(z, \phi); M\gamma(z, \phi)) d(z, \phi) \\ &= \sum_{\substack{M \in \Gamma_0 \\ M \neq I}} \int_{\tilde{D}_{\Gamma_\infty}} \omega_\delta(z, \phi; M(z, \phi)) d(z, \phi), \end{aligned}$$



where  $\tilde{D}_{\Gamma_\infty}$  denotes a fundamental domain of  $\Gamma_\infty$  in  $\tilde{S}$ . Therefore we have

$$J(\infty) = \lim_{Y \rightarrow \infty} \left\{ \int_0^Y \int_0^1 \int_0^\pi 2 \sum_{\substack{M \in \Gamma_0 \\ M \neq I}} \omega_\delta(z, \phi; M(z, \phi)) d(z, \phi) - \int_{\tilde{D}_Y} \tilde{H}_\delta(z, \phi; z, \phi) d(z, \phi) \right\},$$

where  $\tilde{D}_Y = \tilde{D} - \tilde{D}'_Y$  with the direct product  $\tilde{D}'_Y$  of the real torus  $T$  and the subdomain of the strip determined by  $\text{Im } z > Y$  for a sufficiently large  $Y > 0$ . Furthermore

$$\begin{aligned} \text{(A)} \quad & \int_0^Y \int_0^1 \int_0^\pi 2 \sum_{\substack{M \in \Gamma_0 \\ M \neq I}} \omega_\delta(z, \phi; M(z, \phi)) d(z, \phi) \\ &= 2 \sum_{\substack{m \in \mathbb{Z} \\ m \neq 0}} \int_0^Y \int_0^1 \int_0^\pi \omega_\delta(z, \phi; M_0^m(z, \phi)) d(z, \phi) \quad \left( M_0 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right) \\ &= 2\pi \sum_{\substack{m \in \mathbb{Z} \\ m \neq 0}} \int_0^Y \left| \frac{y}{2yi - m} \right|^\delta \frac{y}{2yi - m} \frac{dy}{y^2} \\ &= 2^{\delta+4} \pi \sum_{m=1}^\infty \int_0^Y \frac{1}{(4 + (m/y)^2)^{\delta/2+1}} \frac{dy}{y^2}. \end{aligned}$$

If we put

$$k(t) = \frac{1}{(4 + t)^{\delta/2+1}} \quad \text{and} \quad f(x) = 2 \int_0^Y k\left(\frac{x^2}{y^2}\right) \frac{dy}{y^2},$$

then

$$f(x) = \frac{1}{x} \int_{x^2/Y^2}^\infty \frac{k(t)}{\sqrt{t}} dt;$$

and hence

$$\int_0^Y \int_0^1 \int_0^\pi 2 \sum_{\substack{M \in \Gamma_0 \\ M \neq I}} \omega_\delta(z, \phi; M(z, \phi)) d(z, \phi) = 2^{\delta+3} \pi \sum_{m=1}^\infty f(m).$$

Now we make use of a summation formula due to Euler-MacLaurin:

$$\text{(3.1)} \quad \sum_{m=1}^\infty f(m) = \frac{1}{2} f(1) + \int_1^\infty f'(x) \{x\} dx + \int_1^\infty f(x) dx,$$

where  $[x]$  denotes the greatest integer in  $x$  and  $\{x\} = x - [x] - 1/2$ . Then we have

$$\frac{1}{2}f(1) \longrightarrow 2^{-(\delta+2)} \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{\delta+1}{2}\right)}{\Gamma\left(1 + \frac{\delta}{2}\right)},$$

and

$$\int_1^\infty f'(x)\{x\} dx = 2^{-(\delta+1)} \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{\delta+1}{2}\right)}{\Gamma\left(1 + \frac{\delta}{2}\right)} \left(C - \frac{1}{2}\right) + o(1),$$

as  $Y \rightarrow \infty$ , where  $C$  is Euler’s constant. As for the third integral of (3.1), we have first the following by [3]:

$$\begin{aligned} &\int_1^\infty f(x) dx \\ &= 2^{-(\delta+1)} \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{\delta+1}{2}\right)}{\Gamma\left(1 + \frac{\delta}{2}\right)} \log Y + \frac{1}{2} \int_0^\infty \frac{\log t \cdot k(t)}{\sqrt{t}} dt + o(1), \end{aligned}$$

as  $Y \rightarrow \infty$ . Furthermore,

$$\begin{aligned} &\frac{1}{2} \int_0^\infty \frac{\log t \cdot k(t)}{\sqrt{t}} dt = \frac{1}{2} \int_0^\infty \frac{\log t}{\sqrt{t}(4+t)^{\delta/2+1}} dt \\ &= 2^{-(\delta+1)} \log 2 \int_0^\infty \frac{1}{\sqrt{t}(1+t)^{\delta/2+1}} dt + 2^{-(\delta+2)} \int_0^\infty \frac{\log t}{\sqrt{t}(1+t)^{\delta/2+1}} dt \\ &= 2^{-(\delta+1)} \log 2 \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{\delta+1}{2}\right)}{\Gamma\left(1 + \frac{\delta}{2}\right)} \\ &\quad + 2^{-(\delta+2)} \left\{ -\sum_{k=0}^\infty \binom{-\delta/2-1}{k} \frac{1}{\left(\frac{1}{2} + k\right)^2} + \sum_{k=0}^\infty \binom{-\delta/2-1}{k} \frac{1}{\left(\frac{\delta+9}{2} + k\right)^2} \right\}, \end{aligned}$$

where  $-2 < \delta < 4$ . Summing up the above results, we obtain for the first half of  $J(\infty)$ ,

$$\int_0^Y \int_0^1 \int_0^\pi 2 \sum_{\substack{M \in \Gamma_0 \\ M=I}} \omega_s(z, \phi; M(z, \phi)) d(z, \phi)$$

$$\begin{aligned}
 &= 2^2\pi \frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{\delta+1}{2}\right)}{\Gamma\left(1+\frac{\delta}{2}\right)}(\log Y + C + \log 2) \\
 &\quad + 2\pi\left\{-\sum_{k=0}^{\infty}\binom{-\delta/2-1}{k}\frac{1}{\left(\frac{1}{2}+k\right)^2}\right. \\
 &\quad \left. + \sum_{k=0}^{\infty}\binom{-\delta/2-1}{k}\frac{1}{\left(\frac{\delta+9}{2}+k\right)^2}\right\} + o(1),
 \end{aligned}$$

as  $Y \rightarrow \infty$ .

(B) We define the following the compact part of  $E_\chi(z, \phi; s)$ :

$$\tilde{E}_\chi(z, \phi; s) = \begin{cases} E_\chi(z, \phi; s) & \text{for } y \leq Y, \\ E_\chi(z, \phi; s) - e^{-i\phi}(y^s + \varphi_\chi(s)y^{1-s}) & \text{for } y > Y, \end{cases}$$

with  $\varphi_\chi(s) = -i\sqrt{\pi} \frac{\Gamma(s)}{\Gamma\left(s + \frac{1}{2}\right)}\varphi_{0,\chi}(s)$ . Then,

$$\begin{aligned}
 &\int_{D_Y} \tilde{H}_\delta(z, \phi; z, \phi) d(z, \phi) \\
 &= \frac{1}{4\pi^2} \int_{D_Y} \int_{-\infty}^{\infty} h(r) E_\chi\left(z, \phi; \frac{1}{2} + ir\right) \overline{E_\chi\left(z, \phi; \frac{1}{2} - ir\right)} dr d(z, \phi) \\
 &= \frac{1}{4\pi^2} \lim_{\sigma \rightarrow \frac{1}{2}} \int_{D_Y} \int_{-\infty}^{\infty} h(r) |\tilde{E}(z, \phi; \sigma + ir)|^2 dr d(z, \phi) + o(1),
 \end{aligned}$$

as  $Y \rightarrow \infty$ . Moreover,

$$\begin{aligned}
 &\int_{D_Y} \int_{-\infty}^{\infty} h(r) |\tilde{E}_\chi(z, \phi; s)|^2 dr d(z, \phi) = \int_{D_Y} \int_{-\infty}^{\infty} h(r) E_\chi(z, \phi; s) \overline{E_\chi(z, \phi; s)} dr d(z, \phi) \\
 &\quad + \int_{D_Y} \int_{-\infty}^{\infty} h(r) \{E_\chi(z, \phi; s) - e^{-i\phi}(y^s + \varphi_\chi(s)y^{1-s})\} \\
 &\quad \times \{\overline{E_\chi(z, \phi; s)} - e^{i\phi}(y^{\bar{s}} + \overline{\varphi_\chi(s)}y^{1-\bar{s}})\} dr d(z, \phi); \\
 &\int_{D_Y} \int_{-\infty}^{\infty} h(r) E_\chi(z, \phi; s) \overline{E_\chi(z, \phi; s)} dr d(z, \phi) \\
 &= \int_{\cup M D_Y} \int_{-\infty}^{\infty} h(r) E_\chi(z, \phi; s) y^{\bar{s}} e^{i\phi} dr d(z, \phi) \\
 &= \int_0^Y \int_0^1 \int_0^\pi \int_{-\infty}^{\infty} h(r) E_\chi(z, \phi; s) y^{\bar{s}-2} e^{i\phi} dr d\phi dx dy
 \end{aligned}$$

$$\begin{aligned}
 & - \int_{\substack{\cup M D_Y \\ M \in F_\infty \setminus \Gamma, M \neq I}} \int_{-\infty}^{\infty} h(r) E_x(z, \phi; s) y^s e^{i\phi} dr d(z, \phi) \\
 & = \pi \int_0^Y \int_{-\infty}^{\infty} h(r) (y^s + \varphi(s) y^{1-s}) y^{s-2} dr dy \\
 & \quad - \int_{D_Y} \int_{-\infty}^{\infty} h(r) E_x(z, \phi; s) \{ \overline{E_x(z, \phi; s)} - e^{i\phi} y^s \} dr d(z, \phi) \\
 & = \pi \int_0^Y \int_{-\infty}^{\infty} h(r) y^{2\sigma-2} dr dy + \pi \int_0^Y \int_{-\infty}^{\infty} h(r) \varphi_x(s) y^{-1-2i\tau} dr dy \\
 & \quad - \int_{D_Y} \int_{-\infty}^{\infty} h(r) \{ E_x(z, \phi; s) - e^{-i\phi} (y^s + \varphi_x(s) y^{1-s}) \} \\
 & \quad \times \{ \overline{E_x(z, \phi; s)} - e^{i\phi} (y^s + \overline{\varphi_x(s)} y^{1-s}) \} dr d(z, \phi) \\
 & \quad - \pi \int_Y^\infty \int_{-\infty}^{\infty} h(r) \overline{\varphi_x(s)} y^{-1+2i\tau} dr dy - \pi \int_Y^\infty \int_{-\infty}^{\infty} h(r) |\varphi_x(s)|^2 dr dy .
 \end{aligned}$$

Therefore we have

$$\begin{aligned}
 \int_{D_Y} \tilde{H}_\delta(z, \phi; z, \phi) d(z, \phi) & = \frac{1}{4\pi} \lim_{\sigma \rightarrow \frac{1}{2}} \left\{ \int_0^Y \int_{-\infty}^{\infty} h(r) y^{2\sigma-2} dr dy \right. \\
 & \quad - \int_Y^\infty \int_{-\infty}^{\infty} h(r) |\varphi_x(s)|^2 y^{-2\sigma} dr dy + \int_0^Y \int_{-\infty}^{\infty} h(r) \varphi_x(s) y^{-1-2i\tau} dr dy \\
 & \quad \left. - \int_Y^\infty \int_{-\infty}^{\infty} h(r) \overline{\varphi_x(s)} y^{-1+2i\tau} dr dy \right\} + o(1),
 \end{aligned}$$

as  $Y \rightarrow \infty$ . Here we calculate the integrals appearing in the above expression.

$$\begin{aligned}
 \text{(i)} \quad \int_0^Y \int_{-\infty}^{\infty} h(r) y^{2\sigma-2} dr dy & = \frac{Y^{2\sigma-1}}{2\sigma-1} \int_{-\infty}^{\infty} h(r) dr \\
 & = \frac{Y^{2\sigma-1} - Y^{1-2\sigma}}{2\sigma-1} \int_{-\infty}^{\infty} h(r) dr + \frac{Y^{1-2\sigma}}{2\sigma-1} \int_{-\infty}^{\infty} h(r) dr ; \\
 \int_{-\infty}^{\infty} h(r) dr & = 2^{2+\delta} \pi \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1+\delta}{2}\right)}{\Gamma(\delta) \Gamma\left(1 + \frac{\delta}{2}\right)} \int_{-\infty}^{\infty} \left| \Gamma\left(\frac{\delta}{2} + ir\right) \right|^2 dr \\
 & = 2^{2+\delta} \pi \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1+\delta}{2}\right)}{\Gamma(\delta) \Gamma\left(1 + \frac{\delta}{2}\right)} \cdot \sqrt{\pi} \Gamma\left(\frac{\delta}{2}\right) \Gamma\left(\frac{\delta+1}{2}\right) \\
 & = 2^3 \pi^{5/2} \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1+\delta}{2}\right)}{\Gamma\left(1 + \frac{\delta}{2}\right)} ;
 \end{aligned}$$

$$\lim_{\sigma \rightarrow \frac{1}{2}} \frac{Y^{2\sigma-1} - Y^{1-2\sigma}}{2\sigma - 1} = 2 \log Y.$$

$$\begin{aligned} \text{(ii)} \quad \int_0^Y \int_{-\infty}^{\infty} h(r)\varphi_x(s)y^{-1-2ir} dr dy &= \int_0^Y \int_{-\infty}^{\infty} h(r)\varphi_x(s)e^{-2\pi i \cdot \log y} dr d(\log y) \\ &= \int_{-\infty}^{\log Y} \int_{-\infty}^{\infty} h(r)\varphi_x(s)e^{-ir(2y')} dr dy' \\ &= \pi h(0)\varphi_x\left(\frac{1}{2}\right) \end{aligned}$$

as  $Y \rightarrow \infty$ . By a similar calculation as in the above,

$$\int_Y^{\infty} \int_{-\infty}^{\infty} h(r)\overline{\varphi_x(s)}y^{-1+2ir} dr dy = \int_{\log Y}^{\infty} \int_{-\infty}^{\infty} h(r)\overline{\varphi_x(s)}e^{ir(2y')} dr dy'$$

as  $Y \rightarrow \infty$ . Hence we have

$$\begin{aligned} \lim_{\sigma \rightarrow \frac{1}{2}} \left\{ \int_0^Y \int_{-\infty}^{\infty} h(r)\varphi_x(s)y^{-1-2ir} dr dy - \int_Y^{\infty} \int_{-\infty}^{\infty} h(r)\overline{\varphi_x(s)}y^{-1+2ir} dr dy \right\} \\ = \pi h(0)\varphi_x\left(\frac{1}{2}\right) \end{aligned}$$

as  $Y \rightarrow \infty$ . We note that  $\varphi_x\left(\frac{1}{2}\right) = \pm 1$ .

$$\begin{aligned} \text{(iii)} \quad - \int_Y^{\infty} \int_{-\infty}^{\infty} h(r)|\varphi_x(s)|^2 y^{-2\sigma} dr dy + \frac{Y^{1-2\sigma}}{2\sigma - 1} \int_{-\infty}^{\infty} h(r) dr \\ = - \frac{Y^{1-2\sigma}}{2\sigma - 1} \int_{-\infty}^{\infty} h(r)|\varphi_x(s)|^2 dr + \frac{Y^{1-2\sigma}}{2\sigma - 1} \int_{-\infty}^{\infty} h(r) dr \\ = - Y^{1-2\sigma} \int_{-\infty}^{\infty} h(r) \frac{|\varphi_x(s)|^2 - 1}{2\sigma - 1} dr. \end{aligned}$$

By the Maass-Selberg relation ([3], [4]), we know that

$$|\varphi_x(s)| \longrightarrow 1 \quad \text{as } \sigma \longrightarrow \frac{1}{2};$$

and hence

$$\varphi_x\left(\frac{1}{2} + ir\right)\varphi_x\left(\frac{1}{2} - ir\right) = 1,$$

namely,

$$\lim_{\sigma \rightarrow \frac{1}{2}} \int_{-\infty}^{\infty} h(r) \frac{|\varphi_x(s)|^2 - 1}{2\sigma - 1} dr = \int_{-\infty}^{\infty} h(r) \frac{\varphi'_x\left(\frac{1}{2} + ir\right)}{\varphi_x\left(\frac{1}{2} + ir\right)} dr.$$

By (i), (ii) and (iii), we obtain for the second half of the expression for  $J(\infty)$ ,

$$\begin{aligned} & \int_{D_Y} \tilde{H}_\delta(z, \phi; z, \phi) d(z, \phi) \\ &= 2^2\pi \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{\delta+1}{2}\right)}{\Gamma\left(1+\frac{\delta}{2}\right)} \log Y - \frac{1}{4\pi} \int_{-\infty}^{\infty} h(r) \frac{\varphi'_x\left(\frac{1}{2}+ir\right)}{\varphi_x\left(\frac{1}{2}+ir\right)} dr \\ &+ \frac{1}{4} h(0) \varphi_x\left(\frac{1}{2}\right) + o(1), \end{aligned}$$

as  $Y \rightarrow \infty$ . Summing up the above results (A) and (B), we obtain

$$\begin{aligned} J(\infty) &= 2^2\pi \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{\delta+1}{2}\right)}{\Gamma\left(1+\frac{\delta}{2}\right)} (C + \log 2) + \frac{1}{4\pi} \int_{-\infty}^{\infty} h(r) \frac{\varphi'_x\left(\frac{1}{2}+ir\right)}{\varphi_x\left(\frac{1}{2}+ir\right)} dr \\ &- \frac{1}{4} h(0) \varphi_x\left(\frac{1}{2}\right) \\ &+ 2\pi \left\{ -\sum_{k=0}^{\infty} \binom{-\delta/2-1}{k} \frac{1}{\left(\frac{1}{2}+k\right)^2} \right. \\ &\left. + \sum_{k=0}^{\infty} \binom{-\delta/2-1}{k} \frac{1}{\left(\frac{\delta+9}{2}+k\right)^2} \right\}. \end{aligned}$$

(C) We shall now calculate the limit  $\lim_{\delta \rightarrow 0} \delta J(\infty)$ .

(iv) we use the formula:

$$\binom{-\delta/2-1}{k} = (-1)^k \frac{\Gamma\left(k + \frac{\delta}{2} + 1\right)}{k! \Gamma\left(\frac{\delta}{2} + 1\right)}.$$

Then,

$$\lim_{\delta \rightarrow 0} \delta \left\{ -\sum_{k=0}^{\infty} \binom{-\delta/2-1}{k} \frac{1}{\left(\frac{1}{2}+k\right)^2} + \sum_{k=0}^{\infty} \binom{-\delta/2-1}{k} \frac{1}{\left(\frac{\delta+9}{2}+k\right)^2} \right\}$$

$$\begin{aligned}
 &= -\lim_{\delta \rightarrow 0} \frac{\delta}{\Gamma\left(\frac{\delta}{2} + 1\right)} \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma\left(k + \frac{\delta}{2} + 1\right)}{k! \left(\frac{1}{2} + k\right)^2} \\
 &\quad + \lim_{\delta \rightarrow 0} \frac{\delta}{\Gamma\left(\frac{\delta}{2} + 1\right)} \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma\left(k + \frac{\delta}{2} + 1\right)}{k! \left(\frac{\delta + 9}{2} + k\right)^2} \\
 &= -\lim_{\delta \rightarrow 0} \delta \sum_{k=0}^{\infty} \frac{(-1)^k}{\left(\frac{1}{2} + k\right)^2} + \lim_{\delta \rightarrow 0} \delta \sum_{k=0}^{\infty} \frac{(-1)^k}{\left(\frac{9}{2} + k\right)^2} = 0.
 \end{aligned}$$

(v) By the expression (2.2) of  $h(r)$ , we have

$$\lim_{\delta \rightarrow 0} \delta h(0) = 16\pi^2.$$

Therefore

$$\lim_{\delta \rightarrow 0} \delta \left( -\frac{1}{4} h(0) \varphi_x \left( \frac{1}{2} \right) \right) = -4\pi^2 \varphi_x \left( \frac{1}{2} \right).$$

By (2.2), we also have

$$h(r) \sim \frac{c(\delta) |r|^3}{|r| e^{\pi|r|}}$$

as  $r \rightarrow \infty$ , where  $c(\delta)$  is independent of  $r$  and  $\lim_{\delta \rightarrow 0} c(\delta)$  is finite. On the other hand, if we put

$$f(r) = \varphi'_x \left( \frac{1}{2} + ir \right) / \varphi_x \left( \frac{1}{2} + ir \right),$$

then

$$\begin{aligned}
 &\lim_{\delta \rightarrow +0} \delta \int_{-\infty}^{\infty} h(r) \frac{\varphi'_x}{\varphi_x} \left( \frac{1}{2} + ir \right) dr \\
 &= \lim_{\delta \rightarrow +0} \delta \left\{ \int_{-\infty}^{-N} h(r) \frac{\varphi'_x}{\varphi_x} \left( \frac{1}{2} + ir \right) dr + \int_{-N}^N h(r) \frac{\varphi'_x}{\varphi_x} \left( \frac{1}{2} + ir \right) dr \right. \\
 &\quad \left. + \int_N^{\infty} h(r) \frac{\varphi'_x}{\varphi_x} \left( \frac{1}{2} + ir \right) dr \right\}.
 \end{aligned}$$

Since the function  $f(r)$  is bounded on  $[-N, N]$ , we have

$$\begin{aligned}
 &\int_{-N}^N h(r) f(r) dr = O\left( \int_{-\infty}^{\infty} h(r) dr \right), \text{ i.e.,} \\
 &\lim_{\delta \rightarrow +0} \delta \int_{-N}^N h(r) f(r) dr = 0.
 \end{aligned}$$

Moreover, Since the operator  $\tilde{K}_\delta$  is completely continuons on  $L^2(\Gamma \backslash \tilde{S})$ , there exists some constant  $\delta_1$  such that

$$\int_N^\infty |r|^{\beta_1} \frac{f(r)}{|r|e^{\pi|r|}} dr < +\infty .$$

Then, for any  $\delta$  such that  $0 \leq \delta < \delta_1$ , the function  $|r|^\delta(f(r)/|r|e^{\pi|r|})$  is integrable on  $[N, \infty)$  and its convergence is uniform for  $\delta$ . Thus

$$\lim_{\delta \rightarrow +0} \delta \int_N^\infty |r|^\delta \frac{f(r)}{|r|e^{\pi|r|}} dr = 0 .$$

Therefore, we have

$$\lim_{\delta \rightarrow +0} \delta \int_{-\infty}^\infty h(r) \frac{\varphi'_\chi}{\varphi_\chi} \left( \frac{1}{2} + ir \right) dr = 0 .$$

Remark 1. The function  $(\varphi'_\chi/\varphi_\chi)(1/2 + ir)$  satisfies

$$\left| \frac{\varphi'_\chi}{\varphi_\chi} \left( \frac{1}{2} + ir \right) \right| < c \log (2 + |r|)$$

for some constant  $c$ . By this estimation we have again

$$\lim_{\delta \rightarrow 0} \delta \int_{-\infty}^\infty h(r) \frac{\varphi'_\chi}{\varphi_\chi} \left( \frac{1}{2} + ir \right) dr = 0 .$$

It is now clear that the above result, combined with the formula (1), proves the following.

**THEOREM F.** *Let  $\Gamma$  be a fuchsian group of the first kind containing the element  $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$  ( $= -I$ ) and suppose that  $\Gamma$  is reduced at infinity. Let  $\chi$  be a one-dimensional unitary representation of  $\Gamma$  such that  $\chi(-I) = -1$ ,  $\chi\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right) = 1$  and  $\chi^2 \neq 1$ , and denote by  $d_1$  the dimension for the linear space consisting of all cusp forms of weight one with respect to  $\Gamma$  with  $\chi$ . Then the dimension  $d_1$  is given by*

$$(3.2) \quad d_1 = \frac{1}{2} \sum_{\{M\}} \frac{\chi(M)}{[\Gamma(M) : \pm I]} \frac{\bar{\zeta}}{1 - \zeta^2} + \frac{1}{2} \operatorname{Res}_{s=0} \zeta^*(s) - \frac{1}{4} \varphi_\chi \left( \frac{1}{2} \right),$$

where the sum over  $\{M\}$  is taken over the distinct elliptic conjugacy classes of  $\Gamma/\{\pm I\}$ ,  $\Gamma(M)$  denotes the centralizer of  $M$  in  $\Gamma$ ,  $\zeta$  is one of the eigenvalues of  $M$ , and  $\zeta^*(s)$  denotes the Selberg type zeta-function defined in Section 0.



We may call the formula (3.2) a kind of Riemann-Roch type theorem for automorphic forms of weight one.

Example. Let  $p$  be a prime number such that  $p \equiv 3 \pmod{4}$ , ( $p \neq 3$ ), and let  $\Phi_0(p)$  be the group generated by the group  $\Gamma_0(p)$  and the element  $K = \begin{pmatrix} 0 & -\sqrt{p}^{-1} \\ \sqrt{p} & 0 \end{pmatrix}$ . Let  $\varepsilon$  be the Legendre symbol on  $\Gamma_0(p)$ :  $\varepsilon(L) = (d/p)$  for  $L = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(p)$ . Since  $\varepsilon(K^2) = \varepsilon(-I) = -1$ , we can define the odd characters  $\varepsilon^\pm$  on the Fricke group  $\Phi_0(p)$  such that  $\varepsilon^\pm(K) = \pm i$ . Then we have

$$S_1(\Gamma_0(p), \varepsilon) = S_1(\Phi_0(p), \varepsilon^+) \oplus S_1(\Phi_0(p), \varepsilon^-).$$

We put

$$\mu_1^\pm = \dim S_1(\Phi_0(p), \varepsilon^\pm).$$

Then

$$d_1 = \dim S_1(\Gamma_0(p), \varepsilon) = \mu_1^+ + \mu_1^-.$$

If  $\sigma^*(p)$  is the parabolic class number of  $\Phi_0(p)/\{\pm I\}$ , then  $\sigma^*(p) = 1$ . As shown in [6], the contribution from elliptic classes to  $\mu_1^\pm$  is given by

$$\frac{1}{2} \sum_{(M)} \frac{1}{[\Gamma(M): \pm I]} \frac{\bar{\zeta}}{1 - \bar{\zeta}^2} \varepsilon^\pm(M) = \mp \frac{1}{4} h.$$

We also have  $\varphi_\varepsilon^\pm(1/2) = \mp 1$ . Let  $P_\alpha$  ( $\alpha = 1, 2, 3, \dots$ ) be a complete system of representatives of the primitive hyperbolic conjugacy classes in  $\Gamma_0(p)/\{\pm I\}$  and let  $\lambda_{0,\alpha}$  be the eigenvalue ( $\lambda_{0,\alpha} > 1$ ) of a representative  $P_\alpha$ . We put

$$Z^*(s) = \sum_{\alpha=1}^\infty \sum_{k=1}^\infty \frac{\varepsilon(P_\alpha)^k \log \lambda_{0,\alpha}}{\lambda_{0,\alpha}^k - \lambda_{0,\alpha}^{-k}} |\lambda_{0,\alpha}^k + \lambda_{0,\alpha}^{-k}|^{-s}.$$

Then, we have the following formula for  $d_1$  which is our conclusion

$$d_1 = \mu_1^+ + \mu_1^- = \frac{1}{2} \operatorname{Res}_{s=0} Z^*(s).$$

Remark 2. For a general discontinuous group  $\Gamma$  of finite type containing the element  $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ , we obtain, in the same way as in the case of a group reduced at  $\infty$ , the contribution from parabolic classes to  $d_1$ .

## Addendum

Taking this opportunity, I would like to comment on our previous paper:

T. Hiramatsu and Y. Mimura,  
The modular equation and modular forms of weight one,  
Nagoya Math. J., **100** (1985), 145-162.

By an exchange of letters, it has been shown that the paper "Hohere Reziprozitatgesetze und Modulformen von Gewicht Eins, Jour. reine angew. Math., **361** (1985), 11-22", remarkably overlapping with our one, was written after its author had read a preprint of ours.

## REFERENCES

- [ 1 ] T. Hiramatsu, On some dimension formula for automorphic forms of weight one I, Nagoya Math. J., **85** (1982), 213-221.
- [ 2 ] H. Jacquet and R. P. Langlands, Automorphic forms on  $GL(2)$ , Lecture Notes in Math., **114**, Berlin, Heidelberg, New York: Springer 1970.
- [ 3 ] T. Kubota, Elementary theory of Eisenstein series, Tokyo-New York: Kodansha and Halsted 1973.
- [ 4 ] A. Selberg, Discontinuous groups and harmonic analysis, in Proc. International Math. Congr., Stockholm, 177-189, 1962.
- [ 5 ] J.-P. Serre, Modular forms of weight one and Galois representations, in Proc. Symposium on Algebraic Number Fields, 193-268, London: Academic Press 1977.
- [ 6 ] T. Hiramatsu, A formula for the dimension of spaces of cusp forms of weight one, to appear in Advanced Studies in Pure Math.

*Department of Mathematics  
Faculty of Science  
Kobe University  
Nada-ku, Kobe 657  
Japan*