

Correspondence

Edited by Kiriakos Xenitidis and
Colin Campbell

Contents

- Behavioural and psychiatric symptoms in people with dementia admitted to acute hospitals
- More data on speed of remission with ECT in geriatric depression
- The 'unknown' safety concern for aripiprazole once monthly

Behavioural and psychiatric symptoms in people with dementia admitted to acute hospitals

We would like to commend Sampson *et al*¹ on undertaking the difficult task of identifying and monitoring behavioural and psychological symptoms of dementia (BPSD) in people admitted to acute medical wards. The authors have also done their best to untangle the BPSD syndrome from similar clinical symptomatology seen in delirium, which still remains an ongoing conundrum for many of us working in liaison psychiatry. The study not only provides a wealth of information, but also raises a number of issues about how BPSD presentation might differ in older people when admitted to acute medical settings and how it influences their outcomes.

The authors used the Behavioral Pathology in Alzheimer's Disease scale (BEHAVE-AD),² which has been widely used to detect behavioural problems in people with dementia, in particular Alzheimer's disease. However, this scale has a number of shortcomings, neglecting some important symptoms of dementia, such as apathy, irritability and/or disinhibition,³ that are frequently present in dementia. The low–medium BPSD scores (2.6–4.4 mean BEHAVE-AD severity) are somewhat surprising, since the majority of the enrolled participants had higher Functional Assessment Staging Test (FAST)⁴ staging, corresponding to more advanced stages of dementia. Such low BPSD scores are usually associated with mild cognitive impairment.² Only five individuals coming from an 'other' place of residence (continuing NHS care?) had substantially higher BPSD severity scores. Similarly, the severity of the dementia (as measured via the FAST) did not influence BPSD scores, suggesting that either the medical problems modified the BPSD presentation, or the BPSD were pharmacologically managed. The only factor to have a significant effect on BPSD was presence of delirium, thus highlighting the difficulties in routine clinical settings of differentiating between BPSD and clinical symptoms of delirium.

Nevertheless, Sampson *et al*'s work undoubtedly indicates that hospitals make people with dementia worse, trebling their paranoia and delusional beliefs, causing hallucinations, making them more aggressive and disturbed, and substantially worsening their moods and anxieties. These findings support the public's widespread beliefs that hospitals are dangerous places, not only filled with sick people and germs, but with a wide potential for something to go amiss in lieu of wrong. And this 'wrong' ranges from having newly acquired diagnosis of dementia when physically unwell to worsening BPSD,¹ further complicating their polypharmacy and making them more frail and with poorer functional outcomes,⁵ as well as increasing their likelihood of death.¹ Not surprisingly, this also affects their formal and informal

caregivers. There is a striking discord between the severity of the recorded BPSD and the caregiver's distress, arguing that the problems around the escalating in-hospital behavioural changes are much more serious than the physical illness itself.

Many of the highlighted BPSD could be easily regulated with non-pharmacological approaches, including better orientation, information and knowing our patients. What is happening to make our hospitals more dementia-friendly? A number of hospitals have already introduced dementia-friendly wards that should be fully equipped with the professional experts in dementia care. However, surprisingly, there is a void of research evidence regarding how the newly introduced dementia-friendly policy in acute medical settings influences the behaviour of people with dementia. Since most of the people with BPSD come from residential and nursing care,¹ one wonders whether there should be another way of introducing dementia-friendly management. Perhaps an 'in-home health dementia care' approach would avoid (unnecessary) hospital admissions, and involve medical care professionals treating people with dementia in their own home whenever possible. This would reduce substantially not only the distress that both people with dementia and their caregivers face when in the acute medical setting, but also reduce hospital admissions, and thus result in substantial financial savings. The data from a recent study⁶ on people with advanced dementia stages prove that this can be successfully done, stressing that managing the distress, rather than behaviours that challenge, is central when treating people with dementia. We now need to take these lessons on board and implement them not only within our acute medical wards, but also adapt them to use in the community.

- 1 Sampson EL, White N, Leurent B, Scott S, Lord K, Round J, et al. Behavioural and psychiatric symptoms in people with dementia admitted to the acute hospital: prospective cohort study. *Br J Psychiatry* 2014; **205**: 189–96.
- 2 Reisberg B, Auer SR, Monteiro IM. Behavioral pathology in Alzheimer's disease (BEHAVE-AD) rating scale. *Int Psychogeriatr* 1996; **8** (suppl 3): 301–8.
- 3 Robert P, Ferris S, Gauthier S, Ihl R, Winblad B, Tennigkeit F. Review of Alzheimer's disease scales: is there a need for a new multi-domain scale for therapy evaluation in medical practice? *Alzheimers Res Ther* 2010; **2**: 24.
- 4 Reisberg B. Functional assessment staging (FAST). *Psychopharmacol Bull* 1988; **24**: 653–9.
- 5 Runganga M, Peel NM, Hubbard RE. Multiple medication use in older patients in post-acute transitional care: a prospective cohort study. *Clin Intervent Aging* 2014; **9**: 1453–62.
- 6 Treloar A, Crugel M, Adamis D. Palliative and end of life care of dementia at home is feasible and rewarding. Results from the 'Hope for Home' study. *Dementia* 2009; **8**: 335–47.

Elizabeta B. Mukaetova-Ladinska, Senior Lecturer in Old Age Psychiatry, Institute of Neuroscience, Newcastle University, Newcastle University Institute for Ageing and Newcastle Liaison Team for Older Adults, Tyne and Wear NHS Foundation Trust, Newcastle upon Tyne. Email: Elizabeta.Mukaetova-Ladinska@ncl.ac.uk. **Ann Scully**, Consultant in Old Age Psychiatry, Newcastle Liaison Team for Older Adults, Tyne and Wear NHS Foundation Trust, Newcastle upon Tyne.

doi: 10.1192/bjp.206.2.166

Author's reply: Mukaetova-Ladinska & Scully's comments on the challenges of conducting clinical research involving people with dementia on acute hospital wards¹ raise the issue of whether BPSD may present differently in the acute hospital.

In our study, those with more severe dementia were unwell, often bed-bound, and may have been less able to display BPSD (e.g. wandering or pacing). Additionally, 12% of our cohort was taking an antipsychotic (details available from the authors on request), and many of these patients had more severe dementia. We note the problem of recall bias in staff who may report problems that are harder to manage; for example, depression in