

¹Department Of Psychiatry, Clinical Psychology And Mental Health, La Paz University Hospital, Madrid, Spain and ²Idipaz, Department Of Psychiatry, Clinical Psychology And Mental Health, La Paz University Hospital, Madrid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.2083

Introduction: There are no studies which address the relationship between mindfulness and cognitive flexibility in interventions carried out online. This is the first study to examine the effect of two online mindfulness-based interventions on this cognitive function.

Objectives: To assess changes on cognitive flexibility after two online mindfulness-based group interventions in adult patients with anxiety disorders.

Methods: This study was carried out in a Mental Health Unit in Spain (Colmenar Viejo, Madrid). Thirteen adult patients (age mean = 51.69 years, ranging from 33 to 69 years, S.D. = 11.56) with anxiety disorders completed the interventions. The group treatments were Acceptance and Commitment Therapy and a Mindfulness-based Emotional Regulation intervention, during 8 weeks, guided by two Clinical Psychology residents. Both interventions were carried out online. The dependent variable was the score on the TMT-B (seconds). A comparison of paired-means was conducted. Statistical significance was set at $p < .05$.

Results: The normality assumption was met. Statistical power observed = 70.0%. The paired t-test showed statistically significant change between pre-treatment and post-treatment ($p = 0.019$; Cohen's $d = 0.75$), indicating improvement on cognitive flexibility.

Conclusions: These results show a statistically significant and medium/large effect-size change in cognitive flexibility after the two online interventions based on mindfulness. A larger sample size is required to confirm these results. Moreover, other studies need to examine the reliable change on this neuropsychological outcome.

Disclosure: No significant relationships.

Keywords: acceptance and commitment therapy; cognitive flexibility; Mindfulness-based Emotional Regulation; Online treatments

EPV0564

On gender and TMT-A. The REM-ACT study: Acceptance and commitment therapy versus a mindfulness-based emotional regulation intervention in anxiety disorders. A randomized controlled trial

E. Vidal-Bermejo¹, E. Fernández-Jiménez^{2*}, T. Castellanos-Villaverde¹, I. Torrea-Araiz¹, G. Navarro-Oliver¹ and A. Hospital-Moreno¹

¹Department Of Psychiatry, Clinical Psychology And Mental Health, La Paz University Hospital, Madrid, Spain and ²Idipaz, Department Of Psychiatry, Clinical Psychology And Mental Health, La Paz University Hospital, Madrid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.2084

Introduction: There is paucity of empirical studies which compare various mindfulness-based interventions on speed of visuomotor tracking and also analyse the differential effect of gender.

Objectives: To compare the effectiveness of Acceptance and Commitment Therapy (ACT) versus a Mindfulness-based Emotional Regulation (MER) intervention on speed of visuomotor tracking according to gender.

Methods: This study was carried out in a Mental Health Unit in Spain (Colmenar Viejo, Madrid). Firstly, 80 adult patients with anxiety disorders were randomized according to the score on the Acceptance and Action Questionnaire-II (blocking factor), of whom, 64 patients decided to participate (mean age = 40.66, S.D. = 11.43; 40 females). Each intervention was weekly, during 8 weeks, guided by two Clinical Psychology residents. A 2x2x2 mixed ANOVA (pre-post change x intervention type x gender) was conducted, with Sidak-correction post-hoc tests. The dependent variable was the score on TMT-A.

Results: Normality and homoscedasticity assumptions were met. No statistically significant differences were observed on age or gender between interventions. No statistically significant interaction effect was observed between pre-post change x intervention x gender on TMT-A [$F_{(1, 52)} = 2.867$, $p = .096$, statistical power observed = 38.3%]. However, simple effects were statistically significant: while males improved on TMT-A after MER ($p = .000$; Cohen's $d = 1.092$), females did so after ACT ($p = .000$; Cohen's $d = 1.506$).

Conclusions: These results show that gender moderates the improvement of the two mindfulness-based interventions examined on the speed of visuomotor tracking. More research is needed to confirm these findings.

Disclosure: No significant relationships.

Keywords: Mindfulness-based Emotional Regulation; speed of visuomotor tracking; acceptance and commitment therapy; randomized controlled trial

EPV0566

Online group psychotherapy during the “lock down”

D. Romac

Department Of Mental Health And Addiction Prevention, Teaching Institute for public health Dr Andrija Štampar, Zagreb, Croatia
doi: 10.1192/j.eurpsy.2021.2085

Introduction: The Covid-19 pandemic has limited the classic psychotherapy treatment. EAP provided temporary recommendations for online psychotherapy in March of 2020 which brought new possibilities and limitations and contains rules, ethics and techniques. From the perspective of a psychodynamic therapist, the specificity of online group psychotherapy in the context of strong stressors is described.

Objectives: During the “lockdown” period and the earthquakes that occurred simultaneously in Zagreb in March of 2020, the secure Internet platform enabled the continuity of the group's work in a video link modality.

Methods: Online group had regular weekly meetings which lasted 1.5 hours. The classic rules of group analytical therapy were adapted to the new setting in virtual space. A risk assessment was also performed.

Results: The six group members and therapist have connected online through more freedom, developed new levels of sensitivity, flexibility and creativity. They have also accepted limitations. The online group is able to provide holding, but deeper connections require a physical presence to exchange emotions that technology

interferes with as well as the process of empathizing. Previous live sessions crucial to maintaining emotional connections have served as reservoirs for a period in which communication over the Internet was insufficient.

Conclusions: Internet technology can temporarily enable the continuity of a group psychotherapy. Technical and institutional support is recommended. The advantages of technology can be used if the technique is adapted, realistic goals set, and clinical limitations accepted. Many questions about the possibilities of “online psychotherapy” are open and unexplored.

Disclosure: No significant relationships.

Keywords: group; online psychotherapy; “lock down”

EPV0570

The effect of motivational interview based on WhatsApp on the psychological domains of quality of life in infertile women with pcos: A randomized clinical trial

F. Ansari^{1*}, Z. Hamzehgardeshi¹, F. Elyasi², M. Moosazadeh³ and I. Ahmadi⁴

¹Nursing And Midwifery Faculty, Mazandaran University of Medical Sciences, Sari, Iran; ²Neurology, School Of Medicine Psychiatry And Behavioral Sciences Research Center Addiction Research Institutes Sari Imam Khomeini Hospital, Mazandaran University of Medical Sciences, Sari, Iran; ³Health Sciences Research Center Addiction Research Institutes, Mazandaran University of Medical Sciences, Sari, Iran and ⁴Department Of Obstetrics And Gynecology., School Of Medicine Sexual And Reproductive Health Research Center Sari Imam Khomeini Hospita, Mazandaran University of Medical Sciences, SARI, Iran

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.2086

Introduction: Polycystic ovary syndrome (Pcos) disease significantly decreased quality of life for women. Mental health is one of the components affecting the quality of life of these patients that attention to it is necessary to improve their quality of life.

Objectives: The present study was conducted to determine the effect of Motivational Interview Based on WhatsApp on the Psychological Domains of Quality of Life in Infertile Women with PCOS.

Methods: This randomized controlled clinical trial enrolled 60 Infertile Women with PCOS from the city of Sari-Iran in 2020. Participants were assigned to MI and control groups using block randomization. The intervention group received 5 weekly of MI online via WhatsApp. While the control group received only routine care. The psychological Domains score of quality of life in these individuals was measured using the quality of life questionnaire of polycystic ovary syndrome (MPCOSQ) before and after the intervention. Then, the data were entered into the SPSS software, version 25 and were analyzed using descriptive statistics, chi-square test, t-test, and repeated measures analysis of variance.

Results: No significant difference was observed between the two groups before the intervention mean The Psychological Domains scores ($p > 0.05$). After the intervention, mean (SD) of The Psychological Domains score was 34.8 (11.8) in the intervention group and 30.7 (11.6) in the control. No significant increasing in the mean between the two groups. The effect size (0.35) was calculated.

Conclusions: The results of the study showed that motivational interviewing is effective in improving the quality of life of women with pcos.

Disclosure: No significant relationships.

Keywords: Infertile women; polycystic ovary syndrome; motivational interview; quality of life; psychosis

EPV0571

An assessment of the content of discharge summaries at Mount Carmel Hospital, Malta

E. Camilleri*, G. Grech, Y. Farrugia and J. Padovani

Psychiatry, Mount Carmel Hospital, Attard, Malta

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.2087

Introduction: Discharge summaries are the mainstay of intra and inter-departmental communication, ensuring continuity of care. Local instructions fail to provide clear guidance to foundation doctors to ensure standardised discharge summaries.

Objectives: The audit aimed to assess the inclusion of information within discharge summaries at Mount Carmel Hospital, Malta. A secondary objective was to update the current online discharge summary framework.

Methods: Stratified random sampling was used to select 120 discharge summaries, issued between October 2018 and September 2019. These were chosen out of a total of 956 discharge summaries issued during the period. The inclusion of information was analysed against the National Standard for Patient Discharge Summary Information issued by the Health Information and Quality Authority, Ireland. Data was collected and grouped into seven categories each containing multiple data points.

Results: Patient details were present in all discharge summaries while no details relating to the primary care healthcare professional were documented. The average information inclusion rate regarding admission, discharge and medications was 85%. Average clinical information was documented in 50% while that of future management and person completing discharge summary was found in 41% and 28% respectively (as per Table 1). Encouragingly, discharge summaries contained mandatory information more frequently than conditional or optional information.

