

was lower only in the age group 30–40 years ($P < 0.05$), prolactin was lowered in the first month ($P < 0.05$), F.S.H. and L.H. were lower in the first month and first year ($P < 0.001$). Also F.S.H. and L.H. were lower in patients with Hypoactive sexual desire ($P < 0.01$).

Conclusions: hormonal therapy of sexual dysfunction may be tailored according to our findings using either androgens, bromocryptine or a combination of F.S.H. & L.H. according to the age, duration of the disorder or if hypoactive sexual desire is present.

Tues-P105

A SEXUAL BEHAVIOUR CORRECTION OF YOUNG PHYSICALLY HANDICAPPED MEN

V.G. Taktarov. *Moscow Medical Stomatological Institute, 103473, Moscow, Russia*

A comparison studies of the principal sexuality characteristics in physically handicapped men of 18–29 years old with motor disorders were performed in two groups. The first group was consisted of the invalids with inborn or early occurred motor lesions, namely, cerebral palsy (CP), spina bifida (SB) and herdoneurologic diseases (HND) mostly including myopathies. The second group involved the patients, who were invalidated after puberty in consequence of an amputation of extremities or different lesions of joints. Totally 96 patients were examined.

The peculiarities of libido, erection, ejaculation and orgasm were established by means of interview and questionnaire analysis. Furthermore we carried out a clinical examination of patients, during which the erectile function was estimated by nycturnal penial tumescence registration using Rigiscan-machine and by ultrasound flowmetry of penial arteries using simple or bidirectional dopplerography. The instrumental urodynamic methods served for an evaluation of a functional status of lower urinary tract.

The prominence of libido was not decreased in the great majority of examined persons. The sexual disorders in invalids of the second group were imaginary whereas in the first group the erectile dysfunction and ejaculatory problems were associated with organic abnormalities, particularly, in patients having HND and SB. Almost all members of both study groups showed high sexual activity and therewith the most part of patients preferred also physically defective persons as sexual partners.

It might be concluded that psychotherapeutic programmes is sufficient for a sexual behaviour correction of patients with gained invalidation. On the contrary a sexual rehabilitation of CP, HND and especially SB patients requires additional examinations and cooperative help of psychologists, andrologists and sexologists.

Tues-P106

PECULIARITIES OF SEXUAL DYSFUNCTION IN MEN OF DECLINING YEARS

M. Yagubov¹*, N. Kibrik¹. ¹*Moscow Research Institute of Psychiatry, Moscow, Russia*

The purpose of the investigation was a comparative assessment of sexual dysfunction in men of two age group: aged patients and patients in declining years.

148 aged men (from 46 to 74 years old) and 20 declining years men (from 75 to 89 years old) have been observed. The results of the investigation make it possible to come to the following conclusions:

1. For the moment of addressing for medical care the aged persons had twice as much protracted sexual dysfunction than the declining years patients.
2. Sexual function preservation and its psychological significance in the declining years patients depend predominantly on patient's sexual constitution.
3. Sexual problems differ in two groups: the declining years patients complains were less variative and concerned mostly erection problems and aspermatism.
4. The main feature of sexual dysfunction development in the declining years patients was absence of psychogenous and somatical factors of sexual function impairment.
5. Neurotic-like disturbances are typical in psychological background of sexual dysfunction in the declining age patients; psychological reactions are predominant in the aged persons.
6. Psychological capacity of sexual function in the declining years patients depends on marriage retaining.
7. Sexual function maintenance in the declining years patients correlates with active way of life, work and social activity, physical exercises.

Tues-P107

AGING PECULIARITIES OF DEVELOPMENT OF THE SEXUAL DYSFUNCTIONS

N.D. Kibrik¹. ¹*Moscow Research Institute of Psychiatry Moscow, Russia*

During research of the age factor in clinic of sexual disorders, it is necessary to find a reason of disease, and attempt of rational explanation of the reason of it own sexual insolventation. For young age (before 25), important rule in development of sexual dysfunction's play such moments, as presence of the psycho-physical infantilism, hysterical and epileptic marks of character, violation of the temps of sexual ripen. In young age violations of sexual functions are combined with hypochondriac manifestations, deep affective experience with anxiety, fear, ideas of masturbation hypochondria, self accusation, self humiliation. In middle age group (25–55 years old) maximal important became situation factors, related with matching of sexual function. In the basis of ideas about presence of sexual dysfunction lies wrong evaluation of present sexual possibilities which relate with age changes of rate of sexual manifestations. Disharmonical relations in family, more often cause by psychotraumatic influence of sexual partner. In response to psychotrauma there is development one-sided exaggerated in affect reaction. In basis of such acute developed, situational provoked reaction lies feeling of loneliness, accompanied by thoughts of self uselessness, low importance which related with absence understanding from surround people, distance from the family, all those things could be reviled during deep clinical-psychological investigation. In elder age group (after 55) there is prevalence of persons with epileptic and infantile-primitive features of character. In person characteristic we can see such features as sensitiveness, hypotimness with high level of claims, pessimistic attitude to life. The age reduce of sexual possibilities, loss of usual stereotype of sexual manifestations and related with it loss of self conceit in prestigious for this type of person sphere, all this results in development of steady depressive states with loosing of life perspective, sense and aim of life. Definite importance in this have situational circumstances, such as loneliness, related with the death of one of partners in marriage, somatic diseases, especially in cardiovascular and urogenital spheres, which hinder from manifestation of sexual activity and causing socio-sexual isolation.