

BECOME A LIFE MEMBER AND YOU'LL JOIN PRESTIGIOUS COMPANY



Become a Life Member of Fauna & Flora International to show an extraordinary commitment to an organisation that has been at the forefront of conservation for over 100 years.

Since its inception in 1903 as the world's first international conservation organisation Fauna & Flora International has been working tirelessly for the preservation of plants and animals across the world. For over 100 years Life Members have provided the support and commitment that has helped us to achieve so much, not least key achievements such as:

- Reintroducing the Arabian oryx to the wild
- Establishing the International Gorilla Conservation Programme
- Establishing Vietnam's first locally managed conservation organisation

Collaboration is key to our approach, and wherever possible we work with other global conservation organisations to ensure we are effective and efficient. That's why in 2015 we moved into the David Attenborough Building of the Cambridge Conservation Initiative with several other global conservation groups. Now we are asking you to be part of our wider collaborative work by becoming a Life Member.

As a Life Member you will receive Oryx—The International Journal of Conservation and our annual magazine Fauna & Flora, and you will also be invited to special events, where you can network with some of the world's leading conservationists. You will be joining a select group of supporters who have shown an extraordinary commitment to international conservation.

By joining Fauna & Flora International as a Life Member with a one-off payment of £1,500 you will be making a genuine difference to our conservation work and will forever be part of our global conservation organisation.



Gary Morrisroe/FFI

"I have been a member of Fauna & Flora International since the 1950s... investment in the work of FFI is truly an investment in the future of our planet"

Sir David Attenborough

To join as a Life Member, you can:

Call us on +44 (0)1223 749 019

Email us at members@fauna-flora.org

Visit www.fauna-flora.org/life-membership



751 Primate conservation—new reports from the field,
A.B. Rylands, R.A. Mittermeier & E.A. Williamson
 753 Briefly
 759 Conservation news

Primate conservation

763 New estimates of orangutan density in Sungai Palin watershed, Kapuas Hulu, West Kalimantan, Indonesia, *A. Yanuar, J. Fehse, D. Chivers & J. Sugardjito*
 767 Identifying priority areas for the conservation of the Critically Endangered northern white-cheeked gibbon *Nomascus leucogenys* in northern Lao, *K. Syxaiyakhathor, D. Ngoprasert, N. Asensio & T. Savini*
 776 Transboundary conservation of the last remaining population of the cao vit gibbon *Nomascus nasutus*, *C.-Y. Ma, H. Trinh-Dinh, V.-T. Nguyen, T.-D. Le, V.-D. Le, H.-O. Le et al.*
 784 Using occupancy-based camera-trap surveys to assess the Critically Endangered primate *Macaca nigra* across its range in North Sulawesi, Indonesia, *C.L. Johnson, H. Hilser, M. Linkie, R. Rahasia, F. Rovero, W. Pusparini et al.*
 794 Potential distribution of and priority conservation areas for the Endangered Caatinga howler monkey *Alouatta ululata* in north-eastern Brazil, *R. Freire Filho & J.M. Palmeirim*
 803 Habitat suitability for primate conservation in north-east Brazil, *B. Moraes, O. Razgour, J.P. Souza-Alves, J.P. Boubli & B. Bezerra*
 814 First record of the Peruvian yellow-tailed woolly monkey *Lagothrix flavicauda* in the Región Junín, Peru, *S.M. McHugh, F.M. Cornejo, J. McKibben, M. Zarate, C. Tello, C.F. Jiménez & C.A. Schmitt*
 819 Range and conservation updates for the Critically Endangered blue-eyed black lemur *Eulemur flavifrons* and the Vulnerable black lemur *Eulemur macaco*, *J.C. Tinsman, C.L. Eschmann, J.S. Solofondranohatra, J.R. Ralainirina, M. Holderied & G. McCabe*

Carnivore conservation

828 Effects of habitat alteration and disturbance by humans and exotic species on fosa *Cryptoprocta ferox* occupancy in Madagascar's deciduous forests, *S.D. Merson, L.J. Dollar, C.K.W. Tan & D.W. Macdonald*
 837 Spatial dynamics and activity patterns of the fosa *Cryptoprocta ferox* in Ankarafantsika National Park, Madagascar: carnivores navigating a human-influenced landscape, *E. Wyza, L. Dollar, L.P. Rahajanirina, V. Popescu & N.J. Stevens*

847 Corridors of tolerance through human-dominated landscapes facilitate dispersal and connectivity between populations of African lions *Panthera leo*, *S. Dolrenry, L. Hazzah & L. Frank*
 851 Updated ranges of the Vulnerable cheetah and Endangered African wild dog in Angola, *P. Monterroso, F. Rocha, S. van Wyk, T. António, M. Chicomo, S. Kosmas et al.*
 854 Priority areas for jaguar *Panthera onca* conservation in the Cerrado, *M.P. Portugal, R.G. Morato, K.M.P.M. de B. Ferraz, F.H.G. Rodrigues & C.M. Jacobi*
 866 Factors influencing local attitudes towards the conservation of leopard cats *Prionailurus bengalensis* in rural Taiwan, *I. Best & K.J.-C. Pei*
 873 Conservation status of the dhole *Cuon alpinus* in north-east India, with a focus on Dampa Tiger Reserve, Mizoram, *P. Singh, A. Srivathsa & D.W. Macdonald*
 878 New insights into Sulawesi's apex predator: the Sulawesi civet *Macrogalidia musschenbroekii*, *I. Hunowu, A. Patandung, W. Pusparini, I. Danismend, A. Cahyana, S. Abdullah et al.*
 882 Land-cover change threatens tropical forests and biodiversity in the Littoral Region, Cameroon, *M.I. Mahmoud, M.J. Campbell, S. Sloan, M. Alamgir & W.F. Laurance*
 892 Ecology and conservation of the endemic Bawean warty pig *Sus verrucosus blouchi* and Bawean deer *Axis kuhlii*, *E.J. Rode-Margono, H. Khwaja, M. Rademaker & G. Semiadi*
 901 Camera trapping reveals a diverse and unique high-elevation mammal community under threat, *R. Pillco Huarcaya, C. Beirne, S.J. Serrano Rojas & A. Whitworth*
 909 Acknowledgements

Cover Monitoring populations of conservation concern is much improved if baseline data are available. This is of particular importance for Critically Endangered taxa such as the Bornean orangutan. A survey of orangutan nests in the peat-swamp forests of the Sungai Palin watershed, of the upper Kapuas river, West Kalimantan, Indonesia, has estimated the density of this subspecies in the unprotected area of Sungai Palin and compared this to data from the nearby area of Nanga Awen, which was surveyed in 1991. For further details, see pp. 662–665. (Photograph © Ondrej Chvatal/Shutterstock)