

**Results:** Valerian activity on sleep disturbances has been attributed to the presence of isovaleric acids and valepotriates with reported calming action and GABA reuptake inhibition with sedative effects. Considering the data presented in the literature, despite controversial and conflicting, several studies showed that valerian (160-600mg/day) improved sleep quality and reduced sleep latency and duration; also valerian seems more effective for chronic insomnia than acute episodes.

Hop has different properties: calming, sleep inducing, gastric secretion stimulating and spasmolytic.

Increasing GABAergic activity seems to be the main mechanism of action, thus inhibiting the central nervous system and also has demonstrated binding affinities to some of the melatonin and serotonin receptor. Its sedative characteristics have been confirmed in a clinical trial in association with valerian, where sleep latency and quality were improved. However, monotherapy studies showed no relevant effectiveness in sleep.

Kava Kava plant showed promising results, in rats and humans, with decrease sleep latency, better sleep quality and recuperation after sleep. However, raised concern about its potential of hepatotoxicity.

There is also promising evidence of the lavender efficacy for sleep disorders in a wide variety of populations and diseases, it was actually mentioned to be as effective as lorazepam in adults with anxiety and sleeping problems. With studies with dose of 80mg it was observed a reduction in sleep awakenings, sleep duration and overall sleep quality and anxiety.

**Conclusions:** There is a clear preference from the patient to natural compounds, and with almost nonexistent side effects, some herbal derivatives are showed to have positive effectiveness in mild insomnia, but nonetheless much more studies on this field are needed.

**Disclosure of Interest:** None Declared

## EPV1024

### Sleep disorders among university students as underestimated mental health problem

E. L. Nikolaev<sup>1</sup>, T. Nikolaeva<sup>2</sup> and M. Alhasan<sup>2\*</sup>

<sup>1</sup>Department of Social and Clinical Psychology and <sup>2</sup>Medical Faculty, Ulianov Chuvash State University, Cheboksary, Russian Federation

\*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1609

**Introduction:** The effect of sleep disorders on the weakening of the students' mental health potential is still underestimated. Students might not openly complain of having problems with sleep, considering them insignificant. Nevertheless, sleep disorders may be the sign of actual or developing mental health problems.

**Objectives:** To reveal the prevalence of parasomnic and insomnic disorders in university students, who do not have health related complains.

**Methods:** We surveyed 77 first and second-year students of both genders by means of a questionnaire that included questions describing the signs of various sleep disorders.

**Results:** One third of the students revealed having parasomnic disorders in the form of dissociated sleep states – 35.1% of the respondents talk in sleep (states of somniloquy or sleep talking), 6.5% get seated on their beds, 5.2% get up from their beds (states of partial awakenings and confusional arousals), 5.2% walk around the room or house (sleepwalking, or somnambulism). Over half of the students experience night phobias (53.2%), 2.6% out of them experience them constantly. Some students' fears grow into nightmares. Half of the respondents (50.6%) state they very rarely see nightmares. Every fifth student (20.8%) sees nightmares only from time to time. 10.4% of the students see them very often or constantly. Over half of the respondents (55.8%) complain of insomnic disorders in the form of insomnia. 3.9% of them experience it constantly, 10.4% – often, 16.9% – sometimes, and 24.7% – rarely

**Conclusions:** The frequency of sleep disorders in students is very high. Consequently, it is important to inform university students timely about potential risks and ways to avoid them.

**Disclosure of Interest:** None Declared

## EPV1025

### How effective is ketogenic diet in sleep disorders ?

N. Kouki

outpatient and emergency department, razi hospital, tunis, Tunisia  
doi: 10.1192/j.eurpsy.2024.1610

**Introduction:** Sleep disorders vary widely and its treatment are based on a combination of life style changes and pharmacological therapy adapted to the primer health issue. Ketogenic diet has shown not only its efficacy in different health conditions, but it is also becoming a popular health trend. Could the therapeutic spectrum of ketogenic diet cover sleep disturbances ?

**Objectives:** The aim of our study is to evaluate the effect of ketogenic diet on sleep disorders

**Methods:** To identify relevant studies ,our literature review was based on the Pubmed interface and adapted for 2 databases : science direct and google scholar. We used the following key words (ketogenic diet [meSH terms]) and (sleep disorders [meSH terms]).

**Results:** Our research revealed 14 articles published between 2012 and 2022. We selected 8 which corresponded to the purpose of our review. The ketogenic diet affects sleep hemostasis indirectly. In fact, this diet is associated with weight loss and therefore reduction of metabolic and cardiovascular complications disturbing sleep quality. From a neurobiological perspective, this regimen based on limited carbohydrates is associated with a low Tryptophan intake which is the precursor of melatonin. But on the other hand, Ketone bodies trigger adenosine activity which promotes melatonin liberation, the sleep inducing hormone.

**Conclusions:** ketogenic diet modulates melatonin activity therefore affects sleep architecture. Meanwhile, Its impact on sleep disorders is still controversial due to the variation of its pathophysiological mechanisms.

**Disclosure of Interest:** None Declared