

Aims Methodological approach to a CVD risk screening, risk stratification and specific life-style interventions development is presented. In a context of the psychiatric hospital with c. 7000 admissions per year.

Methods There are no concise data on CVD risk of psychiatric patients in the Czech Republic so cross-sectional analysis of one day hospitalized patients was performed.

Results A sample of 1056 pts. was obtained. Database allowed extraction data on CVD risk factors (RF): diagnosis, age, sex, BMI and blood pressure (BP). The most common diagnosis were F20 and F10. Multicriterial analysis according to diagnosis (frequency of highest BMI and BP) showed the worst results in the F20 followed by F10 group. Would we define the CVD RF as BMI \geq 30, age \geq 65, sBP \geq 140, dBP \geq 90, then no RF is present in 368, one in 238, two in 191, three in 92 and four RF in 33 pts. Two step screening protocol was developed – SOMA score. Variables and cut-offs for positivity were set based on the results.

Conclusion Methodological process of SOMA score screening is presented as well as consecutive health care interventions.

Disclosure of interest The authors have not supplied their declaration of competing interest.

Acknowledgement The study was realized in a framework of SOMA project, Program CZ11 Initiatives in public health care, Norway grants, NF-CZ11-OV-2-030-2015.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.763>

EW647

Kynurenine pathway metabolites and suicide attempters among psychiatric inpatients: Preliminary results

M. Pompili^{1,*}, D. Erbuto², A. Forte², F. Montebovi², L. Lionetto², P. Girardi²

¹ Rome, Italy

² Sapienza University of Rome, Department of Neurosciences-Mental Health and Sensory Organs, Rome, Italy

* Corresponding author.

Introduction Increased concentrations of kynurenine (KYN) pathway metabolites have been associated to several groups of psychiatric disorders. As for suicide risk literature is still inconclusive due to the limited evidence.

Objectives to test the hypothesis of Increased concentrations of kynurenine as related to suicide attempter.

Aims We aim to investigate the association between kynurenine pathway metabolites blood levels and suicidal behaviour, in affective disorder patients, in order to explore if kynurenine pathway metabolites could be potential diagnostic biomarkers.

Sample We enrolled a sample of affective disorder patients and perform detailed diagnoses, as well as detailed assessment of suicidal behaviour using validated questionnaires. We also aim to follow-up individuals included in the current study.

Methods Plasma KYN was assayed by high performance liquid chromatography in three groups: healthy volunteers ($n=90$), patients with mood disorders with a recent suicide attempt ($n=44$) and without ($n=44$) history of suicide attempt. Analysis of variance tested for group differences in KYN levels. Each was evaluated with psychometric scales. Patients were sampled for 10 cc of venous blood for assay. The preparation of blood samples and assay was processed by a specialist using high performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS).

Results Preliminary results will be presented as to shed light if KYN levels differed across groups. According to preliminary calculations we expect that KYN is higher in suicide attempters compared with non-attempters, who did not differ from healthy volunteers.

Conclusions Our work-in-progress study suggests that KYN and related molecular pathways may be implicated in the precipitation of suicidal behavior.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.765>

EW648

The psychiatry recruitment crisis across Europe: Evaluation by the European Federation of psychiatric trainees

H. Ryland^{1,*}, F. Baessler², M. Casanova Dias³, L. De Picker⁴, M. Pinto Da Costa⁵, A. Kanellopoulos⁶, E. Sonmez⁷, P. Alfimov⁸, D. Sebbane⁹, S.M. Birkle¹⁰

¹ West London Mental Health Trust, Broadmoor Hospital, London, United Kingdom

² University of Heidelberg, Department of Psychiatry, Heidelberg, Germany

³ Camden and Islington NHS Foundation Trust, General Adult Psychiatry, London, United Kingdom

⁴ University of Antwerp, Collaborative Antwerp Psychiatric Research Institute, Antwerp, Belgium

⁵ University of Porto, Hospital de Magalhães Lemos, Porto, Portugal

⁶ National & Kapodistrian University of Athens, Evgenidion Therapeftirion, Athens, Greece

⁷ Marmara University, Department of Psychiatry, Istanbul, Turkey

⁸ Moscow Research Institute of Psychiatry, Department of Psychiatry, Moscow, Russia

⁹ University of Lille, Lille, France

¹⁰ Ruhr-University Bochum, LWL-University Hospital for Child and Adolescent Psychiatry, Hamm, Germany

* Corresponding author.

Introduction Recruitment of medical students and junior doctors in to psychiatry is a long-standing concern in many countries, with low proportions of medical graduates choosing it as a specialty and ongoing stigma from within the medical profession. In some countries the reverse problem is the case, with too many doctors wishing to enter psychiatry, and insufficient training places available.

Objectives To understand the current situation within Europe with regards to recruitment in to psychiatry and to identify existing recruitment initiatives.

Methods The European Federation of Psychiatric Trainees conducts an annual survey of all member organisations. A delegate of each national association of psychiatric trainees is asked to identify if their country has a problem with recruitment and if so, whether there were too many or too few applicants for training places. Delegates from countries with recruitment initiatives were contacted to provide further details.

Results In 2014, a total of 31 countries completed the survey, with 17 stating that too few medical practitioners choose psychiatry. In total 8 countries with recruitment problems reported that initiatives exist to encourage doctors to enter psychiatric training. Of these, 7 responded to describe the initiatives, which included national recruitment strategies, financial incentives, careers fairs, mentoring schemes and a whole host of other projects.

Conclusions Recruitment in to psychiatry remains a serious problem in a significant proportion of European countries, but a wide range of initiatives exist which aim to combat this shortfall. It will be important over the coming years to establish which initiatives are most effective at increasing recruitment.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.766>