

compulsive buying. Twenty outpatients with a lifetime diagnosis of kleptomania and ten suffering from compulsive buying were evaluated by means of the Structured Clinical Interview for DSM-IV, of a specially-designed semi-structured interview and of a modified version of the Family History Research Diagnostic Criteria. The majority of patients reported an early and abrupt onset, with an episodic course of the disorder with no gender prevalence. Lifetime comorbidity for other Axis I disorders was relevant, in particular for mood disorders, obsessive-disorder (OCD), separation anxiety, panic disorder and OCD-related disorders such as pathological gambling and tricotillomania. Family history showed a high prevalence of mood disorders, alcohol abuse and OCD. Our study indicated a clear connection between kleptomania, mood disorders and OCD, the exact nature of which has yet to be clarified.

## S04. Behaviour and motor control in psychiatric disorders

*Chairs:* M.R. Lemke (D), B.G.C. Sabbe (NL)

### S04.01

#### IMPULSIVITY, BEHAVIORAL DYSCONTROL, AND CONSCIOUS AWARENESS

E.S. Barratt, *Department of Psychiatry, The University of Texas Medical Branch, Galveston, TX, 77555-0189, USA*

Impulsivity has emerged as one of the leading risk factors for behavioral disorders including, for example, substance abuse, ADHD, borderline and conduct/antisocial personality disorders. Impulsive acts have been defined clinically as "occurring suddenly, quickly, without much forethought, and without consideration of consequences" (Granit). Theoretical causal models of impulsivity emphasize arousal, varying forms of information processing, attention, and sensitivity to reinforcement in learning paradigms. Barratt and colleagues have proposed that impulsivity is related to temporal information processing which is related to performance on timing and rhythm tasks. In construct validity studies they have demonstrated that self-report measures of motor impulsivity, performance on a wide range of behaviors involving timing and rhythm requirements (verbal tasks [e.g., reading], pursuit rotor and finger tapping) and selected cognitive psychophysiological measures related to behavioral inhibition ( $N_{200}$ ) and information processing ( $P_{300}$ ), converge to define impulsivity. In predictive validity studies they have demonstrated that these measures are significantly related to ADHD, conduct disorder and impulsive aggression. This paper will discuss proposed neural circuits (e.g., basal ganglia, thalamus, and frontal lobes) which may be related to the behavioral measures of impulsivity and will include a discussion of a scale of conscious awareness which has impulsivity and premeditated behaviors at opposite extremes. Current research relating the lateralized readiness potential to impulsivity will also be broached.

### S04.02

#### A NEW MODEL TO ASSESS STIMULUS PROCESSING AND BEHAVIOR CONTROL IN HUMANS FOLLOWING EXPERIMENTAL INDUCTION OF EMOTIONS

M.R. Lemke\*, T. Wendorff, G. Fritzer, C.J. Fischer, Z. Rupp, S. Tetzlaff, *Department of Psychiatry and Psychotherapy, University of Kiel, Kiel, Germany*

**Objective:** Many psychiatric disturbances including borderline and antisocial personality disorders, substance abuse, eating disorders and suicidal behavior present with dysfunction of processing stimuli into adequate actions. Perceptive and executive components of this process may be modulated by affective stimuli. It was hypothesized that stimuli processing and generation of movements are affected by induction of positive and negative emotions.

**Methods:** Healthy subjects were subjected to visual stimuli which they could turn off by releasing and pressing two buttons. Neurophysiological methods including EEG, startle reflex, EMG and kinematic measures of hand movements by infrared detection (Proflex) were used to analyze the neuronal process from stimulus perception to movement execution with a specific software program for continuous chronological assessment of the signals with high precision.

**Results:** Stimuli (International Affective Picture System) induced different affective valence ratings (neutral/positive/negative) and startle response amplitudes (neutral  $48.85 \pm 3.28$ , positive  $49.95 \pm 4.14$ , negative  $52.04 \pm 8.95$ ) Movement analysis revealed differences in onset of movement (neutral  $4.86 \pm 1.56$ , positive  $5.65 \pm 2.12$ , negative  $4.73 \pm 1.32$  sec), movement duration (neutral  $1.48 \pm 0.59$ , positive  $1.49 \pm 0.42$ , negative  $1.36 \pm 0.44$  sec), max. velocity (vmax) (neutral  $0.45 \pm 0.13$ , positive  $0.46 \pm 0.15$ , negative  $0.46 \pm 0.13$  m/s), vmax latency (neutral  $37.88 \pm 17.18$ , positive  $39.87 \pm 14.35$ , negative  $36.11 \pm 12.7$  sec).

**Discussion:** We showed for the first time that not only perceptive, but also executive components of CNS behavior control can be experimentally modulated by affective stimuli. The efficacy of psychotherapeutic and psychopharmacological interventions on behavioral control can be evaluated using our model. Future studies will include experimental variation of serotonergic CNS activity and its effects on behavior control.

### S04.03

#### ANXIETY/AGGRESSION-DRIVEN DEPRESSION: A MANIFESTATION OF SEROTONERGIC PATHOLOGY?

H.M. van Praag

No abstract was available at the time of printing.

### S04.04

#### FACIAL EXPRESSIONS AND PERSONALITY

G. Juckel\*, M. Vogel, A. Präßl, B. Graf, P. Mavrogiorgou, U. Hegerl, *Dept. of Psychiatry, Ludwig-Maximilians-University Munich, Germany*

Facial expressions are often disturbed in psychiatric patients. Subjects with personality styles such as sensation seeking, extraversion or impulsivity are thought to have a risk to develop psychiatric disorders. To identify those subjects, kinematic analysis of facial expressions could be helpful. An active measurement device was used, allowing kinematic analysis of facial movements in detail. Markers which are fixed in distinct points of the face and send light or ultrasonics in high frequency give a direct measure of facial movements with high spatial-temporal resolution. Healthy