

# Seed Dormancy and Seed Morphology Related to Weed Susceptibility to Biofumigation – CORRIGENDUM

## Corrigendum

**Cite this article:** Lefebvre M, Leblanc ML, Watson AK (2019) Seed Dormancy and Seed Morphology Related to Weed Susceptibility to Biofumigation – Corrigendum. *Weed Sci.* **67**: 474. doi: 10.1017/wsc.2018.17

---

Maxime Lefebvre, Maryse L. Leblanc and Alan K. Watson

---

In the original publication of this article, quantities of dried Indian mustard tissues in the petri dish germination tests were reported in  $\text{mg cm}^{-2}$ . The calculations were based on a 10 cm diameter petri dish.

However, the petri dishes had precisely 9.11 cm in diameter. Consequently, the values of all further references to rates in  $\text{mg cm}^{-2}$ , quantities of ITCs reported by  $\text{cm}^{-2}$ , to the **b** coefficient parameter of the nonlinear response (Table 2), to the coefficient and intercept of the linear regression analyses (Table 5) in regard to  $\text{EC}_{50}$  and  $\text{LD}_{50}$  should be multiplied by a factor of 1.205. This does not alter the analyses nor conclusions.

The authors apologize for the error.

## Reference

Lefebvre M, Leblanc ML, Watson AK (2018) Seed Dormancy and Seed Morphology Related to Weed Susceptibility to Biofumigation. *Weed Sci* 66:199–214. doi: 10.1017/wsc.2017.66