

of the six patients who survived hospitalization. All case-patients, but no non-case-patients, were exposed to 11.5-year-old cellulose acetate dialyzers (all of these dialyzers were discarded by the hospital before the investigation). Laboratory investigation of field-retrieved 0- to 13.6-year-old dialyzers of similar type indicated significant chemical degradation in the older membranes. In vivo injection of extracts of membrane degradation products produced iritis and hemorrhages in rabbits' eyes.

It was concluded that severe patient injury was associated with exposure to aged cellulose acetate membranes of dialyzers, allowing cellulose acetate degradation products to enter the blood. Clinicians should be aware that aged cellulose acetate membranes might cause severe adverse reactions.

FROM: Hutter JC, Kuehnert MJ, Wallis RR, Lucas AD, Sen S, Jarvis WR. Acute onset of decreased vision and hearing traced to hemodialysis treatment with aged dialyzers. *JAMA* 2000;283:2128-2134.

Molecular Epidemiology of *S epidermidis* in an NICU

Coagulase-negative staphylococci, especially *Staphylococcus epidermidis*, are increasingly important nosocomial pathogens, particularly in critically ill neonates. Villari and coinvestigators, from the Department of Health and Preventive Sciences, University Federico, Naples, Italy, conducted a 3-year prospective surveillance of nosocomial infections in a neonatal intensive care unit (NICU) using traditional epidemiological methods, as well as molecular typing of microorganisms. The objectives of the study were (1) to quantify the impact of *S epidermidis* on NICU-acquired infections; (2) to establish if these infections are caused by endemic clones or by incidentally occurring bacterial strains of this ubiquitous species; (3) to evaluate the use of different methods for the epidemiological typing of the isolates; and (4) to characterize the occurrence and the spread of staphylococci with decreased glycopeptide susceptibility.

The results showed that *S epidermidis* is one of the leading causes of NICU-acquired infections and that the reduced glycopeptide susceptibility, if investigated by appropriate detection methods such as population analysis, is more common than is currently realized. Typing of isolates, which can be performed effectively through molecular techniques such as pulsed-field gel electrophoresis but not through antibiograms, showed that many of these infections are due to clonal dissemination and thus are potentially preventable by strict adherence to recommended infection control practices and the implementation of programs aimed toward the reduction of the unnecessary use of antibiotics. These strategies are also likely to have a significant impact on the frequency of the reduced susceptibility of staphylococci to glycopeptides, since this phenomenon appears to be determined either by more resistant clones transmitted

from patient to patient or, to a lesser extent, by strains that become more resistant as a result of antibiotic pressure.

FROM: Villari P, Sarnataro C, Iacuzio L. Molecular epidemiology of *Staphylococcus epidermidis* in a neonatal intensive care unit over a three-year period. *J Clin Microbiol* 2000;38:1740-1746.

Growth of *S epidermidis* and *P aeruginosa* on Biomedical Polymers

The infection risk of biomaterials implants varies between different materials and is determined by an interplay of adhesion and surface growth of the infecting organisms. Gottenbos and colleagues from the University of Groningen, The Netherlands, conducted a study that compared initial adhesion and surface growth of *Staphylococcus epidermidis* HBH(2) 102 and *Pseudomonas aeruginosa* AK1 on poly(dimethylsiloxane), Teflon, polyethylene, polypropylene, polyurethane, poly(ethylene terephthalate), poly(methyl methacrylate), and glass. Initial adhesion was measured in situ in a parallel plate flow chamber with microorganisms suspended in phosphate-buffered saline, while subsequent surface growth was followed in full and in 20 times diluted growth medium. Initial adhesion of both bacterial strains was similar to all biomaterials. In full growth medium, generation times of surface growing *S epidermidis* ranged from 17 to 38 minutes with no relation to wettability, whereas in diluted growth medium, generation times increased from 44 to 98 minutes with increasing surface wettability. For *P aeruginosa*, no influence of surface wettability on generation times was observed, but generation times increased with decreasing desorption rates, maximal generation times being 47 minutes and minimal values down to 30 minutes. Generally, generation times of adhering bacteria were shorter than of planktonic bacteria.

The authors concluded that surface growth of initially adhering bacteria is influenced by biomaterials surface properties to a greater extent than initial adhesion.

FROM: Gottenbos B, van der Mei HC, Busscher HJ. Initial adhesion and surface growth of *Staphylococcus epidermidis* and *Pseudomonas aeruginosa* on biomedical polymers. *J Biomed Mater Res* 2000;50:208-214.

Catheter-Associated UTIs Are Rarely Symptomatic

Catheter-associated urinary tract infection (CA UTI) is the most common nosocomial infection, accounting for more than 1 million cases each year in US hospitals and nursing homes. Up to one half of the patients requiring an indwelling urethral catheter for 5 days or longer will develop bacteriuria or candiduria. Silent catheter-associated bacteriuria comprises a huge reservoir of antibiotic-resistant organisms in the hospital, particularly on the critical-care unit. Although there have been recommen-