

15435

THE JOURNAL
OF
LARYNGOLOGY
AND
RHINOLOGY;

*AN ANALYTICAL RECORD OF CURRENT LITERATURE RELATING TO
THE THROAT AND NOSE.*

EDITED BY

SIR MORELL MACKENZIE, M.D. Lond.

AND

R. NORRIS WOLFENDEN, M.D. Cantab.

WITH THE CO-OPERATION OF

DR. G. HUNTER MACKENZIE (EDINBURGH)
DR. FAUVEL (PARIS)
DR. JOAL (MONT DORE)
PROF. MASSEI (NAPLES)
PROF. GUYE (AMSTERDAM)
DR. CAPART (BRUSSELS)
DR. CONSTANTIN KARWOWSKI
(WARSAW)

DR. MICHAEL (HAMBURG)
PROF. RAMON DE LA SOTA Y LASTRA
(SEVILLE)
DR. HOLGER MYGIND (COPENHAGEN)
DR. SMYLY (DUBLIN)
DR. DAVID COLLINGWOOD (SYDNEY)
DR. G. W. MAJOR (MONTREAL)
DR. MAXWELL ROSS (EDINBURGH)

AND

Dr. JOHN N. MACKENZIE
for America.

VOL. II.—1888.

LONDON:
R. ANDERSON & CO.,
14, COCKSPUR STREET,
CHANCING CROSS, S.W.

PHILADELPHIA:
P. BLAKISTON, SON & CO.,
1012, WALNUT STREET.

CONTENTS.

Original Article, p. 223.—MAYO COLLIER: The Epiglottis: Its Attachments and Relations to Surrounding Structures.

New Instruments and Therapeutics, p. 224.—CARPENTER: Lamp for Examination. POTTER: Nasal Scissors. YEO: Address on Therapeutics of Uric Acid Diathesis. PARKER: Incompatibility in Pharmacy. TREVES: Treatment of Carotid Hæmorrhage. COCHRANE: Ergotism in Hæmoptysis. ALLEN STURGE: Antipyrin; a warning. EDMUNDS: Cocaine. POTTER: Curious Effect of Cocaine. MOREAU AND COCHEZ: Hydrofluoric Acid for Diseases of Respiratory Passages. THORNER: Salol. WEILL: Carbonic Acid for Dyspnoea. SCHUSTER: Bergeon's Clysmata of Carbonic Acid, etc., in Diseases of Lungs. SPENCER: Antiseptic Treatment of Phthisis. WAGNER: Colorado Springs and Davos Platz. THOMSON: Winter Climate of Florence.

Diphtheria, p. 228.—BRISTOWE: Diphtheritic Paralysis. LEGROUX: Diphtheria and Creosote. MOHAMMED: Treatment of Diphtheria. BEAUPÈRE: Chloroform Anæsthesia in Tracheotomy.

Nose and Naso-Pharynx, p. 231.—LERMOYER: Hay Asthma. COUPARD: Foreign Bodies in Right Nostril. POTTER: Tuberculosis of Nose, Mouth, and Pharynx. HINKEL: Irregularities of Septum Narium. SEILER: Echinodroses of Septum Narium. GRADLE: Ocular Symptoms in Nasal Disease. SCHMIDT: Bilateral Amaurosis. CORIVEAUD: Choreiform Convulsions of Reflex Origin. STONHAM: Naso-Pharyngeal Polypus. FONTAN: Temporary Maxillo-malar Resection. HERMET: Cauterization of Posterior Region of the Turbinate Bodies. POESCHEL: Diseases of the Naso-pharynx.

Mouth, Tonsils, Pharynx, &c., p. 234.—BROCA: Hare-lip. LLOYD: Composite Odontome of the Upper Jaw. QUENU: Maxillary Fistula. HADDEN: Suppression of the Salivary and Buccal Secretions. ROWLANDS: Permanent Suppression of the Salivary Secretion. BUNTON: Ibid. COOPER: Communicability of Syphilis through Saliva. PHILIP: Contribution towards the Etiology of Phthisis. VERCHÈRE: Treatment of Ranula. GUINARD: Congenital Ranula. WHITEHEAD: Excision of the Tongue. GILLOT: Ampullary Dilatations of the Tongue. MAURIAC: Tertiary Syphilitic Glossopathies. FERRIER: Paralysis of 5th Cranial Nerve. HILLABY: Treatment of Tonsillitis. POTTER: Galvano-Cautery in Treatment of Enlarged Tonsils. VILLAR: Lymphadenoma of the Tonsils. TISSIER: Follicular Amygdalitis. CARPENTER: Sore Throat Affections. WEIL: Chronic Pharyngitis. HAGER: Infectious Phlegmon of Pharynx. BARUCK: Ibid. M.D.: Pharyngeal Syphilis. HOUTANG: Softening Gumma of the Pharynx. MOURE: Mycosis of the Pharynx. BURKHARDT: Retro-Pharyngeal Abscesses. MELTZER: Case of Dysphagia. TERRILLON: Stricture of the Œsophagus. AUDRY: Ibid.

Larynx, p. 242.—SEILER: Physiology of the Voice and Speech. BRACHET and COUPARD: Hygiene of the Vocal Organs. BRUGELMANN: Asthma. SEIFFERT and HOFFA: Congenital Membranous Formation in the Larynx. LENICKE: Cicatricial Laryngo-Stenoses. KOCH: Phonetic Spasm of the Glottis. REMAK: Traumatic Paralysis of the Sympathetic Nerves. SEMON: Vulnerability of the Fibres of the Recurrent Nerve. ROSENBACH: Ibid. FRANKEL: Remarks on Remak's case. OKELL: Tracheotomy in Tubercular Laryngitis. MACKENZIE: Thyrotomy for Growths of the Larynx. SMITH: Removal of Foreign Bodies from Air-Passages. L'ABBÉ: Foreign Body in the Larynx. DEMONS: Extirpation of the Larynx. DU CAZAL: Wounds of the Larynx. LUC: Tracheal Ozena.

Neck, &c., p. 246.—PETIT: Gaseous Tumours of the Neck. SEGOND: Cystic Goitre. GAUTHIER: Thyroïdean Cachexia. VIERORDT: Goitre and Laryngitis Hypoglottica. BALLET: Exophthalmic Goitre. HAW: Hæmorrhage in Myxœdema. DAVIES: Ibid. JALLAND: Tumours of Left Parotid. TILLAUX: Lipoma of the Neck. WALTER: Report of the Polyclinic in Munich.

Congress Meeting, p. 248.—Congress of Physicians in Vienna.

Reports of Societies, p. 249.—Northumberland and Durham Medical Society; Pathological Society of London; Sunderland and North Durham Medical Society; British Medical Association: Nova Scotia Branch; Royal Medical and Chirurgical Society; Medical Society of London; Royal Academy of Medicine in Ireland; Clinical Society of Manchester; Cambridge Medical Society.

Tabloid Triturates { The Latest Advance.

LONDON.]

JUNE.

[1888.

"Tabloids" of Triturated Drugs.

Safer, Pleasanter, and more Efficient and Convenient Medication for Infants, the Fastidious and Idiosyncratic.

AN INNOVATION.

Brunton points out that the introduction of the method of giving small doses at frequent intervals has "the very great advantage that the desired effect can be produced with greater certainty and with less risk of an over-dose being taken."

As fulfilling every requirement possible as regards a safer, pleasanter, more accurate and convenient mode of administering medicines to infants as well as others, and as essentially obviating the disagreeable effects of drugs, the Compressed "TABLOID TRITURATES" are now introduced.

WHAT ARE "TABLOID" TRITURATES?

The Compressed "Tabloid" Triturates are "intimate mixtures of substances with sugar of milk." In no way are they allied to the sugar of milk globules or pellets, dependent so largely upon chance for the absorption of the medicaments poured down the side of the bottle. The following directions are those given in the Pharmacopœia, U.S., for the preparation of Triturates: "Take of the substance ten parts, sugar of milk in moderately fine powder ninety parts, to make one hundred parts; weigh the substance and the sugar of milk separately; then place the substance, previously reduced if necessary to a moderately fine powder, into a mortar, add about an equal bulk of sugar of milk, mix well by means of a spatula, and triturate them thoroughly together. Add fresh portions of the sugar of milk from time to time, until the whole is added, and continue the trituration until the substance is intimately mixed with the sugar of milk and finely comminuted."

RESUMÉ OF ADVANTAGES

Of "Tabloid" * Triturates:—

1. The Tabloid Triturates are made with the pure drug and sugar of milk.
2. The process of trituration employed so finely subdivides and separates the mass of medicament that this is said to be more active than would the same quantity given in the ordinary way.
3. They contain each a very small dose, so that by giving one at a time—they may be repeated often—the taste of the drug is hardly, if at all perceived.
4. Being made with sugar of milk, one of them (if not taken whole) added to a little milk or other fluid, is at once "broken up" and distributed throughout the liquid.

* A very interesting and instructive essay on this subject by William Murrell, M.D., F.R.C.P., appeared in the PRACTITIONER for Dec., 1887.

5. Pulverulent substances, like calomel, are by this means especially distributed well, and for the moment suspended throughout the fluid.

6. They do not roll about the mouth, or stick in the throat, like pills. A great advantage.

7. Being very small, and not globular, they are easy to swallow.

8. They do not harden and become insoluble with time, nor do they crumble, like pills.

9. They are the most convenient form of medicine for infants and fastidious patients, as such a small quantity may be taken at a time that the taste becomes insignificant.

10. They afford the advantages derivable from the administration of small doses repeated often, which are,—1. That if the drug be given in but little liquid, the absorbent power of the mucous membrane of the mouth and gullet are called repeatedly into requisition. 2. That if given on an empty stomach (as is generally desirable) unpleasant symptoms are avoided. 3. In case of idiosyncrasy, the dose can be stopped before large amounts have been given. 4. Administered in this way drugs are better tolerated than is otherwise the case.

11. A greater effect is alleged to be obtainable by this method from a small quantity of medicine than is possible by the usual plan.

12. In some cases, "Tabloid" Triturates are repeated as often as every five or ten minutes, and it is surprising how soon a very small dose of medicine repeated often amounts to a very large quantity.

13. If taken whole, one of the "Tabloid" Triturates dissolves and falls to pieces in the stomach at once, and is never voided unchanged.

14. They afford accuracy of dose, without the trouble and annoyance of weighing or measuring.

15. They can be taken at any time and in any place, even when the patient is following his ordinary avocation.

16. They are only a few lines in thickness, and about quarter the circumference of a lead-pencil.

LIST OF TABLOID TRITURATES.

Price to the medical profession, 83d. per bottle of 100 of the following Tabloid Triturates. Retail price, 1s.	Price to the medical profession, 83d. per bottle of 50 of the following Tabloid Triturates. Retail price, 1s.
Aconite Tinct. 1 min.	Anti-Strych. Aloin 1-5 gr.
Arseous Acid. 1-100 and 1-50 gr.	Constipation Belladon. Ex. 1-8 gr.
Belladonna Tinct. 1 min.	Ipecac. 1-16 gr.
Calcium Sulphide 1-10 gr.	Apomorphine Mur. 1-50 gr.
Capsicum Tinct. 1 min.	Atropin Sulph. 1-100 gr.
Digitalis Tinct. 1 min.	Digitain 1-100 gr.
Hydrarg. Perchlor. 1-100 gr.	Fuonymin Resin 1-8 gr.
Hydrarg. Cum Crete. 1-3 gr.	Hydrarg. Iod. Rub. 1-20 gr.
Hydrarg. Subchlor. (Calomel) 1-10 gr.	Hydrarg. Iod. Vir. 1-8 gr.
Hyo-cyamus Tinct. 1 min.	Morphine Sulph. 1-20 and 1-8 gr.
Nux Vomica Tinct. 1 min.	Opium Tinct. (A. sudanum) 2 min.
Tinct. Camph. Co. (Paregoric). 2 min.	Plilocarpin. Mur. 1-20 gr.
	Podophyllin Resin. 1-4 gr.
	Santonin 1-2 gr.
	Strophanthus Tinct. 2 min.

Waistcoat Pocket Cases, containing eleven tubes of 25 tabloids each (any selection), supplied to the medical profession at 5s. post free.

BURROUGHS, WELLCOME, & CO., Snow Hill Buildings, LONDON, E.C.