

(89.5%). At least one repeat review was indicated in 10 of the 19 patients (52.6%).

Conclusions: Trauma patients have a high rate of comorbid mental illness. Nearly 1/5 are currently referred to the CLP service, which is likely an underestimation of the actual burden of mental health disorders and could be explained by the lack of dedicated services. The liaison psychiatry team provides valuable input into the multi-disciplinary care of trauma patients and the demand for its services is likely to increase with the expansion under the Major Trauma Strategy for Ireland.

Disclosure of Interest: None Declared

Eating Disorders 02

EPP0820

Stages of treatment of eating disorders in endogenous depressions

A. Barkhatova*, A. Smolnikova and M. Bolgov

Department of endogenous mental disorders and affective states, Mental health research center, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1105

Introduction: The problem of eating disorders has become increasingly important in recent years, due to the increase in the number of cases among children and adolescents, as well as the insufficient effectiveness of therapeutic measures. An important role in the course of eating disorders is occupied by the depressive syndrome associated with it, which complicates the process of treatment and rehabilitation in this pathology.

Objectives: Study of the stages of treatment of eating disorders in the structure of depressive states.

Methods: The sample consisted of 63 patients aged from 15 to 25 years old (all female, average age 16.2), who were on outpatient and inpatient observation in the clinic were studied.

Results: In the process of treatment, several stages of treatment of patients were carried out. The first stage was aimed at normalizing the body's vital functions and management of somatoendocrine impairments (the duration of the stage is about 14 days). The next stage was aimed at the psychotropic treatment of eating disorders and concomitant mental pathologies (the duration of the stage is 3-4 weeks). The final stage included rehabilitation, which consisted of working with a psychotherapist (the duration of the stage was 8 weeks or more). It should be noted that in the process of rehabilitation, patients continued to receive psychopharmacotherapy and underwent a comprehensive examination to assess the dynamics of their condition.

Conclusions: Eating disorders in the structure of endogenous depressions require an integrated approach to treatment, including both ensuring adequate vital activity of the organism and the selection of drug treatment depending on the nosological affiliation of the underlying syndrome. Rehabilitation work aimed at social adaptation and prevention of relapses of the disease also plays an important role.

Disclosure of Interest: None Declared

EPP0821

Anorexia Nervosa and Psychiatric Comorbidities – It's not all about food

A. S. Morais*, F. Martins, P. Casimiro, V. Henriques, N. Descalço, R. Diniz Gomes, S. Cruz and N. Costa

Psychiatry, Hospital Garcia de Orta, Lisbon, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1106

Introduction: Anorexia nervosa (AN) is a severe psychiatric disorder that usually begins during adolescence and is associated with a high risk of mortality and morbidity, its treatment is complex and often ineffective. Psychiatric comorbidity is common in patients with eating disorders (with the prevalence of 20–95%), namely 39% in AN.

Objectives: The purpose of the authors is to review the most common areas of psychiatric comorbidity in AN, how it affects the course of both diseases and the potential treatment approaches.

Methods: A brief non-systematized review is presented, using the literature available on PubMed and Google Scholar.

Results: The most common psychiatric comorbidities in AN are: Affective disorders in 24-38% (mainly unipolar depression which can appear in up to 75% of patients, compared to 11% in bipolar disorder); Anxiety disorders in 25.5% (11% with panic disorder, 20% social phobia/social anxiety disorder, 15% specific phobias, 10% generalized anxiety disorder, 13% post-traumatic stress disorder); Obsessive compulsive disorder in 12%; Substance use disorders at 17%; Personality disorders around 30%. Other pathologies occur less commonly but can have a significant impact on the patient, namely Autism spectrum disorder (predictive factor for unfavourable outcome) or Schizophrenia (there are reports of reciprocal relationships between the two pathologies).

Some of these comorbidities may increase mortality in AN, namely unipolar depression, personality disorders, alcohol and illicit drug use. The profound impact that starvation has on mood and cognition is well known. It can condition symptoms that are confused with other psychiatric diseases and change their clinical presentation. As such, the specific clinical characteristics and the therapeutic approach will be presented for each of the psychiatric comorbidities.

Conclusions: Early diagnosis and treatment of psychiatric comorbidities in AN are essential to improve the prognosis of this eating disorder. The additional treatment of these pathologies will increase complexity of the already challenging treatment of AN, with the additional symptomatology often being perpetuated by an uncontrolled eating disorder and a poor compliance to treatment. The limited evidence available for approaching these cases is based on the few studies available, most with insufficient samples.

Disclosure of Interest: None Declared

EPP0822

Intergenerational transmission of childhood maltreatment and eating disorder behaviors: Shedding light on the mother-daughter dyad and grandmother-mother-daughter triad

A. Talmon^{1*} and N. Tsur²

¹Hebrew University, Jerusalem and ²Tel Aviv University, Tel Aviv, Israel

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1107