

Anxiety, and Stress Scale-21 (DASS-21), for measuring spirituality/religiosity we used The Duke University Religion Index.

Results: The highest rate of vaccinated individuals was noted in a group of medical students (69.9%), the lowest - among responders studying science (1.9%). Students who wanted to be vaccinated had higher levels of depressive, anxiety, and stress symptoms compared to those who were already vaccinated ($p=0.04$); they also had higher depressive symptoms than unvaccinated and unwilling participants ($p=0.028$). Students who didn't want to be vaccinated against COVID-19 showed the highest religiosity compared to those who would like to be vaccinated ($p<0.001$) or were vaccinated ($p=0.003$). There was a negative correlation between the level of religiosity and severity of depressive and anxiety symptoms ($p=0.002$).

Conclusions: 1. The attitudes towards vaccination against COVID-19 depended on the fields of study. 2. Religiousness has been linked with the attitudes towards COVID-19 vaccination as well as level of depression and anxiety amongst Polish students.

Disclosure: No significant relationships.

Keywords: mental health; religiosity; Covid-19; vaccination

EPV0399

Changes in Inpatient and Day Patient Treatment during the COVID-19 Pandemic: Insights and Outlook Results of a Study in an Adult Psychiatry Clinic in Germany

T. Barth^{1*} and B. Voigtländer²

¹Klinikum Chemnitz, Psychiatry, Chemnitz, Germany and ²Klinikum Chemnitz, Psychiatry, Dresden, Germany

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1263

Introduction: Possible effects of pandemic-related restrictions and adjustments in psychiatric treatment are currently the focus of interest. This article addresses developments in this respect in our clinic.

Objectives: Changes in clinical practice that occurred during the COVID-19 pandemic are to be analysed with regard to possible risks for affected patients.

Methods: A clinic-internal analysis was carried out, focusing on comparing a period during the COVID-19 pandemic (01.11.2020-30.04.2021) and a pre-pandemic reference period (01.11.2018-30.04.2019).

Results: Following trends were observed during the pandemic period: a. Day patient treatment: - The treatment volume fell to 44%. Notable reductions in the number of treatment cases with main diagnoses [ICD-10] F10.- and F30-F39 by >65% were measured. b. Inpatient treatment: - no significant changes regarding socio-demographic patient data and concerning the type of admission and discharge, - detection of coronavirus SARS-CoV-2 by PCR test in 4.7% of the cases, - a decline in the treatment volume to 87% due to 8% decrease in the number of cases and 5% decrease in the ALOS, with patients with the main diagnosis [ICD-10] F10.- were most affected, - increases regarding the percentage of cases with the main diagnosis [ICD-10] F40-F48 ($p<0.05$) and in the ALOS of this

patient group (by 31%), - a significant rise ($p<0.01$) in the ratio of restrained treatment cases.

Conclusions: Related to the decrease in treatment volume, the increase in psychiatric intensive treatment and possible risks in terms of the care situation for patients with the main diagnoses [ICD-10] F10.-, F30-F39 and F40-F48 should be discussed in particular.

Disclosure: No significant relationships.

Keywords: covid; Changes in Pandemic Situation; Clinical Treatment

EPV0400

Bidirectional Associations Between Covid-19 Infection And Mental Disorders

M. Turki, S. Blajji*, A. Daoud, S. Ellouze, M. Abdellatif, N. Gargouri, N. Halouani and J. Aloulou

Hedi Chaker University Hospital, Psychiatry "b" Department, Sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1264

Introduction: The COVID-19 pandemic brought unbearable psychological pressure to people worldwide, because of serious threats to one's physical health and life. From early stages of this pandemic, concerns have been raised about its effect on mental health. However, we still know little whether pre-existing psychiatric disorder (PD) affects the susceptibility and evolution of this infection.

Objectives: We aimed to assess the interactions between COVID-19 infection and PD.

Methods: We conducted a literature review through pubmed database, using the keywords :«COVID 19», «psychiatry», «mental disorders», «schizophrenia», «anxiety», «depression», «insomnia».

Results: On one hand, prior surveys suggested that the infection is associated with increased incidence of a first psychiatric symptom. Mental health disturbances mostly include anxiety, depression, sleep disturbances, cognitive impairment and post-traumatic stress disorder. On the other hand, recent studies showed that patients with pre-existing mental disorders were associated with high susceptibility to be infected, increased risk of intensive care unit admission and a high mortality. The susceptibility to contracting COVID-19 was associated with pre-existing mood disorders, anxiety, and attention-deficit hyperactivity disorder. Infection severity was associated with pre-existing or subsequent mood disorders and sleep disturbances; or a pre-existing schizophrenia. Mortality is increased in patients diagnosed with schizophrenia.

Conclusions: The complicated interactions between COVID-19 infection and PD have several implications. Enhanced psychiatric follow-up should be considered for survivors of COVID-19. Besides, early detection and intervention for PD are needed to control morbidity and mortality induced by the COVID-19 infection.

Disclosure: No significant relationships.

Keywords: COVID19; psychiatry; Mental Disorders