

Parasitology

Back volumes. Vols. 1–71 : Inquiries should be addressed to Wm. Dawson & Sons Ltd, Cannon House, Folkestone, Kent. Vols. 72 onwards : quotations for parts still in print may be obtained from Cambridge or the American Branch of Cambridge University Press.

Copying. This journal is registered with the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923. Organizations in the USA who are also registered with C.C.C. may therefore copy material (beyond the limits permitted by sections 107 and 108 of US copyright law) subject to payment to C.C.C. of the per-copy fee of \$11.00. This consent does not extend to multiple copying for promotional or commercial purposes. Code 0031–1820/96 \$11.00 + .10.

Organizations authorized by the Copyright Licensing Agency may also copy material subject to the usual conditions.

ISI Tear Sheet Service. 3051 Market Street, Philadelphia, Pennsylvania 19104, USA, is authorized to supply single copies of separate articles for private use only.

For all other use, permission should be sought from Cambridge or the American Branch of Cambridge University Press.

Claims for missing issues can only be considered if made immediately after receipt of the subsequent issue.

Advertising. Details of advertising in *Parasitology* may be obtained from the publisher.

© Cambridge University Press 1996

The Pitt Building, Trumpington Street, Cambridge CB2 1RP
40 West 20th Street, New York, NY 10011–4211, USA
10 Stamford Road, Oakleigh, Melbourne 3166, Australia

Printed in Great Britain by the University Press, Cambridge

Parasitology

CONTENTS

	PAGE
Crandall, I. and Sherman, I. W. The human anion transport protein, band 3, contains a CD36-like binding domain for <i>Plasmodium falciparum</i> -infected erythrocytes	261
Clarebout, G., Gamain, B., Slomianny, C., Camus, D. and Dive, D. The course of <i>Plasmodium berghei</i> , <i>P. chabaudi</i> and <i>P. yoelii</i> infections in β -thalassaemic mice	269
Quijada, L., Requena, J. M., Soto, M. and Alonso, C. During canine viscerocutaneous leishmaniasis the anti-Hsp70 antibodies are specifically elicited by the parasite protein	277
Sullivan, A. D., Ittarat, I. and Meshnick, S. R. Patterns of haemozoin accumulation in tissue	285
Dennis, R. D., Baumeister, S., Lauer, G., Richter, R. and Geyer, E. Neutral glycolipids of <i>Schistosoma mansoni</i> as feasible antigens in the detection of schistosomiasis	295
Upcroft, J. A., Campbell, R. W. and Upcroft, P. Quinacrine-resistant <i>Giardia duodenalis</i>	309
Taylor, M. J., Cross, H. F., Mohammed, A. A., Trees, A. J. and Bianco, A. E. Susceptibility of <i>Brugia malayi</i> and <i>Onchocerca lienalis</i> microfilariae to nitric oxide and hydrogen peroxide in cell-free culture and from IFN γ -activated macrophages	315
Folkard, S. G., Hogarth, P. J., Taylor, M. J. and Bianco, A. E. Eosinophils are the major effector cells of immunity to microfilariae in a mouse model of onchocerciasis	323
Hong, X. Q., Santiago Mejia, J., Kumar, S., Perler, F. B. and Carlow, C. K. S. Cloning and expression of DiT33 from <i>Dirofilaria immitis</i> : a specific and early marker of heartworm infection	331
Uchikawa, R., Yamada, M., Matsuda, S., Tegoshi, T., Nishida, M., Kamata, I., Kuroda, A. and Arizono, N. Dissociation of early and late protective immunity to the nematode <i>Nippostrongylus brasiliensis</i> in Brown Norway and Fischer-344 rats	339
Kennedy, C. R. Establishment, survival and site selection of the cestode <i>Eubothrium crassum</i> in brown trout, <i>Salmo trutta</i>	347

CAMBRIDGE
UNIVERSITY PRESS



0031-1820(199603)112:3;1-W