

save time and money for psychiatric health systems. Lastly, concerning Virtual Reality (VR) in psychiatric care, 57% of participants were familiar with VR technology, but only 43% were acquainted with its applications in psychiatry. Notably, 71% did not view VR as a viable replacement for in-person psychiatric management.

Conclusions: This is the first study conducted on understanding digital psychiatry in Pakistan's healthcare system, which revealed multiple challenges to digital health competency among psychiatrists. This emphasizes on the need for formal training and funding towards resources to overcome obstacles in utilizing mental health technologies.

Disclosure of Interest: None Declared

EPV0434

Patterns of Internet Addiction in an Italian sample: 100% of the sample experience Nomophobia

F. Marcolini*, D. De Ronchi and A. R. Atti

Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1139

Introduction: Internet Addiction Disorder, a concept introduced for the first time by Ivan Goldberg in 1995, is one of the most recently identified forms of addiction, but already considered a real psychosocial phenomenon, capable of having a profound impact on different aspects of social and psychological life of individuals. One of its most recently identified manifestations is Nomophobia, a neologism formed from the combination of terms such as “no mobile”, “phone” and “phobia”, which can be understood as the fear of feeling disconnected. It is today considered a situational phobia, characteristic of contemporary times. The most common symptoms include excessive cell phone use and constant anxiety at the thought of losing the internet connection. Others are, for example, “Ringxiety”, ringing anxiety, or the “phantom vibration syndrome”.

Objectives: This study aims to examine the spread of Nomophobia in the Italian population, evaluating psychopathological correlations that can explain its diffusion.

Methods: Between January and May 2023, an anonymous online questionnaire was randomly sent to the general population. Alongside with tests to evaluate psycho-social features, the instrument used to study Nomophobia was the *Nomophobia Questionnaire* (NMP-Q) (Yildirim *et al.* *Comput Hum Behav.* 2015; 49:130–7), in its Italian version (Adawi *et al.* *JMIR MHealth UHealth.* 2018;6:e24).

Results: The sample consists of 308 people (189 F, 119 M), with an average age of 32 years (*sd* 14). In our sample, 100% of the subjects tested positive for Nomophobia. Values indicating a state of severe Nomophobia are found in 12.3% of the sample (F 15.9%, M 6.7%). The young population, between 18 and 25 years old, represents 54% of the affected population, but more than 60% of severe cases (95% confidence interval 50-65%). The severe cases correlate positively ($p < 0,05$) with findings of high impulsiveness. There are no other studies that investigate the psychopathological correlates of Nomophobia among Italians.

Conclusions: Despite possible *biases*, the data obtained are an alarming sign of the spread of internet addiction that characterizes our times, of which the excessive use of cell phones in the form of

Nomophobia is an expression. Despite their now undisputed usefulness, mobile devices are capable of causing the onset of serious health problems, starting from exposure to radiation capable of causing dermatitis, tumors, and infertility. Furthermore, they dramatically interfere with driving safety, becoming a major cause of road accidents. Considering these consequences, it appears to be extremely important to characterize the phenomenon, as well as its psychosocial determinants, in order to proceed with its better definition and prevention.

Disclosure of Interest: None Declared

EPV0435

Assessing empathy among caregivers: a cross-sectional study

M. Kahloul¹, I. Kacem^{2*}, A. Ghenim², A. Aloui², A. Chouchane², M. Ajmi¹, W. Naija¹, M. Maoua² and N. Mrizak²

¹Sahloul Academic Hospital, Anesthesia and Intensive Care Department and ²Occupational Medicine Department, Farhat Hached Academic hospital, Sousse, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1140

Introduction: Empathy plays an important role in everyday human relationships. It is the ability to put oneself in the place of others, to represent what they think and feel. In healthcare settings, several studies have highlighted its positive effects on patients in terms of physical and psychological well-being.

Objectives: Evaluate empathy among caregivers.

Methods: This is a cross-sectional study, conducted over a 1-month period and enrolling nursing staff working at Farhat Hached Academic hospital. Empathy was assessed using the Jefferson Scale of physician's empathy (JSPE) scale.

Results: A total of 92 caregivers were enrolled in this study. The average age was 40.41 years with a sex ratio of 0.95. The most represented category was nurses (64.1%) with an average seniority of 17.2 years. The average empathy score was 98.4. Scores above half were reported in 69.5% of cases. The presence of empathy was significantly associated with female gender ($p = 0.002$).

Conclusions: Empathy is a key point in the patient-caregiver relationship. Thus, the nursing staff must be aware of this concept in order to improve the quality of care.

Disclosure of Interest: None Declared

EPV0436

The opportunity for e-mental health to overcome stigma and discrimination

K. Subramaniam^{1*}, A. Greenshaw² and A. Thapliyal³

¹Global Medical Affairs, Viatrix, Auckland, New Zealand; ²Dept of Psychiatry, University of Alberta, Edmonton, Canada and ³e-Mental Health International Collaborative, Auckland, New Zealand

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1141