

Marine Record

Cite this article: De Weerd J (2023). A new record of a white humpback whale (*Megaptera novaeangliae*) in Papeete, Tahiti. *Journal of the Marine Biological Association of the United Kingdom* **103**, e78, 1–3. <https://doi.org/10.1017/S002531542300067X>

Received: 4 April 2023

Revised: 24 August 2023

Accepted: 1 September 2023

Keywords:

albinism; French Polynesia; leucism; marine mammal; Oceania; pigmentation; stock F

Corresponding author:

Joëlle De Weerd

Email: eliscientific@gmail.com

© The Author(s), 2023. Published by Cambridge University Press on behalf of Marine Biological Association of the United Kingdom. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



A new record of a white humpback whale (*Megaptera novaeangliae*) in Papeete, Tahiti

Joëlle De Weerd^{1,2} 

¹Association ELI-S, Education, Liberté, Indépendance – Scientifique, Allée de Verdalle 39, 33470 Gujan-Mestras, France and ²Biology Department, Vrije Universiteit Brussel, VUB, Pleinlaan, 1050 Brussel, Belgium

Abstract

A new record of a white humpback whale (*Megaptera novaeangliae*) was made off Papeete, Tahiti in July 2019. The individual was completely white, except for grey patterns on its body and on the ventral part of the fluke. The presence of different colourations indicates that this individual is not an albino but a leucistic individual. According to field observations, the individual was likely a juvenile or a subadult. It was seen breaching several times and travelled further north of Papeete, where the animal was recaptured off Arue. White individuals are rare and offer unique opportunities to understand whale movement patterns and potential population connectivity among humpback whale populations.

Melanin production in mammals is regulated by the enzyme tyrosinase (Behrmann, 1998). Melanocytes and chromatophore cells are responsible for the production of melanin, which results in the dark skin colour of cetaceans (Behrmann, 1998). Albinism is characterized by the complete absence of melanin, and consequently, a lack of pigmentation in the skin, hair, and eyes (Grønsvov *et al.*, 2007). Leucism occurs when a lack of melanin partially affects skin pigmentation in some areas of the body, excluding the eyes (Fertl and Rosel, 2009). Piebaldism occurs when the lack of pigmentation in the skin is random, leading to the display of patchy patterns (Fertl and Rosel, 2009). Anomalous white colouration in marine mammals can negatively affect survival due to the increased visibility of predators (Hain and Leatherwood, 1982), and decreased protection from ultraviolet radiation and sun damage (Martinez-Levasseur *et al.*, 2011). Anomalous white colouration in mammals can cause health problems, such as anaemia, reduced fertility, sensory system defects, and increased susceptibility to infections (Searle, 1968); however, it is currently unclear whether these health problems affect marine mammals. Albinism and anomalous white colouration were recorded in approximately 25 cetacean species (Fertl and Rosel, 2009) including 17 odontocetes (*e.g.* killer whales [Speckman and Sheffield, 2001], pilot whales [Hain & Leatherwood, 1982], harbour porpoises [Tonay *et al.*, 2012; Kopalani *et al.*, 2017], short-beaked dolphins [Kopalani *et al.*, 2017]), and eight mysticetes (*e.g.* grey whales [Willoughby *et al.*, 2018], and humpback whales [Polanowski *et al.*, 2012; Burns *et al.*, 2014; Pirota *et al.*, 2023]).

The humpback whale, *Megaptera novaeangliae*, is a migratory species morphologically characterized by dark grey to black dorsal surface colour with dark or white ventral sides according to the population from which they originate. Northern Hemisphere humpback whales tend to be totally black, while the Southern Hemisphere humpback whales are mostly white in the ventral area (Chittleborough, 1965; Allen *et al.*, 1994; Acevedo *et al.*, 2017). The variation in skin pigmentation is used for individual identification, which allows to assess migratory routes, movement patterns, and site fidelity (Forestell *et al.*, 2001; Lydersen *et al.*, 2013; Pirota *et al.*, 2023). Southern Hemisphere humpback whale populations are divided into seven breeding stocks (A–G) according to the International Whaling Commission. Anomalous skin colour patterns (leucism) were previously reported in humpback whales in Ecuador (stock G, 2002, Castro *et al.*, 2007), in eastern Australia (stock E, 1991, Forestell *et al.*, 2001), in Norway (Cape Verde, 2012, Lydersen *et al.*, 2013), and in South Africa (stock B, 2014, Koper *et al.*, 2017). Albinism was only confirmed genetically in the eastern Australia sighting (Forestell *et al.*, 2001; Polanowski *et al.*, 2012). Two presumed cases of albinos were reported both in eastern Australia in 2011 (Koper *et al.*, 2017) and in Costa Rica in 2022 (Mora *et al.*, 2022).

In this note, I report the first record of a humpback whale with anomalous skin pigmentation sighted opportunistically off the coast of Papeete, Tahiti, French Polynesia. Two opportunistic sightings were made on the same day (Figure 1A, B). On 25 July 2019, at 0930 am a white humpback whale was observed off Papeete (17°31′30.0″S, 149°34′34.9″W) during a private boat trip (Figure 1). Data were collected by a citizen (Montlahuc) during this trip and included photographs of the whale, geographic location, behaviour, and time of the sighting. Photographs were taken with a Sony DSC-Rx10M4 camera. The whale was re-sighted by divers about 5 km away in the early afternoon of the same day farther north near the commune of Arue (Montlahuc, pers. comm.) (Figure 1C). No coordinates were obtained from this resighting. The whale has not been resighted since 2019 by local nonprofits dedicated to whale research (Bennet, pers. comm.). It was impossible to determine the individual's size based on *in situ* observations. The individual was displaying surface-active behaviours such




Figure 1. Location of French Polynesia (A, B) and the observation location in Papeete, Tahiti (C) with a black arrow indicating the track of the boat that followed the whale.

as breaching and tail slapped several times. The individual presented a mixed grey-white body colour with some skin lesions likely made by cookie shark (*Isistius brasiliensis*) bites (Figure 2A–D). The ventral surface of the flukes displayed

some white-grey-coloured patterns (Figure 2D). Pink skin was also observed on the dorsal fin of the animal (Figure 2A). No photograph could be taken of the eyes to assess the presence of red eyes to characterize albinism. This report represents the




Figure 2. Photographs of an anomalous skin-coloured humpback whale observed in Papeete, Tahiti, on 25 July 2019 showing the dorsal surface of (A) the dorsal fin and (B) the dorsal side of the fluke; (C) the right side of the head and right pectoral flipper head and (D) the ventral side of the fluke with skin lesions likely made by cookie shark (*I. brasiliensis*) bites. Photo credit: Olivier Montlahuc.

first documented case of a leucistic white whale in the breeding stock F.

Humpback whales observed in Oceania are part of breeding stock F, which ranges from the Cook Islands (substock F1) to French Polynesia (substock F2) (Pastene *et al.*, 2011). No other sighting of anomalous skin-coloured whales was previously reported in breeding stock F. Despite the disadvantages that white skin colours may present, two adult white whales are known to be at least 20 years old as reported off the Norwegian mainland coast (Lydersen *et al.*, 2013) and eastern Australia (Pirota *et al.*, 2023). Except for those two whales, few photographic or visual recaptures of white whales were reported in the past. The atypical body colouration of the individual sighted off Tahiti can result in a higher likelihood to be resighted compared to normal skin colouration, especially if it occurs in coastal areas with high anthropogenic activity. No recaptures of this animal were made on the Happywhale citizen science platform containing a database of approximately 289,000 encounters and uses fully automated fluke-matching software (Cheeseman *et al.*, 2022).

Anomalous white colouration can be of genetic origin due to mutations in the tyrosinase enzyme, and albinism is the result of homozygous deletion of the nucleobase cytosine in DNA (Polanowski *et al.*, 2012). In the Southern Hemisphere, the recessive allele responsible for depigmentation in humpback whales has been reported in stock E (eastern Australia; Forestell *et al.*, 2001) and potentially in stock G (Central to South America; Mora *et al.*, 2022). It is unclear if this phenomenon is the result of a recessive allele transmission between stocks or if this results from an independent genetic anomaly (e.g. insertions or deletions). Interchange has been found in the past between stocks E and F (Western Australia) through photo-identification methods (Garrigue *et al.*, 2011). If the recessive allele responsible for albinism is present in stock F it might indicate potential inter-stock movements with stock E, where one albino was previously observed (Pirota *et al.*, 2023). Future resighting of anomalous white-coloured humpback whales should include biopsy sampling and genetic analysis of skin samples to understand the origin of the genetic deviation (e.g. gene mutation or allele transfer) to assess potential stock exchanges and reproduction.

Acknowledgements. I would like to thank Mathilde Giry for indicating the existence of this albino sighting and Olivier Montlahuc for sharing his observation, photographs, and related information. Special thanks to Marie Van Bressems for her input on the skin lesion analysis, PC and the two reviewers for revising this manuscript, and Agnes Bennet from Association Mata Tohora confirming the absence of recapture of this white humpback whale in the area. No permits were required for the data collection.

Financial support. This research received no specific grant from any funding agency, commercial, or not-for-profit sectors.

Competing interest. None.

Data availability. All the data are available upon request.

References

- Acevedo J, Aguayo-Lobo A, Allen J, Botero-Acosta N, Capella J, Castro C, Rosa LD, Denkinger J, Félix F, Flórez-González L, Garita F, Guzmán HM, Haase B, Kaufman G, Llano M, Olavarría C, Pacheco AS, Plana J, Rasmussen K, Scheidat M, Secchi ER, Silva S and Stevick PT (2017) Migratory preferences of humpback whales between feeding and breeding grounds in the eastern South Pacific. *Marine Mammal Science* 33, 1035–1052.
- Allen JM, Rosenbaum HC, Katona SK, Clapham PJ and Mattila DK (1994) Regional and sexual differences in fluke pigmentation of humpback whales (*Megaptera novaeangliae*) from the North Atlantic Ocean. *Canadian Journal of Zoology* 72, 274–279.
- Behrmann VG (1998) The origin of the skin colour of toothed whales (Odontoceti). *Übersee-Museum Jahrbuch* 7, 131–142.
- Burns D, Brooks L, Harrison P, Franklin T, Franklin W, Paton D and Clapham P (2014) Migratory movements of individual humpback whales photographed off the eastern coast of Australia. *Marine Mammal Science* 30, 562–578.
- Castro C, Groch K, Marcondes M, Van Bressems M and Van Waerebeek K (2007) Miscellaneous skin lesions of unknown aetiology in humpback whales *Megaptera novaeangliae* from South America. *SC/60/DW18*, 6 pp.
- Cheeseman T, Southerland K, Park J, Olio M, Flynn K, Calambokidis J, Jones L, Garrigue C, Frisch JA, Howard A, Reade W, Neilson J, Gabriele C and Clapham P (2022) Advanced image recognition: a fully automated, high-accuracy photo-identification matching system for humpback whales. *Mammalian Biology* 102, 915–929.
- Chittleborough R (1965) Dynamics of two populations of the humpback whale, *Megaptera novaeangliae* (Borowski). *Marine and Freshwater Research* 16, 33–128.
- Fertl D and Rosel PE (2009) Albinism. In William FP, Würsig B and Thewissen JGM (eds), *Encyclopedia of Marine Mammals*. Elsevier, pp. 24–26.
- Forestell PH, Paton DA, Hodda P and Kaufman GD (2001) Observations of a hypo-pigmented humpback whale, *Megaptera novaeangliae*, off east coast Australia: 1991–2000. *Memoirs of the Queensland Museum* 47, 437–450.
- Garrigue C, Franklin T, Constantine R, Russell K, Burns D, Poole M, Paton D, Hauser NAN, Oremus M, Childerhouse S, Mattila D, Gibbs N, Franklin W, Robbins J, Clapham P and Baker CS (2011) First assessment of interchange of humpback whales between Oceania and the east coast of Australia. *Journal of Cetacean Research and Management*, Special Issue 3, 269–274.
- Gronskov K, Ek J and Brondum-Nielsen K (2007) Oculocutaneous albinism. *Orphanet Journal of Rare Diseases* 2, 1–8.
- Hain JHW and Leatherwood S (1982) Two sightings of white pilot whales, *Globicephala melaena*, and summarized records of anomalously white cetaceans. *Journal of Mammalogy* 63, 338–343.
- Kopaliani N, Gurielidze Z and Ninua L (2017) Records of anomalously white harbour porpoises and atypical pigmented short-beaked common dolphin in the Georgian Black Sea waters. *Journal of the Black Sea/Mediterranean Environment* 23, 66–74.
- Koper RP, Drost E and Plön S (2017) First sighting of a leucistic humpback whale (*Megaptera novaeangliae*) in South African coastal waters. *Aquatic Mammals* 43, 331–334.
- Lydersen C, Oien N, Mikkelsen B, Bober S, Fisher D and Kovacs KM (2013) A white humpback whale (*Megaptera novaeangliae*) in the Atlantic Ocean, Svalbard, Norway, August 2012. *Polar Research* 32, 1–6.
- Martinez-Levasseur LM, Gendron D, Knell RJ, O'Toole EA, Singh M and Acevedo-Whitehouse K (2011) Acute sun damage and photoprotective responses in whales. *Proceedings of the Royal Society B: Biological Sciences* 278, 1581–1586.
- Mora J, Lopez L, Cruz L and Chaves G (2022) First record of a white humpback whale, *Megaptera novaeangliae*, from the Eastern Tropical Pacific. *Caribbean Journal of Science* 52, 389–396.
- Pastene LA, Goto M, Kanda N, Kitakado T and Palsboll PJ (2011) Preliminary mitochondrial DNA analysis of low and high latitude humpback whales of Stocks D, E and F. Paper SC/63/SH9 presented to IWC Scientific Committee, June 2011, Tromsø, Norway (unpublished). 16pp.
- Pirota V, Franklin W, Mansfield L, Lowe J and Peterson O (2023) Sighting records of 'Migaloo' the white humpback whale provide evidence of Australian site fidelity and use of New Zealand waters as a migratory route. *Australian Zoologist* 42, 1014–1028.
- Polanowski AM, Robinson-Laverick SM, Paton D and Jarman SN (2012) Variation in the tyrosinase gene associated with a white humpback whale (*Megaptera novaeangliae*). *Journal of Heredity* 103, 130–133.
- Searle A (1968) Pigmentation and inheritance: comparative genetics of coat colour in mammals. London; Academic Press, New York, 308pp.
- Speckman SG and Sheffield G (2001) First record of an anomalously white killer whale, *Orcinus orca*, near St. Lawrence Island, northern Bering Sea, Alaska. *Canadian Field-Naturalist* 115, 501–502.
- Tonay AM, Bilgin S, Dede A, Akkaya A, Yeşilçiçek T, Köse Ö and Ceylan Y (2012) First records of anomalously white harbour porpoises (*Phocoena phocoena*) in the Turkish seas with a global review. *Hystrix* 23, 76–87.
- Willoughby AL, Ferguson MC, Clarke JT and Brower AA (2018) First photographic match of an anomalously white gray whale (*Eschrichtius robustus*) in the Northeastern Chukchi Sea, Alaska, and Baja California, Mexico. *Aquatic Mammals* 44, 7–12.