

Conclusions: Infertility can interfere negatively in women sexuality. The investigation of sexual difficulties in infertility consultations must be systematic.

Keywords: sexuality; infertility; woman; Dysfunction

EPP1282

Improving contraceptive and family planning awareness on a perinatal inpatient unit

G. Xia

General Psychiatry, Central and North West London NHS Trust, Wembley, London, United Kingdom
doi: 10.1192/j.eurpsy.2021.1468

Introduction: Unplanned pregnancies are a significant risk factor in perinatal mental health. They also have the potential to result in adverse health impacts for mother, baby and children into later in life. Women from disadvantaged backgrounds are less likely to access contraception. Women are more likely to on board health advice during pregnancy and post partum period due to high level of surveillance by health professionals.

Objectives: Our aim was for 90% of patients on Coombe Wood Mother and Baby Unit (MBU) to feel supported to make an informed decision about their contraception by October 2020.

Methods: A questionnaire was completed by fifteen inpatients at the Mother and Baby Unit over a 4 month period (April- August 2020) to assess areas around their pregnancy and contraceptives of choice. Contraceptive training was provided by a Sexual Health Specialist to staff across multiple disciplines on Coombe Wood MBU. Sexual Health discussion groups were delivered by doctors to inpatients on a monthly basis. A post-intervention questionnaire was given to patients.

Results: •53% of patients reported unplanned pregnancies. •40% of women felt lacking confidence in choosing the right contraceptive •The most frequent question asked during the sexual health groups was regarding hormonal contraceptives impacting on mental health. •By September 100% of patients felt they were able to make an informed decision about their contraception on discharge.

Conclusions: Facilitating women to make informed decisions regarding their contraception empowers them to gain autonomy, reduces the risks of physical and mental illness, improves the quality of life for mothers and babies.

Keywords: sexual health; Contraception; Perinatal Psychiatry; Perinatal Mental Health

EPP1283

The importance of mental and sexual health in addressing people with hiv - the double stigma

S. Nascimento* and M. Gonçalves

Psychiatry, Centro hospitalar psiquiátrico de lisboa, Lisboa, Portugal
*Corresponding author.
doi: 10.1192/j.eurpsy.2021.1469

Introduction: Having a mental disorder is associated with increased vulnerability to the transmission of the Human Immunodeficiency Virus (HIV) and the prevalence of HIV is higher in people with a severe mental disorder. People with psychiatric comorbidities such as bipolar affective disorder and depressive disorder, post-traumatic stress disorder (physical or sexual abuse)

and/or psychoactive substance use have a higher risk of HIV infection.

Objectives: This work is intended to expose the importance of integrating mental health care with the care of HIV patients.

Methods: The authors conducted a non-systematic review of the literature, conducting research through Pubmed and Medscape using the keywords 'Preexposure prophylaxis', 'HIV', 'Mental health problems'.

Results: Several factors may contribute to the high comorbidity between HIV and Mental Disorders, including socio-demographic factors, weak social and environmental structures, as well as internalized stigma, social and experienced discrimination. Mental health problems may interfere with the care needed for prevention, including regular HIV testing and/or adherence to Preexposure Prophylaxis (PrEP); and influence access to and adherence to antiretroviral treatment.

Conclusions: This compelling evidence makes the necessary contribution of integrating mental health into an assessment and continuous treatment of the HIV patient, on the other hand, the assessment and treatment of mental disorders should address sexual health.

Keywords: Preexposure prophylaxis; HIV; Mental health problems; sexual health

EPP1284

Sexual function assessment in men with PTSD

R. Maalej¹, G. Hamdi^{1,2}, D. Felfel^{1*}, H. Ben Ammar¹, A. Maamri^{1,2}, Y. El Kissi² and H. Zalila¹

¹Psychiatry Department, Razi, manouba, Tunisia and ²Psychiatry, Tunisian Society of Clinical Sexology, TUNIS, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1470

Introduction: Exposure to extreme traumatic events can lead to post-traumatic stress disorder (PTSD). This disorder affects emotional, social and professional functioning. Recent studies suggest that it can lead to sexual dysfunction.

Objectives: The aim of this study is to compare the level of sexual dysfunction between men with PTSD and control subjects.

Methods: A total of 30 male PTSD patients and 30 controls were included in this study. We used the Post-Traumatic Stress Disorder Check Scale (PCLS) to assess the intensity of PTSD symptoms and the International Erectile Function Index (IIEF15) to assess sexual dysfunction of both patients and controls.

Results: The mean IIEF-15 score was 51.16 ± 6.82 in patients followed for PTSD versus 77.33 ± 2.02 in healthy controls with a non-significant difference ($p = 0.26$). Three patients (10%) had an alteration of desire while the control reported only dysfunction but there was no significant difference between the mean scores of IIEF-SD ($p = 0.22$). No patient or control had erectile dysfunction and there was no significant difference between the IIEF-EF sub-scores in the 2 groups ($p = 0.20$). The mean sexual intercourse satisfaction (SD) score in the patients was 5.13 ± 1.10 versus 8.86 ± 0.40 with a non-significant difference ($p = 0.09$). Altered satisfaction with intercourse was noted in 15% ($n = 5$) of subjects with PTSD versus a single control.

Conclusions: It is important that practitioners address the subject of sexuality in patients followed for PTSD and refer their patients, if necessary, to a sexology consultation.

Keywords: SEXUAL FUNCTION; ptsd

EPP1285

Predictors of male sexual dysfunction in post traumatic stress disorder

R. Maalej¹, G. Hamdi^{1,2}, D. Felfel^{1*}, H. Ben Ammar¹, A. Maamri^{1,2}, Y. El Kissi² and H. Zalila¹

¹Psychiatry Department, Razi, manouba, Tunisia and ²Psychiatry, Tunisian Society of Clinical Sexology, TUNIS, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1471

Introduction: Post-traumatic stress disorder affects emotional, social and professional functioning. It can also affect physical and sexual health.

Objectives: The aim of this study is to write down the prevalence of sexual dysfunction in men with PTSD and to economize on potential predictors of sexual dysfunction.

Methods: A total of 30 male patients with PTSD were included in this study. We collected socio-demographic and clinical data and we used Post-Traumatic Stress Disorder (PCL) and International Erectile Function Index (IIEF15) scales.

Results: The mean PTSD severity score was 65.43 ± 2.95 . The mean score for revitalization, avoidance, cognitive and mood alteration, and hypervigilance were 15.80 ± 1.44 , respectively; 8 ± 0 ; 24.07 ± 1.20 and 17.57 ± 2.95 . The mean IIEF-15 score was 51.16 ± 6.82 . The mean sub-scores were 3.93 ± 0.52 for sexual desire; 18.80 ± 5.68 for erectile function; 8.93 ± 8.97 for orgasmic function; 5.13 ± 1.10 for satisfaction with intercourses and 4.13 ± 1.16 for overall satisfaction. The IIEF15-EF score was negatively correlated with the presence of a personal medical history ($p = 0.02$) and the impairment cognitions and mood score ($p = 0.023$). The IIEF-OF score was significantly associated with reviviscence, hypervigilance, cognition and mood alterations ($p = 0.015$; 0.041 ; 0.045). The IIEF-15 SD score was negatively correlated with altered cognition and mood ($p = 0.007$).

Conclusions: Our study focused on the importance of assessing sexual function in men followed for PTSD and helps to understand the association of PTSD with different types of sexual dysfunction.

Keywords: ptsd; Sexual Dysfunction

EPP1286

Sexual counselling and sexual therapy in chemsex users in an NGO in Spain

J. Curto Ramos^{1,2*}, I. Azqueta¹, M.T. Heredia Soriano¹, J.F. Cabrera Solano¹ and L. Ibaruchi¹

¹Chemsex Program “sexo, Drogas Y Tú”, Apoyo Positivo, madrid, Spain and ²Mental Health Service, University Hospital La Paz, MADRID, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1472

Introduction: The intentional use of drugs before or during sexual intercourse (chemsex) is a phenomenon of special importance in the MSM (men who have sex with men) population due to its impact on mental, physical and sexual health. Sexual health issues related to chemsex practice have been described such as difficulties in achieving sober sex, erectile dysfunction or problems with sexual desire.

Objectives: To describe the sexual health interventions (including sexual counselling and sexual therapy) for patients with chemsex practices in the NGO Apoyo Positivo in Madrid. We describe the main sexual problems.

Methods: Descriptive analysis.

Results: The main sexual problems were dissatisfaction in sexual intercourse without substance and difficulties with sexual desire activation (70%); compulsive sexual behaviour (70%), difficulties with sexual orientation and non normative gender expression, difficulties in erection (34%), premature ejaculation (7%) and delayed ejaculation (10%).

Conclusions: Chemsex is a phenomenon that needs a multidisciplinary approach and mental and sexual health must be taken into account. “Sexo, Drogas y Tu” is a model of collaborative approach which is a pioneering intervention developed by an NGO in Spain.

Keywords: chemsex; NPS; sexuality

EPP1287

Chronic inflammatory bowel diseases and sexuality: Inevitable disorders?

N. Charfi¹, A. Guermazi^{1*}, F. Guermazi¹, S. Omri¹, N. Smaoui¹, R. Feki¹, J. Ben Thabet¹, L. Zouari¹, M. Maalej Bouali¹, N. Tahri², M. Boudabous² and M. Maalej¹

¹Psychiatry C Department, Hedi chaker University hospital, sfax, Tunisia and ²Gastroenterology Department, Hedi chaker University hospital, sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1473

Introduction: Improving the quality of sexual life of patients has become a major therapeutic objective in the management of Chronic Inflammatory Bowel Diseases (CIBD).

Objectives: To assess the prevalence of sexual dysfunction (SD) in patients with CIBD in remission and compare it to healthy controls (HC), and to determine the associated factors

Methods: This was a cross-sectional study, conducted over 8 months, involving 36 patients with CIBD, who attended the gastroenterology outpatient of Hedi Chaker University Hospital in Sfax (Tunisia). They were compared to 36 HC. Sexual function was assessed with the “Female sexual Function Index” and the “International Index of Erectile Function”.

Results: In the sample of CIBD, the prevalence of SD was 65.4% in women and 50% in men. Compared to controls, patients with male gender had significantly more impaired erection and orgasm ($p=0.005$; $p=0.002$ respectively), and those with female gender had significantly more impaired sexual arousal and desire ($p=0.003$; $p=0.028$ respectively). In the sample of patients, having a poor marital harmony and a fewer sexual attraction towards partner were correlated with decreased desire ($p=0.017$) in men and with sexual arousal ($p=0.024$) and decreased desire ($p=0.048$) in women. The number of relapses negatively affects erection ($p=0.038$) and orgasm ($p=0.048$). Depression correlated with a decreased orgasm ($p=0.001$) and desire ($p=0.048$) in men, and with a decreased sexual arousal ($p=0.006$) in women.

Conclusions: SD is common in CIBD, hence the need for a multidisciplinary approach to allow improvement of the quality of life of these patients, and of their partners.

Keywords: Chronic Inflammatory Bowel Diseases; Sexual Dysfunction; quality of sexual life