

Letters to the Editor

Interpretation of Blood Culture Results

To the Editor:

We read with interest the article by Lorian and Amaral published in *the* May issue of *Infection Control and Hospital Epidemiology*.¹ Although the problems related to the interpretation of blood culture results are well explained, there are a few imprecisions in the example given to illustrate the predictive value of positive blood culture.

The chances of detecting a positive blood culture depend on the sensitivity of the method employed and not on its specificity. Because the sensitivity of these cultures is less than 100%, not all episodes of bacteremia are actually detected (false-negative results). The value of 99% used in the example actually refers to the sensitivity of three blood culture sets for the detection of bacteremia found by Washington and Ilstrup.² The sensitivity of a single blood culture is considerably lower (around 80%),² and thus, if a single blood culture is obtained from 1,000 patients of whom 10% are truly bacteremic, only 80% true-positive cultures should be expected.

Pablo Yagupsky, MD
Ben Gurion University of the Negev
Beer Sheva, Israel

REFERENCES

1. Lorian V, Amaral L. Predictive value of blood cultures. *Infect Control Hosp Epidemiol*.1992;13:293-294.

2. Washington JA, Ilstrup DM. Blood cultures: issues and controversies. *Rev Infect Dis*.1986;8:792-802.

To the Editor:

With much interest I read the article "Predictive Value of Blood Cultures" by Dr. Victor Lorian in *Infection Control and Hospital Epidemiology*.¹

Some figures seem to not be quite correct. If the rate of bacteremia was 3%, the number of positive cultures would be 30 minus 0.3 (1%) = 29.7, divided by 49.7, which gives a predictive value of 59.7%.

J. De Jong, MD
Hospital Eemland
Amersfoort. The Netherlands

REFERENCE

1. Lorian V, Amaral L. Predictive value of blood cultures. *Infect Control Hosp Epidemiol*.1992;13:293-294.

The authors reply

We are sorry for the calculation error, but we also note that, even with the error, the results remain the same.

V. Lorian, MD
L. Amaral, MD
Bronx-Lebanon Hospital Center
Bronx, New York

Transmission of *Clostridium difficile*

To the Editor:

Brooks and colleagues reported data which suggested

that the rectal route may be important in the transmission of *Clostridium difficile* in their acute care and skilled nursing care facilities.¹ They were able to significantly reduce the incidence of *C difficile*-associated diarrhea by replacing the use of electronic thermometers with single-use disposable thermometers. A total of 20.8% of the electronic rectal thermometer handles were contaminated with *C difficile*.

A cluster of five cases of *C difficile*-associated diarrhea prompted an investigation in a chronic facility of 200 beds where this was a rare occurrence. Four additional cases were uncovered. These cases occurred in one hospital wing where rectal probes were used for temperature recording. There were no cases of *C difficile*-associated diarrhea in the other wings where disposable glass thermometers were used. Because rectal probes were suspected as the common vehicle of transmission, each probe was cultured on three sites: the probe tip, the surface of the probe covered by disposable probe cover, and the handle of the probe. The swabs were put into chopped meat glucose broth to look specifically for anaerobes. Although *C difficile* was not specifically cultured, all parts of the probes were found to be contaminated with fecal flora (Table) despite cleaning with alcohol wipes between patients and the use of disposable rectal sheaths. The use of rectal probes was therefore discontinued. No