

# ***Jimaodanus*, a replacement name for the algal genus *Heterocladus***

## **LoDuca, Kluessendorf, and Mikulic, 2003**

Pu Huang\* 

Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China <[puhuang@nigpas.ac.cn](mailto:puhuang@nigpas.ac.cn)>

The fossil genus *Heterocladus* LoDuca, Kluessendorf, and Mikulic, 2003 was established based on specimens from the Silurian (Llandovery) of southeastern Wisconsin, USA (LoDuca et al., 2003). This genus is monotypic and contains only the type species *Heterocladus waukeshaensis* LoDuca, Kluessendorf, and Mikulic, 2003, regarded as a noncalcified dasycladalean alga (LoDuca et al., 2003). *Heterocladus waukeshaensis* shows a narrow, undivided main axis and two types of laterals. Hair-like, branched laterals are arranged in whorls at lower part of the main axis, whereas clavate, undivided laterals appear at the upper part (LoDuca et al., 2003). The generic name is derived from the Greek *hetero* and *klados*, meaning ‘different’ and ‘branches,’ respectively (LoDuca et al., 2003).

However, *Heterocladus* LoDuca, Kluessendorf, and Mikulic, 2003 is a later homonym of *Heterocladus* Turczaninow, 1847, a name that was validly published based on an extant angiosperm (Turczaninow, 1847). *Heterocladus* Turczaninow, 1847 is currently regarded as a taxonomic synonym of *Coriaria* Linnaeus, 1753 (Good, 1930; Skog, 1972).

According to Articles 11.3 and 53.1 of the International Code of Nomenclature for Algae, Fungi, and Plants (Shenzhen Code) (Turland et al., 2018), *Heterocladus* LoDuca, Kluessendorf, and Mikulic, 2003 should not be used because this name is illegitimate. Hence, the replacement name *Jimaodanus* is proposed here to replace *Heterocladus* LoDuca, Kluessendorf, and Mikulic, 2003. The replacement name is derived from *jīmáodānzi*, the Chinese name of a feather duster, referring to morphological characters of the laterals.

### **Systematic paleontology**

Genus *Jimaodanus* nom. nov.

2003 *Heterocladus* LoDuca, Kluessendorf, and Mikulic, p. 1152, fig. 1.

**Type species.**—*Jimaodanus waukeshaensis* (LoDuca, Kluessendorf, and Mikulic) comb. nov., from Wisconsin of USA, monotypy; figured by LoDuca et al. (2003, fig. 1).

**Diagnosis.**—See LoDuca et al. (2003, p. 1152).

**Occurrence.**—Brandon Bridge Formation, Silurian (Llandovery); Waukesha Lime and Stone Company quarry, north of Waukesha, Wisconsin, USA.

\*Corresponding author.

**Etymology.**—From *jīmáodānzi*, the Chinese name of feather duster, referring to morphological characters of the laterals.

**Remarks.**—The genus *Heterocladus* LoDuca, Kluessendorf, and Mikulic, 2003 is a later homonym of *Heterocladus* Turczaninow, 1847. Therefore, the replacement generic name *Jimaodanus* is proposed to replace *Heterocladus* LoDuca, Kluessendorf, and Mikulic, 2003.

*Jimaodanus waukeshaensis* (LoDuca, Kluessendorf, and Mikulic, 2003) comb. nov.

2003 Basionym: *Heterocladus waukeshaensis* LoDuca, Kluessendorf, and Mikulic, p. 1155, fig. 1-1, 1-2.

**Holotype.**—Field Museum of Natural History, FMNH PP 45959a, b, Wisconsin, USA.

**Diagnosis.**—See LoDuca et al. (2003, p. 1155).

**Occurrence.**—Brandon Bridge Formation, Silurian (lower Llandovery); Waukesha Lime and Stone Company quarry, north of Waukesha, Wisconsin, USA.

**Description.**—See LoDuca et al. (2003, p. 1155).

**Remarks.**—The species *Heterocladus waukeshaensis* LoDuca, Kluessendorf, and Mikulic, 2003 was illegitimate due to the generic name used previously for the extant angiosperm. Hence, a new combination is here proposed.

### **Acknowledgments**

I thank J. Xiu of Friedrich Schiller University Jena for providing important literature. The work was supported by the National Natural Science Foundation of China (grant no. 41802003).

### **Declaration of competing interests**

The author declares none.

### **References**

- Good, R.D.O., 1930, The geography of the genus *Coriaria*: New Phytologist, v. 29, p. 170–198.  
Linnaeus, C., 1753, Species Plantarum, Volume 2: Stockholm, Laurentius Salvius, 1037 p.  
LoDuca, S.T., Kluessendorf, J., and Mikulic, D.G., 2003, A new noncalcified dasycladalean alga from the Silurian of Wisconsin: Journal of Paleontology, v. 77, p. 1152–1158, [https://doi.org/10.1666/0022-3360\(2003\)077<1152:ANNDAF>2.0.CO;2](https://doi.org/10.1666/0022-3360(2003)077<1152:ANNDAF>2.0.CO;2).



- Skog, L.E., 1972, The genus *Coriaria* (Coriariaceae) in the western Hemisphere: *Rhodora*, v. 74, p. 242–253.
- Turczaninow, N., 1847, Decas tertia generum adhuc non descriptorum, adjectis discriptionibus nonnullarum specierum Myrtacearum xerocarpicarum atque Umbelliferarum imperfectarum: *Bulletin de la Société Impériale des Naturalistes de Moscou*, v. 20, p. 148–174.
- Turland, N.J., Wiersema, J.H., Barrie, F.R., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Kusber, W.H., Li, D.Z., Marhold, K., May,

T.W., McNeill, J., Monro, A.M., Prado, J., Price, M.J., and Smith, G.F., 2018, International Code of Nomenclature for Algae, Fungi, and Plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017 (Regnum Vegetabile 159): Glashütten, Germany, Koeltz Botanical Books, 254 p.

Accepted: 14 March 2024