

2. We standardised the blood test parameters that are required following admission to our PICU ward and measured if all relevant blood tests\* were completed during admission.
3. Re-audit after implementation of a standardised policy.\* (Full blood count, urea and electrolytes, liver function tests and lipid profile, CRP, thyroid function tests, vitamin B12, folate, prolactin, vitamin D, calcium and HbA1c%).

**Methods.** First cycle is an audit of electronic patient records for all admissions to the PICU ward in a two-month period by using an audit proforma to capture information to answer the questions above.

After implementing a standardised guideline for PICU, second cycle was conducted over a two-month period for the re-audit.

**Results.** In the first cycle, a total of 20 patient records were audited. 30% had bloods taken within 24 hours of admission to PICU. 10% of patients had bloods taken on their previous ward. 40% of patients declined/unsafe, and 20% were not done in the timeframe.

69% of patients directly admitted to PICU had all the relevant admission blood tests completed and 23% had some of the blood tests done. These numbers were lower for the patients transferred to PICU from other wards.

In the second cycle, a total of 16 patient records were audited. There was an improvement in number of patients having bloods done within 24 hours of admission with only 6% not done in the timeframe.

Of the patients directly admitted to PICU, all the consenting patients (89%) had all the standardised blood tests. These figures remained lower for those patients transferred from other wards without an indication to retest the bloods.

**Conclusion.** The standardised guideline for PICU that was implemented after the first cycle consisted of a pre-filled blood test request form which included all relevant parameters detailed above.

This had a positive impact with a significant percentage increase in the total number of patients directly admitted to PICU having all relevant admission blood tests completed at re-audit (77% to 100%).

There was also a significant percentage increase (80% to 94%) of bloods taken within 24 hours of admission to PICU.

The next step will be to implement these pre-filled forms across all wards in the KMPT trust.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

## Prescription of High Dose of Antipsychotics in Two Rehabilitation Wards of Chronically Unwell Patients

Dr Danae Liolitsa<sup>1\*</sup> and Dr Azmathulla Khan<sup>2</sup>

<sup>1</sup>Cygnets Lodge Kenton, London, United Kingdom and <sup>2</sup>Cygnets Hospital Harrow, London, United Kingdom

\*Corresponding author.

doi: 10.1192/bjo.2023.442

**Aims.** The Royal College of Psychiatry (RCPsych) Consensus and the NICE guidelines have set out quality standards for the prescription of antipsychotics. There are concerns, however, regarding the over prescription across different psychiatric units: doses above BNF limits, multiple antipsychotics prescribed, lack of adequate monitoring of physical health and lack of availability of a clear rationale when high dose antipsychotic therapy (HDAT) is used. In the current audit, we sought to review the frequency HDAT is used in a female rehabilitation ward of patients with psychosis and a male rehabilitation ward of patients with

dual diagnosis psychosis and/or autism. We also examined whether appropriate considerations are taken prior to prescribing HDAT and if the mandatory monitoring is in place when someone is on HDAT.

**Methods.** A set of standards was set out and an audit tool was formulated that took the format of a table for data collection. Data were collected and anonymised, by looking at the medication charts and patients' records, to identify if essential steps prior to/during HDAT prescribing are followed. All service users, 10 in each ward, that were in admission during the period between 7th and 13th November 2022, were included in the study.

Data were compared directly with standards set out based on NICE guidelines and the RCPsych consensus. The data were input into Excel spreadsheet, then analysed and presented in tabular and graphs' format.

**Results.** A total of 20 medication charts were reviewed over the five-day period. Only one out of ten individuals with ASD in Springs Centre was treated on HDAT and a clear rationale was documented to justify its use. All patients in both wards had the HDAT form and side effects scale form completed and filed. All patients had documentation of HDAT use on T2/T3 forms.

40% of patients at Kenton Lodge were on HDAT. These patients required use of at least one depot due to lack of compliance. There was four patients overall treated on combination of antipsychotics. One patient did not have physical health monitoring at baseline due to refusing consent.

**Conclusion.** In a small proportion of cases, HDAT may be justified, as long as the safety implications are considered and monitoring requirements observed.

We recommend that audits of high dose antipsychotic prescribing can be performed periodically as a matter of routine practice. Also, to introduce a formal psychosis assessment scale to identify whether improvement has been accomplished since HDAT initiation.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

## The Management of Short-Term Insomnia for Inpatients in an Adult Acute Admissions Ward: Audit of Compliance With Local and National Guidelines

Dr Marie Long\* and Dr Rose Widdowson

Inverclyde Royal Hospital, Glasgow, United Kingdom

\*Corresponding author.

doi: 10.1192/bjo.2023.443

**Aims.** This study sought to establish local adherence to the following recommendations for the management of short-term insomnia (under 3 months duration): 1. Sleep hygiene advice should be utilised prior to pharmacological management. 2. A non-benzodiazepine hypnotic medication should be prescribed. 3. A one-off trial dose of benzodiazepine or Z-drug should be prescribed prior to regular prescription. 100% of patients reviewed by medical staff for insomnia are expected to meet the set standards.

**Methods.** Retrospective analysis of a cohort of patients (n=21), in an Adult Acute Admissions Ward in the West of Scotland. Taken over seven days in January 2023. Electronic prescriptions and chronological notes were reviewed in order to identify patients reviewed by medical staff for short-term insomnia.

**Results.** Of the patients reviewed for insomnia (n=8):

- 12.5% were offered sleep hygiene advice (n=1).