

estimate adjusted odds ratios (OR) and 95% confidence intervals (CI) after adjustment for age, sex and HIV status as potential confounders. RESULTS/ANTICIPATED RESULTS: Among 466 HHCs, the median age was 29 years (IQR 23-38), 58.8% were female, 3.4% were HIV-positive, and median BMI was 20.9 kg/m² (IQR 18.9-23.8). Overall, 329 HHCs (70.6%) had LTBI, 26 (5.6%) had DM and 73 (15.7%) had pre-DM. Compared to HHC without DM, the prevalence of LTBI was higher in those with pre-DM (68.9% vs. 72.6%; OR 1.19, 95% CI 0.69-2.13) and those with DM (88.5%; OR 3.45, 95% CI 1.17-14.77). In multivariable analysis, there was a trend towards increased LTBI risk among HHCs with DM vs. without DM (OR 2.16, 95% CI 0.67-9.70) but the difference was not statistically significant. Among HHCs with LTBI, the median IFN- γ response to TB1 antigen was modestly greater in those with DM (5.3 IU/mL; IQR 3.0-7.8) and pre-DM (5.4 IU/mL; IQR 2.0-8.4) compared to HHCs without DM (4.3 IU/mL; IQR 1.4-7.7). DISCUSSION/SIGNIFICANCE OF FINDINGS: Our results suggest that DM may increase the risk of LTBI among HHCs recently exposed to active TB. Among those with LTBI, increased IFN- γ antigen response in the presence of DM and pre-DM may indicate an exaggerated but ineffectual response to TB. Further investigation is needed to assess how dysglycemia impacts susceptibility to *M. tuberculosis*.

67863

Insulin use and depigmentation: a survey of real-world evidence

James Lukasik¹, Zachary A.P. Wintrob^{1,2}, G. Emilia Costin^{1,3} and Alice C. Ceacareanu^{1,2}

¹Hartwick College, ²ROAKETIN Inc. and ³Institute for In Vitro Sciences

ABSTRACT IMPACT: Long-acting insulin containing protamine is more likely to be associated with skin depigmentation. OBJECTIVES/GOALS: An acquired disorder, skin depigmentation was found to be significantly correlated with diabetes. While a recent meta-analysis pointed at a possible similar pathogenesis, the possibility of vitiligo occurring as a drug-induced disease was never explored. This study aimed at elucidating whether utilization of specific insulins may play a role. METHODS/STUDY POPULATION: Records from the Medical Panels Expenditure Survey (MEPS) database made available by the Agency for Healthcare Research and Quality were used to identify all injectable insulin users (n=8867). ICD-9/10 codes were abstracted from the medical conditions files for all the subjects reporting any type of injectable insulin use (1996-2017). Skin depigmentation codes identified in our dataset were 709 and L81. Insulins were categorized based on duration of action, short-acting (SA), intermediate-acting (IA), and long-acting (LA), as well as based on formulation ingredients (zinc, protamine-zinc, other), and insulin combination (SA with or without IA/LA containing or not protamine-zinc). The association between skin depigmentation occurrence and insulin type and/or category was assessed by Fisher's exact test. RESULTS/ANTICIPATED RESULTS: A total of 225 out of 8867 individuals were diagnosed with skin depigmentation. Incidence of skin depigmentation was 2.25% in SA users (n1=3606, p=0.355), 2.24% in LA users (n2=3910, p=0.337), and 2.39% in IA users (n3=4015, p=0.062). Occurrence of skin depigmentation was similar between users of insulin mono- or combo therapy (p=0.758). Interestingly, among IA insulins, insulin protamine-zinc insulin (n4=3992) distinguished as being mainly responsible for the association with the occurrence of skin depigmentation (p=0.062), whereas insulin zinc was not (n5=37, p=1.000). The highest skin depigmentation incidence was observed among Pacific Islanders (2.66%, p=0.110). Males distinguished by a skin depigmentation incidence of 2.34% vs. 1.91% in females (p=0.086). DISCUSSION/

SIGNIFICANCE OF FINDINGS: We report that presence of protamine-zinc may play a role in the development of skin depigmentation. It is uncertain whether this risk may be shared equally by insulin users diagnosed with type 1 and type 2 diabetes. Of note, we observed a higher skin depigmentation incidence than that reported by community- (0.2%) or hospital-based (1.8%) studies.

68127

High Sensitivity Troponins Predicts Mortality in Patients Who Present to the ED with Severe Sepsis or Septic Shock*

Kendrick Williams¹, Ryan Tucker, MD², James Cranford, PhD² and Christopher Fung, MD²

¹University of Michigan Medical School and ²Michigan Medicine Department of Emergency Medicine

ABSTRACT IMPACT: Our may suggest that delta hsTrop could be of prognostic value in patients with sepsis. OBJECTIVES/GOALS: - METHODS/STUDY POPULATION: We analyzed data of those presenting to the ED over an 18-month period with sepsis and at least one episode of hypotension after 1 liter of IV fluids. We performed a retrospective analysis using a cohort derived from modified inclusion and exclusion criteria from the CLOVERS study. The outcomes of patients found to have a delta (at least 6 pg/dL) in high sensitivity troponin T were compared to patients who did not have a delta or have a troponin level measured. We examined demographic and treatment characteristics of this cohort and the incidence of adverse outcomes were determined. We used multivariable logistic regression analysis to test the association of hsTrop and mortality. RESULTS/ANTICIPATED RESULTS: 778 patients met criteria to be included in the cohort. 279 patients had a change in high sensitivity troponins, an incidence of 35.9%. Patients with a delta were more likely to be older, male, and have a higher Charlson index than patients without a delta or those that had no troponin measured. They were also more likely to have a history of chronic lung disease, heart failure and hypertension. Change in high sensitivity troponins were associated with higher in-hospital mortality. When adjusted for age, gender, and Charlson Index, the association between a positive delta troponin and mortality remained statistically significant. DISCUSSION/SIGNIFICANCE OF FINDINGS: In patients with severe sepsis and septic shock, the presence of a positive or negative delta hsTrop at 2 hours is associated with increased mortality. Measurement of high sensitivity troponin early in the patient's hospital course may have prognostic utility.

75561

Association of childhood hypertension with early adulthood obesity and hypertension*

Mindy Pike, T. Alp Iikizler, Loren Lipworth and Cassianne Robinson-Cohen

Vanderbilt University Medical Center

ABSTRACT IMPACT: This study establishes the association between childhood hypertension and health outcomes in early adulthood, identifying the need to understand blood pressure during early life for primary prevention of hypertension and cardiovascular disease. OBJECTIVES/GOALS: There is evidence that blood pressure level in early life can influence hypertension and other cardiovascular risk factors later in life. We examined whether hypertension before the age of 18 is associated with higher odds of obesity and hypertension after the age of 18. METHODS/STUDY POPULATION: We studied 19,367