
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editors, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
395A Hatfield Road,
St Albans, Herts AL4 0XQ.
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

ASHFORD, R. W. & BARNISH, G. (1989) *Strongyloides fuelleborni* and similar parasites in animals and man. In: *Strongyloidiasis: a major roundworm infection of man* (editor, D. I. Grove) pp. 271–286. Taylor & Francis: London.

GARSDIE, P. & BEHNKE, J. M. (1989) Primary infection of hamsters with *Ancylostoma ceylanicum*. *Parasitology*, **98**, 283–289.

Contents

	Pages
The mouse as a suitable host for an isolate of <i>Schistosoma haematobium</i> from Niger. D. IMBERT-ESTABLET, C. VERA, B. SELLIN and J. JOURDANE	1-5
An insect, <i>Sialis lutaria</i> , as a host for <i>Proteocephalus</i> sp. C. R. KENNEDY, P. NIE, and J. ROSTRON	7-16
<i>Proteocephalus neglectus</i> as a possible indicator of changes in the ecological balance of aquatic environments. V. HANZELOVÁ	17-24
Molecular cloning of the cDNA encoding a 42 kDa antigenic polypeptide of <i>Anisakis simplex</i> larvae. K. SUGANE, S. SUN and T. MATSUURA	25-32
The identity of <i>Anisakis</i> type II larvae with <i>Anisakis physeteris</i> confirmed by restriction fragment length polymorphism analysis of genomic DNA. T. MATSUURA, S. SUN and K. SUGANE	33-37
<i>Trichinella spiralis</i> : specificity of ES antigens from pre-encysted larvae. R. C. KO and T. P. WONG	38-44
On the biological and biochemical nature of cloned populations of <i>Strongyloides ratti</i> . M. E. VINEY, B. E. MATTHEWS and D. WALLIKER	45-52
Life cycle of <i>Gnathostoma nipponicum</i> Yamaguti, 1941. K. ANDO, H. TOKURA, H. MATSUOKA, D. TAYLOR and Y. CHINZEI	53-61
Stage-specific expression of a developmentally regulated gene in <i>Dirofilaria immitis</i> . S. SUN, T. MATSUURA and K. SUGANE	62-67
Effects of patent <i>Echinostoma caproni</i> infection on the sterol composition of the digestive gland-gonad complex of <i>Biomphalaria glabrata</i> as determined by gas-liquid chromatography. P. H. SHETTY, B. FRIED and J. SHERMA (Research Note)	68-71
Seasonal variation in the prevalence and intensity of canine <i>Gnathostoma spinigerum</i> infection in northeastern Thailand. W. MALEEWONG, S. PARIYANONDA, P. SITTHITHAWORN, W. DAENGSEEGAWE, V. PIPITGOOL, S. TESANA, C. WONGKHAM, P. INTAPAN and N. MORAKOTE (Research Note)	72-74
Seroprevalence of <i>Toxocara canis</i> antibodies in humans in northern Jordan. M. N. ABO-SHEHADA, L. SHARIF, S. N. EL-SUKHON, N. ABUHARFEIL and R. F. ATMEH (Research Note)	75-78
Errata	78