

Introduction: Obsessive–Compulsive Disorder (OCD) is a disabling and chronic illness defined by the presence of obsessions and/or compulsions. Recently it has been proposed that the perinatal period may act as a trigger in this disorder, leading to its onset or exacerbation.

Objectives: Variations in pregnancy-related hormones are believed to be one of the main etiological theories for the development of perinatal OCD (pOCD). Perhaps for that reason research has been almost exclusively focused on the development of this disorder in mothers. We aim to investigate pOCD in fathers.

Methods: A non-systematic review was conducted via electronic searches of PubMed. The keywords used were “Perinatal”, “Father”, “OCD”, “Obsessive-compulsive disorder”.

Results: Unwanted intrusive thoughts are experienced with a similar prevalence in mothers and fathers. The same seems to be true regarding compulsions. However, it does appear that mothers are more distressed by these symptoms, which tend to be baby-related, usually concerning themes of suffocation, accidents or contamination. It is hypothesized that this seemingly different impact is related to the fact that mothers are more often the primary caregivers than fathers, thus feeling more distress because they are imbued with a greater responsibility. Accordingly, pOCD symptoms tend to be more severe in fathers who consider their baby-related obsessions meaningful, often confusing them as a desire to carry out such thoughts. These findings are consistent with the Cognitive-Behavioral Theory of OCD, highlighting that purely biological theories for the development of pOCD might not suffice.

Conclusions: Research indicates a similar presence of OCD symptoms in postnatal mothers and fathers, although it seems that mothers may experience more distress. Underlying dysfunctional beliefs seem to be responsible for the negative appraisal of these symptoms, predicting the development of the disorder in question. Further research of pOCD should seek to better characterize the onset or exacerbation of this disorder in fathers.

Disclosure of Interest: None Declared

EPV0643

Antiandrogenic treatment of obsessive compulsive neurosis: A case review

L. Huerga García*, I. Careno Baez, G. Oropeza Hernández, A. Marcos Rodrigo, C. Delgado Torres, G. Garriga Rocío and P. Gómez Pérez

Psychiatry, Hospital Universitario Nuestra Señora de la Candelaria, Santa Cruz de Tenerife, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1961

Introduction: Obsessive-compulsive disorder (OCD) is a mental disorder in which patients who suffer from it have repetitive and undesirable thoughts, feelings, ideas, sensations (obsessions) and behaviors that drive them to do something over and over again (compulsions).

Often the person tries to get rid of the obsessive thoughts through compulsions, but this only provides short-term relief. Not carrying out the obsessive rituals can cause enormous anxiety and suffering.

Objectives: To describe a 23-year-old male patient, who suffers from anxiety and mood symptoms, reacts to ego-dystonic obsessive ideas and sexual content, of months of evolution, and who manages

to calm down through compulsive masturbation or watching sexual videos on the internet. All this clinic negatively interferes with their quality of life, asking the patient for medical help to calm these ideas.

Methods: We carried out a review in Pubmed with the terms Antiandrogens and TOC, in order to make a better description of the clinical case.

Results: After several treatment attempts (Sertraline, Paroxetine, Clomipramine, Clomipramine + SSRI), reaching maximum doses according to clinical guidelines, and with poor therapeutic response, it was decided to discuss the case with the endocrinology department of our hospital, deciding to start treatment with antiandrogens, in order to alleviate the persistent intrusive ideas of a sexual nature. The administration of antiandrogens in men can cause a decrease or increase in the development or involution of secondary sexual characteristics in men, reducing the activity or function of accessory sexual organs, and hyposexuality, with decreased sexual desire or libido.

After several weeks, there was improvement in the obsessive symptoms with a decrease in compulsive rituals. However, after the 3rd mo, some symptoms reappeared, but not with the same severity and intensity as before treatment. In addition, we cannot ignore the adverse effects that have occurred, such as involution of secondary sexual characteristics. However, and taking into account the negative repercussion that this clinic had on the patient's quality of life, the benefit obtained exceeded the risk, having noted clear improvement with this therapy, and maintaining evolutionary controls by both psychiatry and endocrinology.

Conclusions: Patients suffering from obsessive-compulsive disorder can be effectively treated with anti-androgenic pharmacological agents with various modes of action. The most effective group of such agents is the long-acting analogues of the gonadotropin-releasing hormone. The objective of this review is to elucidate the possibility of using such powerful anti-androgenic agents in the treatment of obsessive-compulsive disorder.

Disclosure of Interest: None Declared

EPV0644

Homosexual Obsessive–compulsive Disorder Comorbid with Bipolar Disorder: A Rare Case report

M. Gardabbou*, R. Feki, N. Smaoui, I. Gassara, S. Omri, M. Maalej Bouali, N. Charfi, J. Ben Thabet, L. Zouari and M. Maalej

psychiatry C department, Hedi Chaker, Sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1962

Introduction: While bipolar disorder–obsessive compulsive disorder overlap is quite common, sexuality remains a largely unexplored area of this clinical entity.

Objectives: Illustrate through a clinical vignette the case of a patient with diagnosed homosexual obsessive –compulsive disorder (OCD) comorbid with bipolar disorder (BD).

Methods: The clinical case report was prepared through the review of the patient's clinical record.

Results: We report a rare case of a 22 year-old man who was diagnosed with Homosexual Obsessive–compulsive Disorder comorbid with Bipolar Disorder, admitted to our department for a suicide attempt. He came from a religious and conservative