

tral response is available at the same time as initial response is required by emergency services.

**Discussion:** By developing such an organization, the benefits of providing early integrated response by medical coordinators ensures on-site medical control is established at the same time as first response occurs, proper liaison is affected with combat commanders, and that a correct assessment is made of the situation. In addition, the problem of inappropriate convergence of medical assistance is minimized, casualties can be distributed to appropriate hospitals more easily, resources can be arranged more easily, and preplanned site procedures can be adhered to by paramedics, hospital medical teams, mental health, and other responders required during the incident. The major factor emanating from providing such an organization is that when disaster strikes, the total medical and health response, which is not a discreet service, is able to work comfortably with other emergency services, because on a daily basis the medical coordinators are members of all relevant planning committees, work together during lesser incidents, and get to know personalities of other services.

#### 084. Physicians Guide for Extremity Trauma

Marcos Musafir, MD, Sergio J. Franco, MD, Paulo Pinheiro, MD,  
Savino Gasparini, MD  
Rio de Janeiro, Brazil

At the Miguel Couto Trauma Center in Rio de Janeiro, Brazil, the trauma team has four orthopedic surgeons and three residents on duty each 24 hours under Professor Nova Monteiro's leadership. Sometimes, all of them are at the operating theater, and other physicians need to take the initial care of extremity trauma. The hospital has 800 admittances at the emergency room per day, and 50% are not in need of emergency orthopedics (28% of these patients each day).

Concerning this, we prepared a training guide including how to recognize the severity of one extremity trauma either closed or open fracture and the initial management of these lesions by non-orthopedic surgeons.

The extremity trauma care is done after the ABCs.

The rule 1 is LOOK, FEEL, and MOVE.

The rule 2 is CLEAN and PROTECT the wounds.

The rule 3 is PREVENT further injury.

The rule 4 is how to IMMOBILIZE.

The rule 5 is MEDICATION and X-RAY examinations.

The authors will distribute this practical guide during the Congress. It has worked very well in Rio, reducing dramatically (70%) of the complications of some extremity injuries when they initially are managed by other physicians at the hospital. This guide also is now in use for the paramedics, working in prehospital care.

#### 035. Fluid Therapy with Middlemolecular Hydroxyethyl Starch (HES 200/0,5) in Surgery, Trauma, Sepsis, and Burns

Michael G. Beez

Pharm. Int. Dept. FRESENIUS AG  
Homburg, Germany

HAES-steril® is a colloidal plasma volume substitute and, therefore, increases the plasma volume, thus improving cardiac output and oxygen transport values. As a result, HAES-steril® is capable of improving organ functions and the overall hemodynamic situation in patients with hypovolemia/shock.

HAES-steril® 6% is used for medium-term, non-expansive, volume replacement therapy in routine surgical patients. Since the hemodynamic efficacy of HAES-steril® 6% is similar to that of 5% human albumin of Plasma Protein Fraction, the use of HAES-steril® in hypovolemic/shock patients substantially cuts down the albumin and PPF consumption.

HAES-steril® 10% is used for medium-term expansive volume replacement therapy in the above patient population with hypovolemia and shock if the overriding therapeutic goal is to achieve a more rapid and more extensive expansion of the plasma volume and a more profound improvement in hemodynamic variables, microcirculatory flow, and oxygen delivery. Examples include ICU patients with massive acute blood losses, surgical patients with protracted shock and disturbed microcirculation and/or those at high risk for developing pulmonary thromboembolism (PTE). HAES-steril® 10%, too, saves considerable amounts of albumin in the treatment of hypovolemic/shock patients.

Nearly all clinical trials of HAES-steril® showed significant improvements in arterial pressures (MAP, PAP), cardiac index (CI), oxygen delivery (DO<sub>2</sub>), and oxygen consumption (VO<sub>2</sub>), as well as in organ functions.

Thus, HAES-steril® 10% and HAES-steril® 6% meet Shoemaker's criteria (cf. section on Hypovolemia/Shock) of reducing the morbidity and mortality of critically ill patients.

The efficacy of HAES-steril® 10% and HAES-steril® 6% proved similar and/or superior to the efficacy of 5% albumin.

The effects of HAES-steril® 10% and HAES-steril® 6% proved superior to the effects of crystalloid solutions. Crystalloid solutions, even when given in a multiple overdose, are not suitable for improving hemodynamics and oxygen transport in patients with hypovolemia and shock.

#### 021. Management of Extremity Injuries in Extreme Circumstances

Endre Cziffer, MD, PhD

Department of Traumatology, Central Army Hospital,  
Budapest, Hungary

External fixation is in its present renaissance due to the rapidly rising frequency of severe road accidents, civilian and military mass casualties, local wars, and terrorist attacks. The goal for