NOTES FOR AUTHORS

The Bulletin of Entomological Research publishes original research papers concerning insects, mites, ticks or other arthropods of economic importance in agriculture, forestry, stored products, biological control, medicine, animal health and natural resource management. The geographical scope of the Bulletin is worldwide but with emphasis on the tropics. Taxonomic papers are accepted if relevant. Short review papers, although normally by invitation, will also be considered for publication.

Page Format. The *Bulletin* is printed in a twocolumn format (column width of 80 mm) with a text area of 170×225 mm.

Text. Papers should be typed, on one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side and with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used. Guidelines for taxonomic papers are available.

When the paper has been accepted word-processed text stored on floppy disk is encouraged, providing the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with fewer type-setting errors.

Abstract. Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words. A short title should be provided as a running head.

Tables. Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or land-scape (if absolutely necessary) format. Table captions should be typed on a separate sheet.

Illustrations. Copies only of artwork should be submitted. The original illustrations should accompany the paper after acceptance and revision. Text figures, line drawings, computer-generated figures and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Half-tone photographs are acceptable where they are a real contribution to the text. Figure and captions should be typed on a <u>separate sheet</u> in the following format:

Figs 23–26. Figs 23–24, <u>Urophora</u> eggs: 23, <u>U.</u> <u>hispanica</u>; 24, <u>U. stigma</u>. Figs 25–26, spermathecae: 25, <u>U. maura</u>; 26, <u>U. stigma</u>; scale lines = 0.05 mm.

Voucher specimens. The deposition of voucher specimens should be considered where appropriate.

References. References must be based on the name and year system, give full journal titles and conform to the following styles:

Powell, W. (1986) Enhancing parasitoid activity in crops. pp. 319–340 <u>in</u> Waage, J. & Greathead, D. (<u>Eds</u>) <u>Insect parasitoids</u>. London, Academic Press (Symposium, Royal Entomological Society of London No. 13).

Southwood, T.R.E. (1978) <u>Ecological methods with</u> <u>particular reference to the study of insect populations</u>. 2nd edn. 524 pp. London, Chapman & Hall.

Zhou, X., Carter, N. & Mumford, J. (1989) A simulation model describing the population dynamics and damage potential of the rose grain aphid, <u>Metopolophium dirhodum</u> (Walker) (Hemiptera: Aphididae), in the UK. <u>Bulletin of Entomological Resear</u>ch 79, 373–380.

Citation of authors in the text should appear in the form: Polaszek (1990) or (Polaszek, 1990). More than one author should be cited in chronological order as: (Holloway <u>et al.</u>, 1987; Walker & Huddleston, 1988).

Offprints. 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

Manuscripts. Three copies of the manuscript and artwork should be submitted to:

The Editors Bulletin of Entomological Research International Institute of Entomology 56, Queen's Gate London SW7 5JR, UK.

JUNE 1993

Bulletin of Entomological Research

Baylis, M. & Nambiro, C. O. The responses of <i>Glossina pallidipes</i> and <i>G. longipennis</i> (Diptera: Glossinidae) to odour-baited traps and targets at Galana Ranch, south-eastern Kenya	145
Bigger, M. Time series analysis of variation in abundance of selected cocoa insects and fitting of simple linear predictive models	153
Boakye, D. A., Post, R. J., Mosha, F. W., Surtees, D. P. & Baker, R. H. A. Cytotaxonomic revision of the Simulium sanctipauli subcomplex (Diptera: Simuliidae) in Guinea and the adjacent countries including descriptions of two new species	171
Feng, M. G., Nowierski, R. M. & Zeng, Z. Binomial sampling plans for the English grain aphid, Sitobion avenae (Homoptera: Aphididae) based on an empirical relationship between mean density and proportion of tillers with different tally thresholds of aphids	187
Hargrove, J. W. Target barriers for tsetse flies (<i>Glossina</i> spp.) (Diptera: Glossinidae): quick estimates of optimal target densities and barrier widths	197
Hollis, D. & Martin, J. H. <i>Padaukia</i> – a new genus of jumping plantlice (Homoptera: Psylloidea) attacking economically important species of <i>Pterocarpus</i> (Leguminosae)	201
Lomer, C. J., Stride, A. B. & Balfas, R. The biology of <i>Hindola</i> spp. (Homoptera: Machaerotidae) vectors of Sumatra disease of clove and some related species in Indonesia	213
Polaszek, A., Ubeku, J. A. & Bosque-Perez, N. A. Taxonomy of the <i>Telenomus busseolae</i> species- complex (Hymenoptera: Scelionidae) egg parasitoids of cereal stem borers (Lepidoptera: Noctuidae, Pyralidae)	221
Seth, R. K. & Reynolds, S. E. Induction of inherited sterility in the tobacco hornworn <i>Manduca sexta</i> (Lepidoptera: Sphingidae) by substerilizing doses of ionizing radiation	227
Shanower, T. G., Schulthess, F. & Bosque-Perez, N. Development and fecundity of <i>Sesamia</i> calamistis (Lepidoptera: Noctuidae) and <i>Eldana saccharina</i> (Lepidoptera: Pyralidae)	237
Simandl, J. Influence of temperature on larval defecation and its use in estimating canopy population size in pine sawflies (Hymenoptera: Diprionidae)	245
Tamo, M., Baumgärtner, J., Delucchi, V. & Herren, H. R. Assessment of key factors responsible for the pest status of the bean flower thrips <i>Megalurothrips sjostedti</i> (Thysanoptera: Thripidae) in West Africa	251
Tingle, C. C. D. Bait location by ground foraging ants (Hymenoptera: Formicidae) in mopane woodland selectively sprayed to control tsetse fly (Diptera: Glossinidae) in Zimbabwe	259
Upton, M. S. Aqueous gum-chloral slide mounting media: an historical review	267
Vale, G. A. Visual responses of tsetse flies (Diptera: Glossinidae) to odour-baited targets	277
Wilson, L. J. Seasonal abundance and distribution of <i>Tetranychus urticae</i> (Acari: Tetranychidae), the two spotted spider mite, on cotton in Australia and implications for management	291
Book Reviews	305

© C-A-B International, 1993

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.

https://doi.org/10.1017/S0007485300034623 Published online by Cambridge University Press by Information Press, Eynsham, Oxford.