

that work needs to be done to ensure that, in every case of an inpatient needing to be subject to HDAT, the clinical rationale for this is documented. Capacity to consent to being prescribed HDAT must be documented for each HDAT patient as a matter of good clinical practice.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

An Audit of Whether Patients on the General Adult Inpatient Wards and the PICU in Mersey Care NHS Foundation Trust Who Are Prescribed High Dose Antipsychotic Therapy Are Being Monitored as per Trust Policy

Dr Declan Hyland*, Dr Roopa Singh and Mrs Kerry Dainton
Mersey Care NHS Foundation Trust, Liverpool, United Kingdom
*Presenting author.

doi: 10.1192/bjo.2024.580

Aims. High Dose Antipsychotic Therapy (HDAT) should only be used in exceptional circumstances, as there is little evidence to suggest that higher than recommended doses of antipsychotics are more clinically effective than standard doses, with potential side effects being greater. In practice, there are several clinical scenarios where HDAT may be prescribed and the potential benefits must outweigh the potential risks. NICE guidelines for psychosis and schizophrenia advise that dosages outside the range given in the British National Formulary should be justified and recorded.

This audit aimed to determine whether patients on the 16 general adult inpatient wards and Psychiatric Intensive Care Unit (PICU) in Mersey Care NHS Foundation Trust who are prescribed HDAT are managed as recommended by the Trust's HDAT policy.

Methods. A list of all inpatients admitted to the 16 general adult inpatient wards and the PICU in the Trust between 17th and 20th of July 2023 was obtained. The electronic prescription record for each patient was scrutinised to determine whether the patient was prescribed HDAT. For each HDAT patient, the patient's electronic psychiatric record was analysed to determine whether baseline physical health assessments – ECG, BMI, waist circumference, BP, pulse rate, FBC, U and Es, LFTs, serum prolactin level, HbA1c level and random serum total cholesterol level and lipid profile were completed before commencing HDAT. Each HDAT patient was reviewed to determine whether a cardiovascular assessment was completed prior to commencing HDAT.

Results. 29 inpatients on the 16 general adult wards and the PICU were prescribed HDAT. The Trust's HDAT policy states 13 baseline physical health assessments should be completed prior to commencing HDAT. For the 29 patients, 13 baseline physical health assessments should have been completed on 378 occasions. On 98 occasions, the patient refused and these were excluded from the compliance. There were 226 (81%) occasions they were completed and 54 (19%) when they weren't completed. 12 (41%) of the 29 HDAT patients had a cardiovascular assessment done prior to commencing HDAT.

Conclusion. There are a significant number of inpatients in whom not all the required baseline physical health assessments prior to commencing HDAT are completed. A cardiovascular assessment is an important aspect of deciding whether to prescribe a patient HDAT and yet commonly not completed. There is a need to ensure that nursing and medical staff on the

inpatient wards are aware of the Trust's HDAT policy and need to refer to and to adhere to it.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

An Evaluation of the High Dose Antipsychotic Therapy Patients on the General Adult Inpatient Wards and the Psychiatric Intensive Care Unit in Mersey Care NHS Foundation Trust – Is Clozapine Being Considered in Each Case?

Dr Declan Hyland*, Dr Roopa Singh and Mrs Kerry Dainton
Mersey Care NHS Foundation Trust, Liverpool, United Kingdom
*Presenting author.

doi: 10.1192/bjo.2024.581

Aims. High Dose Antipsychotic Therapy (HDAT) should only be used in exceptional circumstances, as there is little evidence to suggest that higher than recommended doses of antipsychotics are more clinically effective than standard doses, with potential side effects being greater. In practice, there are several clinical scenarios where HDAT may be prescribed and the potential benefits must outweigh the potential risks. NICE guidelines for psychosis and schizophrenia advise that dosages outside the range given in the British National Formulary should be justified and recorded.

This audit aimed to determine whether the option of clozapine is being considered in those patients on the 16 general adult inpatient wards and Psychiatric Intensive Care Unit (PICU) in Mersey Care NHS Foundation Trust who are prescribed HDAT.

Methods. A list of all inpatients admitted to the 16 general adult inpatient wards and to the PICU in the Trust between 17th and 20th of July 2023 was obtained. The electronic prescription record for each patient was scrutinised to determine whether the patient was subject to HDAT and, if so, whether there was documentation in the patient's electronic record that the option of treatment with clozapine was considered. The authors also wished to determine whether, in those HDAT patients in whom clozapine was considered, the rationale for it not being pursued as a treatment option was documented.

Results. 29 inpatients on the 16 general adult wards and on the PICU were prescribed HDAT. In 9 (39%) of the HDAT patients, the option of treatment with clozapine was considered. It was documented for 6 of the HDAT patients that the option of clozapine was not applicable. Of the 9 HDAT patients that had a trial of clozapine considered, all of them had documented evidence of the decision in their electronic record. Four of the 9 patients accepted the trial of clozapine, 5 did not accept/it was deemed not appropriate. Of the 5 patients who did not accept the trial of clozapine or were deemed not appropriate, the rationale was documented for each patient.

Conclusion. Given the lack of recommendation and evidence base to prescribe HDAT, the option of clozapine, if appropriate, should always be considered in any patients in whom HDAT is being considered or required. There may be barriers to clozapine being chosen – both patient-related and clinician-related. Any such barriers should be explored and addressed to ensure that treatment-resistant patients are commenced on clozapine without unnecessary delay.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.