



proportion of male patients (IRR 1.12, 95%CI 1.10-1.14), Black patients (IRR 1.05, 95%CI 1.03-1.07), or Medicaid beneficiaries (IRR 1.15, 95% CI 1.12-1.17). Effect modification was observed between SDI and rurality, with higher SDI in non-metropolitan areas associated with higher claim rates, whereas SDI in metropolitan areas was inversely related to claim rates. **Conclusion:** This study showed that the distribution of high and low SDI and rates of fluoroquinolone claims were more geographically clustered than expected by random chance alone. Lower fluoroquinolone claim rates among Texas Medicare providers were seen in metropolitan areas with higher SDI, indicating potential barriers to care. Conversely, higher claim rates were observed in rural areas with higher SDI, signifying a possible knowledge or attitude gap towards fluoroquinolone use. These findings provide opportunities for public health professionals to explore gaps in the knowledge and attitudes of patients and providers related to antimicrobial use, particularly in rural regions, and investigate barriers to healthcare access in metropolitan areas.

Antimicrobial Stewardship & Healthcare Epidemiology 2024;4(Suppl. S1):s75-s76
doi:10.1017/ash.2024.205

Presentation Type:

Poster Presentation - Poster Presentation

Subject Category: DEI

Survey of Infectious Diseases and Infection Prevention Practitioners on Diversity, Equity, and Inclusion Experiences

Christopher Evans, Tennessee Department of Health; Erika Kirtz, Tennessee Department of Health; Rekha Murthy, University of Nebraska Medical Center; Cedars-Sinai, University of Nebraska Medical Center; Jasmine Marcelin, University of Nebraska Medical Center and Zanthia Wiley, Emory

Background: Incorporating diversity, equity, inclusion, and justice into healthcare ensures equitable opportunity to achieve optimal health. Infectious diseases, antimicrobial stewardship, and infection prevention teams rely on consultative recommendations to improve patient care which may be influenced by implicit and explicit biases of the recipient treatment teams. Little is known about how race, ethnicity, and other characteristics impact stewardship and infection control recommendations. **Methods:** A survey of infectious diseases, antimicrobial stewardship, and infection prevention practitioners was developed through the Society of Healthcare Epidemiology of America (SHEA) Antimicrobial Stewardship Committee. The survey was sent electronically to members of the SHEA Research Network and was promoted to attendees of two sessions at IDWeek 2022 and SHEA Spring 2023. Survey questions included demographics, awareness of (and participation in) unconscious bias and

microaggression training at their institutions, antibiotic prescribing bias observations, and perceptions of how race, ethnicity, and other characteristics have influenced participants' antimicrobial stewardship and infection prevention recommendations. Descriptive statistics were performed using SAS V.9.4. **Results:** Among 175 survey respondents, 75% (n=129) were White, 16% (n=27) were Asian, 4% (n=7) were Black, 85% (n=150) were non-Hispanic, 5% (n=8) were Hispanic, and 3% (n=5) reported ethnicity as multiethnic. 76% of respondents identified as female, and 2% as non-binary or gender-fluid. 29% of respondents had a medical degree, 12% had a nursing degree, 7% had a pharmacy degree, and 52% had a degree listed as other (7% had a PhD, 23% had an MPH/MSPH degree, and 15% had an MS degree). 65% and 49% of respondents had participated in unconscious bias and microaggression training, respectively. 18% (n=22) of White respondents, 43% (n=3) of Black respondents, and 30% (n=8) of Asian respondents reported witnessing antimicrobial prescribing influenced by race, ethnicity, or other characteristics. 17% and 15% of respondents felt that their antimicrobial stewardship and infection prevention recommendations, respectively, had not been accepted due to their race, ethnicity, gender identity, or other personal identifiers. **Conclusion:** This survey showed demographic characteristics of professionals working in infectious diseases and their perceptions of how certain aspects of their identity have influenced their recommendations. Differences between racial groups were observed in how frequently respondents witnessed inequities in antimicrobial prescribing, and many respondents felt their recommendations had not been accepted due to their identity. A limitation of this analysis is that few Black individuals completed the survey, which makes comparisons by race difficult; however, the respondents were consistent with SHEA membership demographics.

Antimicrobial Stewardship & Healthcare Epidemiology 2024;4(Suppl. S1):s76

doi:10.1017/ash.2024.206

Presentation Type:

Poster Presentation - Poster Presentation

Subject Category: DEI

Patient First Strategies for Reducing Inequities in HAI Prevention

Lisa Stancill, UNC Health; Emily Sickbert-Bennett Vavalle, UNC Health and Lauren DiBiase, UNC Health

Background: Inequities in healthcare-associated infections (HAI) incidence and prevention measures are critically important to understand (Chen,2021). While evaluations are beginning to characterize these disparities by infection type (Gettler, 2023), our work expands this by characterizing disparities by prevention strategies. By better understanding how evidence-based prevention strategies are implemented at the patient level, infection preventionists and hospital epidemiologists can better design strategies that provide equitable care to all patients. **Methods:** Beginning January 2023, gender, race, ethnicity, spoken language, and age group fields were added to daily chlorhexidine gluconate (CHG) treatment and C. difficile test order compliance data captured via electronic medical record. In July 2023, fields on recorded race, ethnicity, and gender were added to well-established foley and vascular access real-time peer audit tools that are used by infection preventionists (IPs). Each prevention strategy variable was summarized by demographic variables and differences in compliance were measured using chi-square tests. **Results:** 899 vascular audits and 420 foley audits were completed by IPs between July – December 2023. In 2023, there were 114,066 opportunities for CHG Treatment and 1,991 C. difficile test orders. Missing data varied by metric but ranged from 0-60%. Statistically significant differences by race were found in 3 of 8 components (i.e., intact seal, secured catheter and absence of dependent loop) in the foley audit ($p < 0.01$) and compliance with C. difficile test ordering ($p < 0.01$). No differences in race were found in vascular access audits or CHG treatment. No differences in gender or ethnicity were noted in foley, vascular access audits, CHG treatment compliance, or C. difficile testing. Differences in gender and age were found in CHG treatment compliance ($p < 0.001$). **Conclusions:** By focusing more on patient level process measures rather than only presenting stratified outcomes data, we can identify targeted