

significantly higher percentage than the one of females (81.47%), (p-value=0.0004).

98.85% of males receive antipsychotic treatments versus 99.56% of females (p-value=0.1185).

35.22% of male patients receive non-drug treatment versus 33.24% of females (p-value=0.3966).

Most often prescribed antipsychotics in this sample are: Risperidone: 28.90%(Males 29.43%, females 28.11%), Haloperidole: 22.96%(23.62%,21.96% respectively), Olanzapine: 21.39%(20.72%, 22.40%), Aripiprazole: 17.60% (15.68%,20.50%) Quetiapine: 9.50%: (9.00%, 10.25%), Amisulpride: 8.51%(9.78%, 6.59%), Ziprasidone: 8.33% (9.00%,7.32%), Clozapin:7.28% (8.52%, 5.42%), Zuclopentixol: 1.98% (2.52%,1.17%).

The average daily dosage for each of the above-mentioned medications for the whole sample and by gender has also been assessed whether they are used as monotherapy or as part of polytherapy.

Conclusions: In Greece, more male Schizophrenic patients compared to females are receiving care for their illness. Second generation antipsychotic treatments represent the treatment of choice.

P144

Most frequently observed adverse events among schizophrenic outpatients in Greece (The Greek 'ACE' study)

E.C. Tsiolka¹, A.A. Karavatos², V.G. Mavreas³, I.A. Chatzimanolis⁴. ¹Medical Department, Bristol-Myers Squibb EPE, Athens, Greece ²First Psychiatric Clinic, General Hospital "Papageorgiou", Thessalonica, Greece ³Psychiatric Clinic, General University Hospital of Ioannina, Ioannina, Greece ⁴First Psychiatric Clinic, University Hospital "Eginitio", Athens, Greece

Purpose: To evaluate the current medical status, clinical care and treatment patterns of schizophrenic patients in Greece.

Methods: The Greek 'ACE' study is a descriptive, cross-sectional, multicentre national survey conducted in outpatient setting by 101 psychiatrists (under 3 Coordinators Psychiatrists) proportionately distributed all over Greece who registered during one visit the first twenty consecutive patients presenting at their consultation. Observational period :29/09/2005 -22/12/2005.

Recruited 2040 patients over 18 years old, with an ICD-10 primary diagnosis of schizophrenia.

Results: A total of 1716 patients (1033 males and 683 females) were treated with antipsychotics agents. Among them, 52.10% experienced adverse events at the time of data collection

The most frequently reported adverse events is weight gain(45.86%), extrapyramidal symptoms (32.88%), sedation and impaired concentration (24.50%), dry mouth (22.26%), sexual(18.23%) and menstrual (9.50%) problems. Distribution of various adverse events separately for males and females by each medication when received as monotherapy, is also presented.

The percentage of males presenting with high cholesterol values (13.44%) is statistically significantly higher than of females (9.46%) (p-value=0.0124), while this does not happen neither with high triglycerides values (9.21%-males vs 6.99%-females) (p-value=0.1011),nor

with high glucose values (6.72%-males vs 6.71%-females) (p-value=0.9920).

Distribution of high levels of cholesterol triglycerides and glucose, separately for males and females by each medication when received as monotherapy, is also presented.

Conclusions: Weight gain is the most frequently reported adverse event in this study. It is important to patients to receive the most appropriate treatment, taking in consideration both symptom control as well as potential side-effects.

P145

Contribution of extramural psychiatric care units in the treatment of psychosis in rural population

P. Panagoutsos, I. Syros, D. Tzavaras, A. Tourka. *Psychiatric Hospital of Tripolis, Tripolis, Arcadia, Greece*

Background and aims: Evidence shows that the hospital-based character of psychiatric services has a negative impact on the personal and social life of the psychotic patients. This situation is worse in rural areas where the Psychiatric Care facilities are fewer. In a 5-year study we investigate the therapeutic role of the Ambulatory Mobile Care Unit in the provinces of Fokida and Evros.

Methods: 45 patients, aged between 17-67, diagnosed with psychosis (schizophrenia, schizoaffective disorder, delusional states, bipolar disorder etc.) were studied. Most of them were unemployed (56%), living with their parents, and belonged to the lower or middle class. An experienced psychiatrist-supervisor, other psychiatrists, trainees, psychologists, social workers and nurses were part of the intervention team that helped the patients in different stages of the disease (acute state, stable condition).

Results: The intervention of the Mobile Care Ambulatory Unit resulted in:

- Increase in the Employment Rate of the Patients (more than half of the patients has been employed in the end of the intervention)
- Improvement in social abilities (more than 4/5 of the patients had improved relations with the local community)
- Leisure time activities: acquisition of new hobbies in more than 2/3 of the patients
- Insight: most of the patients came in terms with their disease
- Nevertheless, very few of them managed to form their own family.

Conclusion: The mobile unit intervention has had a significant effect in most aspects of the life of the psychotic patients.

P146

Extramural psychiatric care units in the treatment of psychosis in rural population: Investigation of the number and duration of relapses in a 5-year study

P. Panagoutsos, D. Tzavaras, A. Tourka, I. Syros. *Psychiatric Hospital of Tripolis, Arcadia, Greece*

Background and aims: The limited number of extramural residential facilities in the rural areas of Greece leads to the diminished effectiveness of the follow-up of the psychotic patients and the relapses of their condition. The goal of our study was to investigate the factors involved in the number and duration of the relapses, as well as the role of the intervention of the Ambulatory Mobile Units in modifying these parameters.

Methods: By using different statistical techniques 45 patients, aged between 17-67, were studied. In a 5-year follow-up, parameters like the number of relapses and their duration were correlated with socio-demographic variables and parameters concerning the personal and family background of the patient. A standard team consisting of an experienced psychiatrist-supervisor, other psychiatrists, trainees, psychologists, social workers and nurses were collaborating with the patients.