

Brownian motion on the golden ratio Sierpinski gasket

Shiping Cao

Department of Mathematics, Cornell University, Ithaca 14853, USA
sc2873@cornell.edu

Hua Qiu

Department of Mathematics, Nanjing University, Nanjing 210093, China
huaqiu@nju.edu.cn

(Received 7 July 2022; accepted 28 February 2023)

We construct a strongly local regular Dirichlet form on the golden ratio Sierpinski gasket, which is a self-similar set without a finitely ramified cell structure, via a study on the trace of an electrical network on an infinite graph. The Dirichlet form is the unique one that is self-similar in the sense of an infinite iterated function system, and is decimation invariant with respect to a graph-directed construction. The proof is based on a fixed point problem of a renormalization map, inspired by Sabot's celebrated work for finitely ramified fractals. Lastly, the Hunt process associated with the Dirichlet form satisfies a two-sided sub-Gaussian heat kernel estimate.

Keywords: golden ratio Sierpinski gasket; infinite graph; Dirichlet forms; heat kernel estimates

2010 *Mathematics Subject Classification:* Primary 28A80; 31E05

1. Introduction

The golden ratio Sierpinski gasket \mathcal{G} is a typical example of a self-similar set satisfying the finite type property ([2], see definition 2.1.), which arises in the study of the Hausdorff dimension of self-similar sets with overlaps [26, 31, 32]. Let $q_0 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$, $q_1 = (0, 0)$, $q_2 = (1, 0)$ be the three vertices of an equilateral triangle in \mathbb{R}^2 , and

$$F_0(x) = \rho^2(x - q_0) + q_0, \\ F_1(x) = \rho(x - q_1) + q_1, \quad F_2(x) = \rho(x - q_2) + q_2,$$

with $\rho = \frac{\sqrt{5}-1}{2}$ being the golden ratio. The gasket \mathcal{G} is the invariant set associated with the iterated function system (IFS for short) $\{F_0, F_1, F_2\}$, i.e. \mathcal{G} is the unique

© The Author(s), 2023. Published by Cambridge University Press on behalf of The Royal Society of Edinburgh




Figure 1. The golden ratio Sierpinski gasket \mathcal{G} .

non-empty compact set satisfying

$$\mathcal{G} = \bigcup_{i=0}^2 F_i \mathcal{G}.$$

See [figure 1](#).

The large overlap $F_1 \mathcal{G} \cap F_2 \mathcal{G}$ makes \mathcal{G} different from the existing examples of self-similar sets on which Brownian motions are constructed.

First, any effort to disconnect the bottom line of \mathcal{G} requires the removal of infinitely many points, so there is not a finitely ramified cell structure [35] on \mathcal{G} . Well-known classes of fractals with finitely ramified cell structures include Lindström's nested fractals [27], Kigami's post-critically finite (PCF) self-similar sets [20, 21], finitely ramified graph-directed fractals [9, 19], and some Julia sets of polynomials [1, 14, 33] or rational functions [10]. See [8, 16, 24] for pioneering works on the Sierpinski gasket, and also books [3, 22] for systematic discussions.

Second, although there is a graph-directed construction related with \mathcal{G} (see §2), by dividing \mathcal{G} into blocks of nearly the same size, the graph is much more complicated. As a result, the deep and famous constructions on the Sierpinski carpet [4–6] by Barlow and Bass, and on certain symmetric fractals [25] by Kusuoka and Zhou would be extremely difficult here. See also [7] for a theorem of uniqueness on the Sierpinski carpet.

Instead, thanks to the golden ratio, there is an ‘infinite cell structure’ on \mathcal{G} . For the first level, we consider the cell $F_0 \mathcal{G}$ and its images under compositions of F_1, F_2 . The union of these cells covers \mathcal{G} except the bottom line. For each such cell, we can find a finite word w , and a contraction map $F_w = F_{w_1} \circ F_{w_2} \circ \cdots \circ F_{w_m}$, so that the cell can be written as $F_w \mathcal{G}$. We name the collection of all such words W_1 , and construct a resistance form [22] on \mathcal{G} , that is self-similar in the sense of the infinite IFS $\{F_w\}_{w \in W_1}$. Roughly speaking, we have the following theorem, see theorems 6.5, 6.6 and 6.8 for detailed and formal results.

THEOREM 1.1. *There exists a unique strongly local regular resistance form $(\mathcal{E}, \mathcal{F})$ on \mathcal{G} such that $f \in \mathcal{F}$ if and only if $f \circ F_w \in \mathcal{F}$ for all $w \in W_1$ and $\sum_{w \in W_1} \rho_w^{-\theta} \mathcal{E}(f \circ F_w, f \circ F_w) < \infty$, where ρ_w is the similarity ratio of F_w and $0 < \theta < 1$ is a constant.*

In addition,

$$\mathcal{E}(f, f) = \sum_{w \in W_1} \rho_w^{-\theta} \mathcal{E}(f \circ F_w, f \circ F_w).$$

Moreover, the form is decimation invariant with respect to the graph-directed construction of \mathcal{G} .

The form $(\mathcal{E}, \mathcal{F})$ is then a strongly local regular Dirichlet form on $L^2(\mathcal{G}, \mu_H)$, where μ_H is the normalized Hausdorff measure on \mathcal{G} . In addition, there is an associated diffusion process on \mathcal{G} . (Readers are suggested to refer the book [15] for more explanations.) Although, our construction is based on an infinite IFS, the behaviour of the process is same on each cell before hitting the boundary, up to a time scaling, since any cell can be decomposed in a same manner.

In addition, by following the well-established method of Hambly and Kumagai [18], which is organized in Barlow's book [3], we can obtain a sub-Gaussian heat kernel estimate (see §7). We refer to [8, 13, 23] for earlier results on transition density estimates on fractals.

THEOREM 1.2. *There is a symmetric transition density $p(t, x, y)$ associated with the form $(\mathcal{E}, \mathcal{F})$ on \mathcal{G} . In addition, there are constants c_1, c_2, c_3, c_4 so that*

$$\begin{aligned} c_1 t^{-d_H/\beta} \exp\left(-c_2 \left(\frac{d(x, y)^\beta}{t}\right)^{\frac{1}{\beta-1}}\right) \\ \leq p(t, x, y) \leq c_3 t^{-d_H/\beta} \exp\left(-c_4 \left(\frac{d(x, y)^\beta}{t}\right)^{\frac{1}{\beta-1}}\right), \end{aligned}$$

for $0 < t \leq 1$, with $\beta = \theta + d_H$, where $d_H \approx 1.6824$ is the Hausdorff dimension of \mathcal{G} , and d represents the Euclidean metric.

The main difficulty in proving this result is establishing an estimate for the resistance metric R on \mathcal{G} of the form $c_1 d(x, y)^\theta \leq R(x, y) \leq c_2 d(x, y)^\theta$ for some constant $c_1, c_2 > 0$.

We organize the structure of the paper as follows. In §2, we will briefly introduce some facts about the geometry of \mathcal{G} . From §3 to 5, we study the trace of forms on an infinite graph. In §3, we establish the resistance forms on the graph. In §4, we study the trace map and a related renormalization map. We will show the joint continuity of the renormalization map. In §5, we show that there is a unique solution to a renormalization problem. With all these preparations, we construct the resistance form on \mathcal{G} in §6, and at the same time we derive an upper-bound estimate for the resistance metric. Lastly, we obtain the transition density estimate through a lower bound for the resistance metric in §7.

Before ending this section, we remark that the result in this paper has a natural extension, by replacing $0 < \rho < 1$ to be a real root of $x^n - 2x + 1$ with $n \geq 4$, and taking the contraction ratios corresponding to F_0, F_1, F_2 to be $1 - \rho, \rho, \rho$. Indeed, by doing so, we obtain a class of gaskets that possess a similar overlapping structure of \mathcal{G} , see figure 2.




Figure 2. A gasket with $0 < \rho < 1$ being a root of $x^4 - 2x + 1 = 0$.

2. Preliminary

The golden ratio Sierpinski gasket \mathcal{G} is a typical example of a self-similar set with overlaps but satisfying the finite type property.

Let K be a general self-similar set associated with an IFS $\{F_i\}_{i=0}^{N-1}$ with contraction ratios $\{\rho_i\}_{i=0}^{N-1}$ with respect to the Euclidean metric. For $m \geq 1$, we call $w = w_1w_2 \cdots w_m$ with $w_i \in \{0, 1, \dots, N - 1\}$, a *word* of length m (denoted by $|w|$), and call \emptyset the empty word. We denote the set of all words by \tilde{W}_* . For any word $w \in \tilde{W}_*$, we write $F_w = F_{w_1} \circ F_{w_2} \cdots \circ F_{w_{|w|}}$, and let F_\emptyset be the identity map for consistency. Let $\rho_* = \min\{\rho_i : 0 \leq i < N\}$.

DEFINITION 2.1 *finite type property.* A self-similar set K is of finite type if there are only finitely many maps $h = F_w^{-1}F_v$ with $w, v \in \tilde{W}_*$ and $F_wK \cap F_vK \neq \emptyset$, and with similarity ratio $\rho_h \in (\rho_*, 1/\rho_*)$.

The finite type property of K , formulated in algebraic terms, was introduced in [2] by Bandt and Rao. It guarantees the existence of an ‘almost non-overlapping’ graph-directed construction (see [2, 31] for details) of K , which is quite useful for calculating the Hausdorff dimension of K . See [26, 32] for more flexible variants of the finite type property.

It is easy to verify that \mathcal{G} satisfies the finite type property, noticing that $F_{122}\mathcal{G} = F_{211}\mathcal{G}$. In particular, it has the following *graph-directed construction* [28].

DEFINITION 2.2 a graph-directed construction of \mathcal{G} .

- (a). Let $K_1 = \mathcal{G}$ and $K_2 = \overline{\mathcal{G} \setminus F_{22}\mathcal{G}}$.
- (b). Let $\Gamma(S, E)$ be a directed graph with the vertex set $S = \{1, 2\}$, and the edge set $E = \{e_i\}_{i=1}^6$, where $e_1 = (1, 2), e_2 = (1, 1), e_3 = (2, 1), e_4 = (2, 2), e_5 = (2, 2), e_6 = (2, 1)$.
- (c). Define $\psi_{e_1} = Id, \psi_{e_2} = F_{22}, \psi_{e_3} = F_0, \psi_{e_4} = F_1, \psi_{e_5} = F_{21}, \psi_{e_6} = F_{20}$.




Figure 3. A graph-directed construction of \mathcal{G} .

Clearly, we have

$$K_1 = \bigcup_{i=1}^2 \psi_{e_i} K_{e_{i,2}} \text{ and } K_2 = \bigcup_{i=3}^6 \psi_{e_i} K_{e_{i,2}},$$

where we use the notation $e_i = (e_{i,1}, e_{i,2})$ for a directed edge. In addition, there exist bounded open sets O_1 and O_2 such that $\bigcup_{i=1}^2 \psi_{e_i} O_{e_{i,2}} \subset O_1$ and $\bigcup_{i=3}^6 \psi_{e_i} O_{e_{i,2}} \subset O_2$, where the unions are disjoint. See figure 3 for an illustration.

Then similar to the open set condition situation, one can calculate the exact value of the Hausdorff dimension of \mathcal{G} to be

$$d_H = \frac{\log \eta}{-2 \log \rho} \approx 1.6824$$

with η being the largest root of $x^3 - 6x^2 + 5x - 1$. In addition, the associated Hausdorff measure of \mathcal{G} is positive and finite. See details in [31] by Ngai and Wang.

Throughout this paper, we use d to denote the Euclidean metric, and take μ_H to be the normalized Hausdorff measure on \mathcal{G} with respect to d , i.e. $\mu_H(\mathcal{G}) = 1$. For $p, q \in \mathcal{G}$, let

$$d_g(p, q) = \inf\{|\gamma| : \gamma \text{ is a path connecting } p, q, \text{ and } \gamma \subset \mathcal{G}\},$$

be the *geodesic metric* between p, q . It is not hard to verify the following lemma.

LEMMA 2.3.

(a). Let $B_s(p) = \{q \in \mathcal{G} : d(p, q) < s\}$. There are constants $c_1, c_2 > 0$ such that

$$c_1 s^{d_H} \leq \mu_H(B_s(p)) \leq c_2 s^{d_H}, \quad \forall p \in \mathcal{G}, 0 < s \leq 1.$$

(b). There exists a constant $c \geq 1$ such that

$$d(p, q) \leq d_g(p, q) \leq cd(p, q), \quad \forall p, q \in \mathcal{G}.$$

The statement (a) is a well-known fact (for example, see [12, corollary 6.4.4]). The proof of (b) relies on the finite type property. The rough idea is to link p, q with a bounded number of cells of diameter approximating to $d(p, q)$.




Figure 4. The infinite graph (V_1, \sim) . (The bottom line equals to $\bar{V}_1 \setminus V_1$.)

By the compactness of \mathcal{G} (for example, see [3, lemma 2.1.1]), there is always a path admitting the infimum length between p, q . So, the metric space (\mathcal{G}, d_g) satisfies the so-called *midpoint property*, i.e. for any $p, q \in \mathcal{G}$, there exists p' so that $d_g(p, p') = d_g(p', q) = \frac{1}{2}d_g(p, q)$. The space $(\mathcal{G}, d_g, \mu_H)$ is then a *fractional metric space*, see [3, definition 3.2].

We will return to look at the geometric properties of \mathcal{G} listed in this section. But first, from § 3 to 5, we will instead consider an infinite IFS and the associated infinite graph.

3. Resistance forms on the infinite graph V_1

The golden ratio Sierpinski gasket \mathcal{G} can be realized as an invariant set of an infinite IFS. For convenience, we introduce some notation. For any word $w, w' \in \tilde{W}_*$, we write ww' for the concatenation of w, w' . For $w = w_1w_2 \cdots w_m$ and $0 \leq l \leq m$, we write $[w]_l = w_1w_2 \cdots w_l$. The following notation is a little different from the standard ones.

Notation. Choose a set of finite words $W_1 \subset \bigcup_{n=0}^\infty \{1, 2\}^n \times \{0\}$ so that

1. for any $w \in \bigcup_{n=0}^\infty \{1, 2\}^n \times \{0\}$, there exists $w' \in W_1$ such that $F_w = F_{w'}$;
2. for different words $w, w' \in W_1$, we have $F_w \neq F_{w'}$.

In addition, based on W_1 , we introduce some more notations.

- (a) For $n \geq 1$, define $W_{1,n} = \{w \in W_1 : |w| = n\}$;
- (b) For $m \geq 2$, define $W_m := W_1^m = \{w_1w_2 \cdots w_m : w_i \in W_1, 1 \leq i \leq m\}$;
- (c) Write $V_0 = \{q_i\}_{i=0}^2$ and for $m \geq 1$, $V_m = \bigcup_{w \in W_m} F_w V_0$. Denote \bar{V}_m the closure of V_m ;
- (d) For distinct $p, q \in V_1$, we denote $p \sim q$ if and only if $p, q \in F_w V_0$ for some $w \in W_1$, which induce an infinite graph (V_1, \sim) . See figure 4 for an illustration.

Obviously, we have

$$\mathcal{G} = \overline{\bigcup_{w \in W_1} F_w \mathcal{G}},$$

and thus $\{F_w\}_{w \in W_1}$ is an infinite i.f.s associated with \mathcal{G} . See [30] for more details about infinite IFSs. The advantage of this IFS lies in the fact that

$$F_w \mathcal{G} \cap F_{w'} \mathcal{G} = F_w V_0 \cap F_{w'} V_0, \quad \forall w \neq w' \in W_1.$$

REMARK. For $n \geq 1$, if we rename the vertices $\{F_w q_0\}_{w \in W_{1,n}}$ to be $\{p_i^{(n)}\}_{i=1}^{N_n}$ with $N_n := \#W_{1,n}$, so that for each i , $p_i^{(n)}$ is on the left of $p_{i+1}^{(n)}$. Then it directly calculates that $d(p_i^{(n)}, p_{i+1}^{(n)})$ is either ρ^n or ρ^{n+1} , and thus $N_n \asymp \rho^{-n}$.

In addition, for $p, q \in V_1$ with $d(p, q) < \rho^{n+2}$, there always exist $w, w' \in \{1, 2\}^n$ such that $p \in F_w V_1, q \in F_{w'} V_1$ and $F_w V_1 \cap F_{w'} V_1 \neq \emptyset$. In fact, by a direct observation, $p \notin \bigcup_{n'=1}^{n+1} \{p_i^{(n')}\}_{i=1}^{N_{n'}}$, and so we can find $\tilde{w} \in \{1, 2\}^{n+1}$ such that $p \in F_{\tilde{w}} V_1$, and one can then see that $q \in \bigcup \{F_{w'} V_1 : w, w' \in \{1, 2\}^n, F_{\tilde{w}} V_1 \subset F_w V_1, F_{w'} V_1 \cap F_w V_1 \neq \emptyset\}$, since otherwise $d(p, q) > \frac{\sqrt{3}}{2} \cdot \rho^{n+1} \geq \rho^{n+2}$.

In the rest of this section, we consider a class of resistance forms generated by decimation. For convenience of readers, we recall the general definition of resistance forms in the following. See [22] for more details.

DEFINITION 3.1. Let X be a set, and $l(X)$ be the space of all real-valued functions on X . A pair $(\mathcal{E}, \mathcal{F})$ is called a (non-degenerate) resistance form on X if it satisfies the following conditions:

- (RF1) \mathcal{F} is a linear subspace of $l(X)$ containing constants and \mathcal{E} is a nonnegative symmetric quadratic form on \mathcal{F} ; $\mathcal{E}(f) := \mathcal{E}(f, f) = 0$ if and only if f is constant on X .
- (RF2) Let \sim be an equivalence relation on \mathcal{F} defined by $f \sim g$ if and only if $f - g$ is constant on X . Then $(\mathcal{F}/\sim, \mathcal{E})$ is a Hilbert space.
- (RF3) For any finite subset $V \subset X$ and any $u \in l(V)$, there exists a function $f \in \mathcal{F}$ such that $f|_V = u$.
- (RF4) For any distinct $p, q \in X$, $R(p, q) := \sup\{\frac{|f(p)-f(q)|^2}{\mathcal{E}(f)} : f \in \mathcal{F}, \mathcal{E}(f) > 0\}$ is finite.
- (RF5) If $f \in \mathcal{F}$, then $\bar{f} = \min\{\max\{f, 0\}, 1\} \in \mathcal{F}$ and $\mathcal{E}(\bar{f}) \leq \mathcal{E}(f)$.

Sometimes, we write $\mathcal{F} = \text{Dom}(\mathcal{E})$, and abbreviate $(\mathcal{E}, \mathcal{F})$ to \mathcal{E} when no confusion occurs. It is well-known ([22]) that $R(p, q)$ defined in (RF3) is a metric on X , named the effective resistance metric.

On the finite set V_0 , a resistance form \mathcal{D} always has the form

$$\mathcal{D}(f, g) = \frac{1}{2} \sum_{i,j} a_{i,j} (f(q_i) - f(q_j)) (g(q_i) - g(q_j)), \quad \forall f, g \in l(V_0), \tag{3.1}$$

where $a_{i,i} = 0$ and the 3×3 matrix $(a_{i,j})$ is positive, symmetric and irreducible. For convenience, we write \mathcal{M} for the collection of all resistance forms on V_0 . We view \mathcal{M} as a subset of \mathbb{R}^3 , which is not closed with the induced topology.

Given a resistance form \mathcal{D} , we define a resistance form on V_1 associated with \mathcal{D} in a self-similar manner, respecting the infinite IFS $\{F_w\}_{w \in W_1}$.

DEFINITION 3.2. For $r > 0$, $\mathcal{D} \in \mathcal{M}$, we define $\Psi_r \mathcal{D}$ as

$$\Psi_r \mathcal{D}(f, g) = \sum_{w \in W_1} r^{-|w|+1} \mathcal{D}(f \circ F_w, g \circ F_w),$$

with $Dom(\Psi_r \mathcal{D}) = \{f \in l(V_1) : \Psi_r \mathcal{D}(f) < \infty\}$.

It is not hard to show that $(\Psi_r \mathcal{D}, Dom(\Psi_r \mathcal{D}))$ is a resistance form on V_1 . However, to get a good resistance form, we need to restrict the range of r .

PROPOSITION 3.3. Let $\mathcal{D} \in \mathcal{M}$ and $r < 1$, then $Dom(\Psi_r \mathcal{D}) \subset C(\bar{V}_1)$ by a natural identification. In addition, if $\rho < r < 1$, then $(\Psi_r \mathcal{D}, Dom(\Psi_r \mathcal{D}))$ extends to a resistance form on \bar{V}_1 , with the associated resistance metric $R(p, q)$ satisfying the estimate

$$R(p, q) \leq \frac{4}{r^3(1-r)} R_0^*(\mathcal{D}) d(p, q)^{\frac{\log r}{\log \rho}}, \quad \forall p, q \in \bar{V}_1, \tag{3.2}$$

where $R_0^*(\mathcal{D}) = \max_{p,q \in V_0} R_0(p, q)$ with R_0 being the resistance metric on V_0 associated with \mathcal{D} .

Proof. By scaling, $R(p, q) \leq R_0^*(\mathcal{D})r^{n-1}$ for any distinct $p, q \in F_w V_0$ with $w \in W_{1,n}$ and $n \geq 1$. For $w \in \{1, 2\}^n$, write $[w]_l = w_1 w_2 \cdots w_l$ and $p_l = F_{[w]_l}(q_0)$, with $0 \leq l \leq n$, then

$$R(p_i, p_j) \leq \sum_{l=i}^{j-1} R(p_l, p_{l+1}) \leq R_0^*(\mathcal{D}) \sum_{l=i}^{j-1} r^l < R_0^*(\mathcal{D}) \frac{r^i}{1-r}, \quad \forall 0 \leq i < j \leq n.$$

In particular, this implies that $R(p, q) \leq \frac{2r^n}{1-r} R_0^*(\mathcal{D})$ for any $p, q \in F_w V_1$ and $w \in \{1, 2\}^n$. Now, if $p, q \in V_1$ and $d(p, q) < \rho^{n+2}$, then by the remark before definition 3.1, there exist $w, w' \in \{1, 2\}^n$ such that $p \in F_w V_1, q \in F_{w'} V_1$, and $F_w V_1 \cap F_{w'} V_1 \neq \emptyset$, which implies that $R(p, q) \leq \frac{4r^n}{1-r} R_0^*(\mathcal{D})$. As a consequence, we have

$$R(p, q) \leq \frac{4}{r^3(1-r)} R_0^*(\mathcal{D}) d(p, q)^{\frac{\log r}{\log \rho}}, \quad \forall p, q \in V_1. \tag{3.3}$$

On the other hand, for any $f \in Dom(\Psi_r \mathcal{D})$, by (RF4), we immediately have

$$|f(p) - f(q)| \leq (R(p, q) \Psi_r \mathcal{D}(f))^{1/2}. \tag{3.4}$$

Combining (3.3) and (3.4), we then get $Dom(\Psi_r \mathcal{D}) \subset C(\bar{V}_1)$ by a natural identification.

To show the second assertion, we let (X, R) be the completion of (V_1, R) , and recall [22, theorem 2.3.10] to get that $(\Psi_\lambda \mathcal{D}, \text{Dom}(\Psi_r \mathcal{D}))$ extends to be a resistance form on X . It suffices to show that the identity map $Id : V_1 \rightarrow V_1$ extends to an homeomorphism from (\bar{V}_1, d) to (X, R) under the assumption $\rho < r < 1$. First, by (3.3), Id is continuous from (V_1, d) to (V_1, R) . Next, let $f \in l(V_1)$ be a restriction of a linear function on \mathbb{R}^2 . We have

$$\begin{aligned} \Psi_r \mathcal{D}(f) &= \sum_{w \in W_1} r^{-|w|+1} \mathcal{D}(f \circ F_w) = \sum_{n=1}^{\infty} \sum_{w \in W_{1,n}} r^{-n+1} \mathcal{D}(f \circ F_w) \\ &= \sum_{n=1}^{\infty} \#W_{1,n} r^{-n+1} \rho^{2(n+1)} \mathcal{D}(f|_{V_0}), \end{aligned}$$

where the last equality follows from the fact that f is linear. Since $\#W_{1,n} \asymp \rho^{-n}$, we have $\Psi_r \mathcal{D}(f) < \infty$ when $r > \rho$, so that $f \in \text{Dom}(\Psi_r \mathcal{D})$. Noticing that for any points $p \neq q \in \bar{V}_1$, we can find a linear function f such that $f(p) \neq f(q)$, we have Id is injective. Finally, due to the fact that (\bar{V}_1, d) is a compact Hausdorff space and (X, R) is the completion of (V_1, R) , we then have that Id is an homeomorphism from (\bar{V}_1, d) to (X, R) . This implies that $(\Psi_r \mathcal{D}, \text{Dom}(\Psi_r \mathcal{D}))$ is a resistance form on \bar{V}_1 , and (3.2) follows immediately from (3.3). \square

REMARK. *The restriction $\rho < r < 1$ is sharp. If $r \leq \rho$, there is no $f \in \text{Dom}(\Psi_r \mathcal{D})$ such that $f(q_1) = 0$ and $f(q_2) = 1$. In fact, for any $f \in C(\bar{V}_1)$ with $f(q_1) = 0$ and $f(q_2) = 1$, by the remark before definition 3.1, the total energy of f on the union of the cells $F_w V_0, w \in W_{1,n} \cup W_{1,n+1}$ (noticing that this union will induce a connected subgraph in (V_1, \sim) , and the resistance between $F_1^n q_0$ and $F_2^n q_0$ is about $r^n \rho^{-n}$) will be bounded away from 0 as $n \rightarrow \infty$.*

4. A renormalization map

In proposition 3.3, we have shown that $\Psi_r \mathcal{D}$ extends to be a resistance form on \bar{V}_1 when $\rho < r < 1$. It is natural to trace it back to V_0 , noticing that $V_0 \subset \bar{V}_1$.

DEFINITION 4.1. *Let $(\mathcal{D}_1, \mathcal{F}_1)$ be a resistance form on \bar{V}_1 , we write*

$$[\mathcal{D}_1]_{V_0}(u) = \inf\{\mathcal{D}_1(f) : f|_{V_0} = u, f \in \mathcal{F}_1\}, \quad \forall u \in l(V_0).$$

Note that by a standard electric network theory, there exists a unique function f so that $\mathcal{D}_1(f)$ attains the infimum above; also $[\mathcal{D}_1]_{V_0}$ induces a resistance form on V_0 by defining

$$[\mathcal{D}_1]_{V_0}(u, v) := \frac{1}{4} ([\mathcal{D}_1]_{V_0}(u + v) - [\mathcal{D}_1]_{V_0}(u - v)).$$

For $\rho < r < 1$ and $\mathcal{D} \in \mathcal{M}$, we define $\mathcal{R}_r \mathcal{D} = [\Psi_r \mathcal{D}]_{V_0}$, and call \mathcal{R}_r the renormalization map. Sometimes, we also write $\mathcal{R}(r, \mathcal{D}) := \mathcal{R}_r(\mathcal{D})$.

The main purpose of this section is to show the continuity of the map $\mathcal{R}(r, \mathcal{D})$.

THEOREM 4.2. *The map $\mathcal{R}(r, \mathcal{D})$ is jointly continuous from $(\rho, 1) \times \mathcal{M}$ to \mathcal{M} .*

To prove theorem 4.2, we need a study on the regularity of the resistance form $\Psi_r \mathcal{D}$.

PROPOSITION 4.3. *Let $\mathcal{D} \in \mathcal{M}$ and $\rho < r_1 < r_2 < 1$. Then*

- (a) *Dom($\Psi_{r_1} \mathcal{D}$) depends only on r_1 , and we have $Dom(\Psi_{r_1} \mathcal{D}) \subset Dom(\Psi_{r_2} \mathcal{D})$.*
- (b) *Dom($\Psi_{r_1} \mathcal{D}$) is dense in $Dom(\Psi_{r_2} \mathcal{D})$ in the sense that for any $f \in Dom(\Psi_{r_2} \mathcal{D})$ and $\varepsilon > 0$, there exists $g \in Dom(\Psi_{r_1} \mathcal{D})$ such that*

$$\Psi_{r_2} \mathcal{D}(f - g) < \varepsilon, \text{ and } f|_{V_0} = g|_{V_0}.$$

Moreover, $Dom(\Psi_{r_1} \mathcal{D})$ is dense in $C(V_1)$ so that the resistance form is regular.

Proof. (a) is obvious since all $\mathcal{D} \in \mathcal{M}$ are comparable up to multiplicative constants, we only need to prove (b). Let $f \in Dom(\Psi_{r_2} \mathcal{D})$, and choose n large enough so that

$$\sum_{l=n}^{\infty} \sum_{w \in W_{1,l}} r_2^{-l+1} \mathcal{D}(f \circ F_w) < \varepsilon. \tag{4.1}$$

For convenience, we rename the vertices $\{F_w q_0\}_{w \in W_{1,n}}$ to be $\{p_i\}_{i=1}^N$ with $N = \#W_{1,n}$, so that for each i , p_i is on the left of p_{i+1} . Then, noticing that the effective resistance between q_1 and p_1 (symmetrically, q_2 and p_N) is bounded above by a multiple of r^{-n} , by (RF4), it is not hard to see

$$\begin{aligned} & r_2^{-n} \left(\sum_{i=1}^{N-1} (f(p_i) - f(p_{i+1}))^2 + (f(q_1) - f(p_1))^2 + (f(q_2) - f(p_N))^2 \right) \\ & \leq c_1 \left(\sum_{l=n}^{\infty} \sum_{w \in W_{1,l}} r_2^{-l+1} \mathcal{D}(f \circ F_w) \right) < c_1 \varepsilon, \end{aligned}$$

where c_1 is a constant depending on \mathcal{D} and r_2 , but not on n .

Write x_i for the x -coordinate of p_i , so we have $0 < x_1 < x_2 < \dots < x_N < 1$. We introduce a piecewise linear function u on \mathbb{R}^2 such that

1. $u(x, y)$ depends only on x ;
2. $u(q_1) = f(q_1)$, $u(q_2) = f(q_2)$, and $u(p_i) = f(p_i)$, $1 \leq i \leq N$;
3. $u(x, 0)$ is linear on each interval $(0, x_1)$, $(x_N, 1)$ and (x_i, x_{i+1}) , $1 \leq i \leq N - 1$.

We define $g \in l(V_1)$ as

$$g(p) = \begin{cases} f(p), & \text{if } p \in \bigcup_{i=1}^{n-1} \bigcup_{w \in W_{1,i}} \{F_w q_0\}, \\ u(p), & \text{if } p \in \bigcup_{i=n}^{\infty} \bigcup_{w \in W_{1,i}} \{F_w q_0\}. \end{cases}$$

By a similar estimate to that applied in the proof of proposition 3.3, one can check that $g \in \text{Dom}(\Psi_{r_1}\mathcal{D})$, and

$$\sum_{l=n}^{\infty} \sum_{w \in W_{1,l}} r_2^{-l+1} \mathcal{D}(g \circ F_w) \leq c_2 r_2^{-n} \left(\sum_{i=1}^{N-1} (f(p_i) - f(p_{i+1}))^2 + (f(q_1) - f(p_1))^2 + (f(q_2) - f(p_N))^2 \right),$$

where c_2 depends only on \mathcal{D} and r_2 . So we have $\Psi_{r_2}\mathcal{D}(f - g) \leq c_3\varepsilon$ for some constant c_3 . Since ε can be arbitrarily small, we have that $\text{Dom}(\Psi_{r_1}\mathcal{D})$ is dense in $\text{Dom}(\Psi_{r_2}\mathcal{D})$. Finally, the claim that $\text{Dom}(\Psi_{r_1}\mathcal{D})$ is dense in $C(\bar{V}_1)$ follows from the same argument. \square

Proof of theorem 4.2. Let $r_n \rightarrow r \in (\rho, 1)$ and $\mathcal{D}_n \rightarrow \mathcal{D} \in \mathcal{M}$. Also, let $u \in l(V_0)$. First, we show that

$$\limsup_{n \rightarrow \infty} \mathcal{R}(r_n, \mathcal{D}_n)(u) \leq \mathcal{R}(r, \mathcal{D})(u). \tag{4.2}$$

We define f to be the unique function in $\text{Dom}(\Psi_r\mathcal{D})$ such that $f|_{V_0} = u$ and

$$\mathcal{R}(r, \mathcal{D})(u) = \Psi_r\mathcal{D}(f).$$

By proposition 4.3, for any $\varepsilon > 0$, there is f_ε such that $f_\varepsilon|_{V_0} = u$, $f_\varepsilon \in \text{Dom}(\Psi_{r_n}\mathcal{D}_n)$ for any $n \geq 1$, and

$$\Psi_r\mathcal{D}(f_\varepsilon) \leq \Psi_r\mathcal{D}(f) + \varepsilon.$$

As a consequence, we have

$$\limsup_{n \rightarrow \infty} \mathcal{R}(r_n, \mathcal{D}_n)(u) \leq \lim_{n \rightarrow \infty} \Psi_{r_n}\mathcal{D}_n(f_\varepsilon) = \Psi_r\mathcal{D}(f_\varepsilon) \leq \mathcal{R}(r, \mathcal{D})(u) + \varepsilon,$$

where the equality is due to the dominated convergence theorem. Since ε can be arbitrarily chosen, we get (4.2).

Next, for each n , let f_n be the unique function in $\text{Dom}(\Psi_{r_n}\mathcal{D}_n)$ such that $f_n|_{V_0} = u$ and

$$\mathcal{R}(r_n, \mathcal{D}_n)(u) = \Psi_{r_n}\mathcal{D}_n(f_n).$$

Then $\{f_n\}_{n \geq 1}$ is uniformly bounded by the Markov property (RF5). In addition, $\Psi_{r_n}\mathcal{D}_n(f_n) \leq \mathcal{R}(r_*, \mathcal{D}_n)(u)$ with $r_* = \inf_{n \geq 1} r_n$, so $\{\Psi_{r_n}\mathcal{D}_n(f_n)\}_{n \geq 1}$ is a bounded sequence. By estimates (3.2) and (3.4), we have

$$|f_n(p) - f_n(q)| \leq c \left(d(p, q)^{\frac{\log r^*}{\log \rho}} \sup_{n \geq 1} \Psi_{r_n}\mathcal{D}_n(f_n) \right)^{1/2}, \quad \forall n \geq 1, \forall p, q \in \bar{V}_1,$$

where $r^* = \sup_{n \geq 1} r_n$ and $c^2 = \sup_{n \geq 1} \left\{ \frac{4}{r_n^3(1-r_n)} R_0^*(\mathcal{D}_n) \right\}$, and so $\{f_n\}_{n \geq 1}$ is also equicontinuous. Thus, there is a subsequence $\{f_{n_k}\}_{k \geq 1}$ such that f_{n_k} converges

uniformly to a function $f \in C(\bar{V}_1)$. Clearly, f is an extension of u . By Fatou’s lemma,

$$\mathcal{R}(r, \mathcal{D})(u) \leq \Psi_r \mathcal{D}(f) \leq \liminf_{k \rightarrow \infty} \Psi_{r_{n_k}} \mathcal{D}_{n_k}(f_{n_k}) = \liminf_{k \rightarrow \infty} \mathcal{R}(r_{n_k}, \mathcal{D}_{n_k})(u).$$

Combining this with (4.2), we see that

$$\mathcal{R}(r, \mathcal{D})(u) = \lim_{k \rightarrow \infty} \mathcal{R}(r_{n_k}, \mathcal{D}_{n_k})(u).$$

Since the argument works for any sequence $(r'_n, \mathcal{D}'_n) \rightarrow (r, \mathcal{D})$, we actually have

$$\mathcal{R}(r, \mathcal{D})(u) = \lim_{n \rightarrow \infty} \mathcal{R}(r_n, \mathcal{D}_n)(u).$$

The theorem follows immediately since u can be any function in $l(V_0)$. □

5. A fixed point problem

In this section, analogous to the case of PCF self-similar sets (see [22, 34]), we consider the renormalization equation

$$\mathcal{R}_r \mathcal{D} = \lambda \mathcal{D}, \tag{5.1}$$

with $\lambda > 0$. We will prove that for each given $\rho < r < 1$, there always exists a positive λ such that (5.1) has a solution \mathcal{D} in \mathcal{M} . Nevertheless, this is not enough for the construction of a satisfying resistance form on \mathcal{G} for our later purposes. In order that cells of same size are assigned with the same renormalization factors, we will in addition require $\lambda = r^2$, i.e.

$$\mathcal{R}_r \mathcal{D} = r^2 \mathcal{D}. \tag{5.2}$$

The existence and uniqueness of such a solution is the main purpose of this section.

It is natural to consider resistance forms on \mathcal{G} that are symmetric with respect to the reflection symmetry of \mathcal{G} . So we look at the resistance forms on V_0 which are symmetric in the sense that $a_{0,1} = a_{0,2}$ in (3.1). We denote \mathcal{M}_S for the set of all such resistance forms.

THEOREM 5.1.

- (a). For each $\rho < r < 1$, there exists a unique pair of $\lambda(r)$ and $\mathcal{D}(r) \in \mathcal{M}$ (up to constants) satisfying (5.1), where $\lambda(r)$ is decreasing and continuous in r , and $\mathcal{D}(r)$ is in \mathcal{M}_S .
- (b). There exists a unique $\rho < r < 1$ such that (5.2) has a unique (up to constants) solution $\mathcal{D} \in \mathcal{M}$.

We will first prove that for each r , there exist a unique $\lambda(r)$ such that (5.1) has a solution $\mathcal{D}(r)$ in \mathcal{M}_S , then prove that $\mathcal{D}(r)$ is indeed a unique solution (up to constants) in \mathcal{M} . The existence and uniqueness of a solution to (5.2) will follow from the properties of $\lambda(r)$. We divide these into two subsections.

5.1. The existence of a symmetric solution

We begin with some simple observations.

LEMMA 5.2. *Let $\rho < r < 1$ be fixed, and suppose that there is a solution to (5.1). Then the constant λ depends only on r .*

Proof. This follows from a standard argument like the finite graph case [29]. Suppose that $\mathcal{D}, \mathcal{D}'$ are two solutions to (5.1) with λ, λ' being the corresponding constant. Let $u \in l(V_0)$ so that $\frac{\mathcal{D}'(u)}{\mathcal{D}(u)} = \sup_{v \neq \text{constants}} \frac{\mathcal{D}'(v)}{\mathcal{D}(v)} := \theta$, and let f be the harmonic extension of u with respect to $\Psi_r \mathcal{D}$. Then

$$\lambda' \mathcal{D}'(u) = \mathcal{R}_r \mathcal{D}'(u) \leq \Psi_r \mathcal{D}'(f) \leq \theta \Psi_r \mathcal{D}(f) = \theta \mathcal{R}_r \mathcal{D}(u) = \theta \lambda \mathcal{D}(u).$$

This implies that $\lambda' \leq \lambda$. A same argument also shows that $\lambda \leq \lambda'$. □

Inspired by lemma 5.2, we can view the constant λ in (5.1) as a function of r . On the other hand, the problem of solvability of (5.1) can be transferred to a fixed point problem.

DEFINITION 5.3.

(a) *Define*

$$\begin{aligned} \tilde{\mathcal{M}}_S = \left\{ \mathcal{D} \in \mathcal{M} : \mathcal{D}(f) = a(f(q_0) - f(q_1))^2 + a(f(q_0) - f(q_2))^2 \right. \\ \left. + (1 - a)(f(q_1) - f(q_2))^2, 0 < a \leq 1 \right\}, \end{aligned}$$

and for $0 < s \leq 1$,

$$\begin{aligned} \tilde{\mathcal{M}}_S^{[s,1]} = \left\{ \mathcal{D} \in \mathcal{M} : \mathcal{D}(f) = a(f(q_0) - f(q_1))^2 + a(f(q_0) - f(q_2))^2 \right. \\ \left. + (1 - a)(f(q_1) - f(q_2))^2, s \leq a \leq 1 \right\}. \end{aligned}$$

(b) *For each $\mathcal{D} \in \mathcal{M}_S$, there is a unique constant c such that $c\mathcal{D} \in \tilde{\mathcal{M}}_S$, and we denote the resulting form $T\mathcal{D}$. We define $\tilde{\mathcal{R}}_r : \mathcal{M}_S \rightarrow \tilde{\mathcal{M}}_S$ to be the map given by $\tilde{\mathcal{R}}_r = T \circ \mathcal{R}_r$. As before, we write $\tilde{\mathcal{R}}(r, \mathcal{D}) = \tilde{\mathcal{R}}_r(\mathcal{D})$.*

The following lemma will play an essential role.

LEMMA 5.4. *For $\rho < r_0 < r_1 < 1$, there exists $0 < s \leq 1$ such that $\tilde{\mathcal{R}} : [r_0, r_1] \times \mathcal{M}_S \rightarrow \tilde{\mathcal{M}}_S^{[s,1]}$.*

Proof. Let $\mathcal{D} \in \mathcal{M}_S$, $r_0 \leq r \leq r_1$ and R be the resistance metric on V_1 associated with $\Psi_r \mathcal{D}$. For convenience, we write $\mathcal{D}(f) = a(f(q_0) - f(q_1))^2 + a(f(q_0) - f(q_2))^2 + b(f(q_1) - f(q_2))^2$, with $a > 0, b \geq 0$.

First, by the series law for resistances, for any $f \in l(V_1)$, we have

$$\Psi_r \mathcal{D}(f) \geq \sum_{n=0}^{\infty} ar^{-n} (f(F_1^n q_0) - f(F_1^{n+1} q_0))^2 \geq a(1 - r) (f(q_0) - f(q_1))^2,$$

so we have $R(q_0, q_1) \leq \frac{1}{a(1-r)} \leq \frac{1}{a(1-r_1)}$.

Next, let f be the linear function on \mathbb{R}^2 such that $f(q_1) = 0, f(q_2) = 1$ and $f(q_0) = \frac{1}{2}$, so f only depends on the x -coordinate. We introduce a ‘horizontal’ edge relation ‘ \sim_h ’ on V_1 : for distinct $p, q \in V_1$, denote

$$p \sim_h q \text{ if there exists } w \in W_1 \text{ so that } p, q \in \{F_w q_1, F_w q_2\}.$$

For each $p \in V_1$, we write

$$[p]_h = \{q \in V_1 : q \sim_h p, \text{ or } q \sim_h q', q' \sim_h p \text{ for some } q' \in V_1\}.$$

Then we modify f on V_1 into a function $g \in l(V_1)$ as

$$g(p) = \frac{\sum_{q \in [p]_h} f(q)}{\#[p]_h}, \quad \forall p \in V_1.$$

By doing this we have

1. $g(p) = g(q)$ if $[p]_h = [q]_h$;
2. $|g(p) - g(q)| \leq c_1 \rho^n$ if $p, q \in F_w V_0$ with $w \in W_{1,n}$.

Thus, we have

$$\begin{aligned} \Psi_r \mathcal{D}(g) &= \sum_{l=1}^{\infty} \sum_{w \in W_{1,l}} r^{-l+1} \mathcal{D}(g \circ F_w) \\ &\leq 2c_1^2 a \sum_{l=1}^{\infty} r^{-l+1} \rho^{2l} \#W_{1,l} \\ &\leq c_2 a \sum_{l=1}^{\infty} r^{-l} \rho^l = \frac{c_2 \rho}{r - \rho} a \leq \frac{c_2 \rho}{r_0 - \rho} a, \end{aligned}$$

where we use the estimate $\#W_{1,l} \asymp \rho^{-l}$. Thus, g extends to $g \in C(\bar{V}_1)$ by proposition 3.3, and it is direct to check that $g|_{V_0} = f|_{V_0}$. As a consequence, we get $R(q_1, q_2) \geq \frac{r_0 - \rho}{c_2 \rho} a^{-1}$.

Due to the above two estimates, there exists $c_3 > 0$ independent of \mathcal{D} such that

$$\frac{R(q_0, q_1)}{R(q_1, q_2)} \leq c_3.$$

Then an effective resistance calculation gives that $\tilde{\mathcal{R}}(r, \mathcal{D}) \in \tilde{\mathcal{M}}_S^{[\frac{1}{2c_3}, 1]}$. The lemma follows. □

By using lemmas 5.2 and 5.4 and theorem 4.2, we can easily prove the following proposition.

PROPOSITION 5.5. *Let $\rho < r < 1$, there always exists a solution to (5.1) in \mathcal{M}_S , with λ uniquely determined by r . In addition, regarding λ as a function of r , $\lambda(r)$ is decreasing and continuous in r .*

Proof. First, we have $\tilde{\mathcal{R}}_r : \tilde{\mathcal{M}}_S^{[s,1]} \rightarrow \tilde{\mathcal{M}}_S^{[s,1]}$ for some $s > 0$ by lemma 5.4. Together with theorem 4.2, the existence of a fixed point of $\tilde{\mathcal{R}}_r$ is then an immediate consequence.

Next, let $r_1 < r_2$, and assume that $\mathcal{R}_{r_1} \mathcal{D}_1 = \lambda(r_1) \mathcal{D}_1$ and $\mathcal{R}_{r_2} \mathcal{D}_2 = \lambda(r_2) \mathcal{D}_2$. Let $u \in l(V_0)$ so that $\frac{\mathcal{D}_2(u)}{\mathcal{D}_1(u)} = \sup_{v \neq \text{constants}} \frac{\mathcal{D}_2(v)}{\mathcal{D}_1(v)} := \theta$, and let f be the harmonic extension of u with respect to $\Psi_{r_1} \mathcal{D}_1$, then we have $\lambda(r_2) \mathcal{D}_2(u) \leq \Psi_{r_2} \mathcal{D}_2(f) \leq \theta \Psi_{r_1} \mathcal{D}_1(f) = \theta \lambda(r_1) \mathcal{D}_1(u)$. So we get $\lambda(r_2) \leq \lambda(r_1)$.

Finally, let $r_n \rightarrow r$, and let $\mathcal{D}_n \in \tilde{\mathcal{M}}_S$ be a sequence of solutions to $\mathcal{R}_{r_n} \mathcal{D}_n = \lambda(r_n) \mathcal{D}_n$. Clearly, we have $\rho < \inf_{n \geq 1} r_n < \sup_{n \geq 1} r_n < 1$, so $\{\mathcal{D}_n\}_{n \geq 1} \subset \tilde{\mathcal{M}}_S^{[s,1]}$ for some $s > 0$ by lemma 5.4. Thus, there exists a subsequence $\{n_k\}_{k \geq 1}$ such that \mathcal{D}_{n_k} converges to some $\mathcal{D} \in \tilde{\mathcal{M}}_S$ and $\lambda(r_{n_k})$ converges. By theorem 4.2, we conclude that $\mathcal{R}_r \mathcal{D} = (\lim_{k \rightarrow \infty} \lambda(r_{n_k})) \mathcal{D}$. So $\lambda(r) = \lim_{k \rightarrow \infty} \lambda(r_{n_k})$. Since the argument works for any sequence $r_n \rightarrow r$, $\lambda(r)$ is continuous in r . □

We have an easy estimate of $\lambda(r)$.

LEMMA 5.6. For $\rho < r < 1$, we have $\left(\frac{1}{1-r} - \frac{r}{2+2r+2r^2}\right)^{-1} \leq \lambda(r) \leq \frac{2}{2+r}$.

Proof. We consider a function $u \in l(V_0)$ with $u(q_0) = 0$ and $u(q_1) = u(q_2) = 1$. Without loss of generality, we assume the solution $\mathcal{D} \in \mathcal{M}_S$ has $\mathcal{D}(u) = 2$. To get an upper bound for $\lambda(r)$, we construct an extension $f \in l(V_1)$ of u by setting

$$f(p) = \begin{cases} 0, & \text{if } p = q_0, \\ \frac{2}{2+r}, & \text{if } p \in \{F_1 q_0, F_2 q_0\}, \\ 1, & \text{if } p \in F_1 V_1 \cup F_2 V_1 \setminus \{F_1 q_0, F_2 q_0\}. \end{cases}$$

Then the upper bound follows easily from the following estimate:

$$\mathcal{R}_\lambda \mathcal{D}(u) \leq \Psi_r \mathcal{D}(f) = \left(\left(\frac{2}{2+r}\right)^2 + 2r^{-1} \left(1 - \frac{2}{2+r}\right)^2 \right) \mathcal{D}(u) = \frac{2}{2+r} \mathcal{D}(u).$$

To get the lower bound, we look at a subgraph in (V_1, \sim) , whose vertices are $\{F_i^l q_0\}_{i,l}$ with $i \in \{1, 2\}$ and $l \geq 0$, together with

$$\begin{aligned} p_{i,0} &= F_i q_0, & p_{i,1} &= F_i F_j q_0, & p_{i,2} &= F_i F_j F_i q_0, \\ p_{i,3} &= F_i F_j F_i^2 q_0, & p_{i,4} &= F_i^2 F_j q_0, & p_{i,5} &= F_i^2 q_0, \end{aligned}$$

with $i, j \in \{1, 2\}$ and $j \neq i$, and edges inherited from (V_1, \sim) (with horizontal edges deleted), see figure 5. Let $f \in l(V_1)$ be the harmonic extension of u , denote $c_l = r^{-l-1}$ for $l \in \{0, 1, 2\}$, and $c_l = r^{l-6}$ for $l \in \{3, 4\}$, then the lower bound follows from the estimate that

$$\begin{aligned} \mathcal{R}_r \mathcal{D}(u) = \Psi_r \mathcal{D}(f) &\geq \sum_{i=1,2} \left(\sum_{l=0}^{\infty} r^{-l} (f(F_i^l q_0) - f(F_i^{l+1} q_0))^2 + \sum_{l=0}^4 c_l (f(p_{i,l}) - f(p_{i,l+1}))^2 \right) \\ &\geq 2 \left(\frac{1}{1-r} - \frac{r}{2+2r+2r^2} \right)^{-1} = \left(\frac{1}{1-r} - \frac{r}{2+2r+2r^2} \right)^{-1} \mathcal{D}(u), \end{aligned}$$




Figure 5. The subgraph of (V_1, \sim) constructed in lemma 5.6.

where the last inequality can be done by an easy computation of the effective resistances on the subgraph (for $i \in \{1, 2\}$, firstly connecting the resistors along $\{p_{i,l}\}_{l=0}^5$ in series, secondly connecting the resulting effective resistor with the resistor between $p_{i,0}$ and $p_{i,5}$ in parallel; lastly connecting all the resistors in series). \square

Using proposition 5.5 and lemma 5.6, we arrive at the main result of this subsection, concerning the solvability of (5.2).

THEOREM 5.7. *There exists a unique $\rho < r < 1$ such that (5.2) has a solution $\mathcal{D} \in \mathcal{M}_S$.*

Proof. By proposition 5.5, we see that there is a continuous function $\lambda(r)$ so that $\mathcal{R}_\lambda(\mathcal{D}) = \lambda(r)\mathcal{D}$ has a solution. Noticing that

$$\lambda(\rho) \geq \left(\frac{1}{1-\rho} - \frac{\rho}{2+2\rho+2\rho^2} \right)^{-1} > 1 - \rho = \rho^2, \text{ and } \lambda(1) \leq \frac{2}{3} < 1,$$

there exists $\rho < r < 1$ such that $\lambda(r) = r^2$ by lemma 5.6. The uniqueness follows from the fact that $\lambda(r)$ is decreasing in r , while r^2 is strictly increasing. \square

REMARK. *We can see the uniqueness of r from another point of view. Let $\theta = \frac{\log r}{\log \rho}$, we will see in § 7 that $\theta + d_H$ is the walk dimension of the resulting diffusion process on the metric measure space (\mathcal{G}, d, μ_H) , whose uniqueness is shown in [17, theorem 4.6] under some weak conditions on the heat kernel.*

5.2. The uniqueness

In this subsection, we consider the uniqueness of the solution to (5.1) or (5.2). The proof is inspired by Sabot’s work [34].

THEOREM 5.8. *Let $\rho < r < 1$ and $\mathcal{D} \in \mathcal{M}_S$ be a symmetric solution to (5.1). Then \mathcal{D} is the unique solution in \mathcal{M} to (5.1).*

For fixed $\rho < r < 1$ and $\mathcal{D} \in \mathcal{M}_S$ satisfying (5.1), for convenience, we always write

1. h_s for the harmonic function with $h_s(q_0) = 0, h_s(q_1) = h_s(q_2) = 1$, and denote $E_s = \{f \in l(V_0) : f(q_0) = 0, f(q_1) = f(q_2) = c, c \in \mathbb{R}\}$;
2. h_a for the harmonic function with $h_a(q_0) = 0, h_a(q_1) = -h_a(q_2) = 1$, and denote $E_a = \{f \in l(V_0) : f(q_0) = 0, f(q_1) = -f(q_2) = c, c \in \mathbb{R}\}$.

Both h_s, h_a are harmonic with respect to $\Psi_r \mathcal{D}$ on $\bar{V}_1 \setminus V_0$, i.e. $\Psi_r \mathcal{D}(h_s, f) = \Psi_r \mathcal{D}(h_a, f) = 0$ for any $f \in \text{Dom}(\Psi_r \mathcal{D})$ such that $f|_{V_0} = 0$.

LEMMA 5.9. For r, \mathcal{D} as above, we have

$$h_s(F_1q_0) = h_s(F_2q_0) = \lambda(r), \quad h_a(F_1q_0) = -h_a(F_2q_0) = \eta,$$

for some $|\eta| < \lambda(r)$.

Proof. For convenience, we write \mathcal{D} in the form $\mathcal{D}(f) = a(f(q_0) - f(q_1))^2 + a(f(q_0) - f(q_2))^2 + b(f(q_1) - f(q_2))^2$, with $a > 0, b \geq 0$.

First, let $h = 1 - h_s$, we have $\mathcal{R}_r \mathcal{D}(h_s, h) = -2a\lambda(r)$. On the other hand, let $f \in l(V_1)$ be defined as $f(p) = \delta_{q_0,p}$, then clearly $f \in \text{Dom}(\Psi_r \mathcal{D})$, and $f|_{V_0} = h|_{V_0}$. Since h_s is harmonic,

$$\Psi_r \mathcal{D}(h_s, h) = \Psi_r \mathcal{D}(h_s, f) = -ah_s(F_1q_0) - ah_s(F_2q_0).$$

This shows the first assertion since $\mathcal{R}_r \mathcal{D}(h_s, h) = \Psi_r \mathcal{D}(h_s, h)$.

Next, by the symmetry of \mathcal{D} , there exists a number η such that $h_a(F_1q_0) = -h_a(F_2q_0) = \eta$. We need to show that $|\eta| < \lambda(r)$. We consider the matrix M such that

$$(h(F_1q_0), h(F_2q_0))^t = M (h(q_1), h(q_2))^t,$$

holds for any harmonic function h with $h(q_0) = 0$. Due to the Perron–Frobenius theorem, it suffices to show that each entry of M is positive. This can be deduced by proving the harmonic function h_1 with boundary value $h_1(q_1) = 1, h_1(q_0) = h_1(q_2) = 0$ is positive on $V_1 \setminus V_0$. To see this, we assume there exists $p \in V_1 \setminus V_0$ such that $h_1(p) = 0$. Let $\psi_p \in \text{Dom}(\Psi_r \mathcal{D})$ be defined as $\psi_p(q) = \delta_{p,q}$, then $\Psi_r \mathcal{D}(\psi_p, h_1) = 0$, so $h_1(p)$ is the weighted average of its neighbours. Thus, h_1 is zero on the neighbours of p . Repeating the argument, we see that $h_1|_{V_1} = 0$. A contradiction. □

Proof of theorem 5.8. Assume there is another solution $\mathcal{D}' \in \mathcal{M}$ to (5.1).

Firstly, we will show that \mathcal{D}' is also symmetric. By diagonalizing \mathcal{D}' with respect to \mathcal{D} , we have two 1-dimensional non-constant subspaces L_1, L_2 of $l(V_0)$ such that

1. L_1, L_2 are orthogonal with respect to both \mathcal{D} and \mathcal{D}' ;
2. $\mathcal{D}'|_{L_1} = \kappa_1 \mathcal{D}|_{L_1}$ and $\mathcal{D}'|_{L_2} = \kappa_2 \mathcal{D}|_{L_2}$, with $0 < \kappa_1 < \kappa_2$.

Let $u \in L_2$ and h_u be the harmonic extension of u with respect to $\Psi_r \mathcal{D}$. Then

$$\begin{aligned} \lambda(r)\mathcal{D}'(u) &= \kappa_2\lambda(r)\mathcal{D}(u) = \kappa_2\Psi_r\mathcal{D}(h_u) = \sum_{w \in W_1} r^{-|w|+1}\kappa_2\mathcal{D}(h_u \circ F_w) \\ &\geq \sum_{w \in W_1} r^{-|w|+1}\mathcal{D}'(h_u \circ F_w) = \Psi_r\mathcal{D}'(h_u) \geq \lambda(r)\mathcal{D}'(u). \end{aligned}$$

This implies that for each $w \in W_1$, $\mathcal{D}'(h_u \circ F_w) = \kappa_2\mathcal{D}(h_u \circ F_w)$, and thus $h_u \circ F_w \in L_2 + constants$. In particular, we have $h_u \circ F_0 \in L_2 + constants$, which means $L_2 + constants$ is an invariant space under the mapping u to $h_u \circ F_0$. By lemma 5.9, we see that $L_2 + constants$ is either $E_s + constants$ or $E_a + constants$. Thus, we have $\mathcal{D}' \in \mathcal{M}_S$.

Secondly, from the above argument, it is not hard to see that $h_s \circ F_w \in E_s + constants$ and $h_a \circ F_w \in E_a + constants$, for any $w \in W_1$.

Lastly, arbitrarily pick a $\tilde{\mathcal{D}} \in \mathcal{M}_S$, we will prove that $\tilde{\mathcal{D}}$ must also solve (5.1). However, this will make $\tilde{\mathcal{R}}_r(\mathcal{M}_S) = \tilde{\mathcal{M}}_S$, which obviously contradicts Lemma 5.4. To achieve this purpose, let \tilde{h}_s and \tilde{h}_a be the harmonic functions with respect to $\Psi_r\tilde{\mathcal{D}}$, with the same boundary value on V_0 as h_s, h_a . By following the same argument as [34, lemma 5.9] due to Sabot, we can see that $\tilde{h}_s = h_s$ and $\tilde{h}_a = h_a$. For convenience of readers, we reproduce the proof here. Write $g = h_s - \tilde{h}_s$. Also, for each $w \in W_1$, let $g_{w,s} \in E_s + constants, g_{w,a} \in E_a + constants$ such that $g_w =: g \circ F_w = g_{w,s} + g_{w,a}$. Then, denoting $h_{w,s} =: h_s \circ F_w$, we can see that

$$\begin{aligned} \Psi_r\tilde{\mathcal{D}}(g) &= \Psi_r\tilde{\mathcal{D}}(h_s, g) = \sum_{w \in W_1} r^{-|w|+1}\tilde{\mathcal{D}}(h_{w,s}, g_w) = \sum_{w \in W_1} r^{-|w|+1}\tilde{\mathcal{D}}(h_{w,s}, g_{w,s}) \\ &= c \sum_{w \in W_1} r^{-|w|+1}\mathcal{D}(h_{w,s}, g_{w,s}) \\ &= c \sum_{w \in W_1} r^{-|w|+1}\mathcal{D}(h_{w,s}, g_w) = c\Psi_r\mathcal{D}(h_s, g) = 0, \end{aligned}$$

for some constant c , with $h_{s,w} =: h_s \circ F_w$, where the first equality is due to the fact that $g|_{V_0} = 0$. Thus, $g = 0$ as desired. As a consequence, we can easily see that, $\tilde{\mathcal{D}}$ is a solution to (5.1), so we arrive at the desired contradiction. \square

Finally, theorem 5.1 immediately follows from proposition 5.5, theorems 5.7 and 5.8.

6. Construction of the Dirichlet form on \mathcal{G}

We will construct a resistance form on the golden ratio Sierpinski gasket \mathcal{G} in this section. Let $\rho < r < 1$, \mathcal{D} be the unique solution to (5.2), i.e. $\mathcal{R}_r\mathcal{D} = r^2\mathcal{D}$. We will focus on this standard form in most contents. For short, we write

$$\theta = \frac{\log r}{\log \rho}, \quad \rho_w = \prod_{n=1}^{|w|} \rho_{w_n}, \quad r_w = \rho_w^\theta,$$

with $\rho_0 = \rho^2$ and $\rho_1 = \rho_2 = \rho$. Obviously, ρ_w is the contraction ratio of F_w .

The following definition is similar to the construction in [22], though we use the infinite graphs at each level.

DEFINITION 6.1.

- (a) For $m \geq 0$ and $f \in C(\bar{V}_m)$, we write $\mathcal{D}^{(m)}(f) = \sum_{w \in W_m} r_w^{-1} \mathcal{D}(f \circ F_w)$, and $\mathcal{F}^{(m)} = \{f \in C(\bar{V}_m) : \mathcal{D}^{(m)}(f) < \infty\}$. In addition, for $f, g \in \mathcal{F}^{(m)}$, we define

$$\mathcal{D}^{(m)}(f, g) = \sum_{w \in W_m} r_w^{-1} \mathcal{D}(f \circ F_w, g \circ F_w).$$

- (b) Define $\mathcal{F} = \{f \in C(\mathcal{G}) : \lim_{m \rightarrow \infty} \mathcal{D}^{(m)}(f) < \infty\}$. For $f, g \in \mathcal{F}$, define

$$\mathcal{E}(f, g) = \lim_{m \rightarrow \infty} \mathcal{D}^{(m)}(f, g).$$

It follows from the definition of $\Psi_r \mathcal{D}$, $\mathcal{D}^{(1)} = r^{-2} \Psi_r \mathcal{D}$. The limit in (b) always exists due to fact that

$$\begin{aligned} \mathcal{D}^{(m+1)}(f) &= \sum_{w \in W_{m+1}} r_w^{-1} \mathcal{D}(f \circ F_w) = \sum_{w \in W_m} r_w^{-1} r^{-2} \Psi_r \mathcal{D}(f \circ F_w) \\ &\geq \sum_{w \in W_m} r_w^{-1} \mathcal{D}(f \circ F_w) = \mathcal{D}^{(m)}(f). \end{aligned}$$

In the rest of this section, we will show that $(\mathcal{E}, \mathcal{F})$ is a good form.

LEMMA 6.2. For $m \geq 0$, $(\mathcal{D}^{(m)}, \mathcal{F}^{(m)})$ is a resistance form on \bar{V}_m . In addition, let

$$R_m(p, q) = \sup_{f \in \mathcal{F}^{(m)}} \frac{|f(p) - f(q)|^2}{\mathcal{D}^{(m)}(f)},$$

then we have $R_n(p, q) = R_m(p, q)$ if $p, q \in \bar{V}_m$ and $n \geq m$.

Proof. (RF1) and (RF5) are trivial. We only need to verify (RF2)–(RF4). For convenience, we focus on $(\mathcal{D}^{(2)}, \mathcal{F}^{(2)})$ only, while for larger m , the same proof works inductively. (RF2). Let $\{f_k\}_{k \geq 1}$ be a Cauchy sequence in $\mathcal{F}^{(2)}$. Then, $f_k|_{\bar{V}_1}$ converges in $\mathcal{F}^{(1)}$ to some \tilde{f} in $\mathcal{F}^{(1)}$, since $(\mathcal{D}^{(1)}, \mathcal{F}^{(1)})$ is a resistance form. Also, for each $w \in W_1$, $f_k \circ F_w$ converges in $\mathcal{F}^{(1)}$ to a function \tilde{f}_w . Now, define $f \in l(\bar{V}_2)$ such that $f \circ F_w = \tilde{f}_w$ and $f|_{\bar{V}_1 \setminus V_1} = \tilde{f}$. We show that $f \in C(\bar{V}_2)$. It suffices to prove that f is continuous at any point $p \in \bar{V}_1 \setminus V_1$. In fact, for any ε , there exists δ and N such that 1. for $q \in B_\delta(p) \cap \bar{V}_1$, we have $|f(p) - f(q)| < \varepsilon$; 2. for $w \in \bigcup_{n=N}^\infty W_{1,n}$ and $q, q' \in F_w \bar{V}_1$, we have $|f(q) - f(q')| < \varepsilon$. This follows from the fact that $\mathcal{D}^{(1)}(f \circ F_w) \leq r_w \sup_{k \geq 1} \mathcal{D}^{(2)}(f_k)$. The continuity of f follows immediately. Lastly, by using Fatou’s lemma, we can directly check that f_k converges to f in $\mathcal{F}^{(2)}$. (RF3). First, we observe that the minimal energy extension of $f \in \mathcal{F}^{(1)}$ to $l(V_2)$ is continuous by a same argument as in (RF2). Let V be a finite set and $u \in l(V)$. First, we always have $f_1 \in \mathcal{F}^{(1)}$ such that $f_1|_{V \cap \bar{V}_1} = u|_{V \cap \bar{V}_1}$. Then we can extend f_1 to be a desired function in $\mathcal{F}^{(2)}$. (RF4). Let $p, q \in \bar{V}_2$ and $f \in \mathcal{F}^{(2)}$. If

$p \in \bar{V}_1$, we let $p' = p$; otherwise we choose $p' \in V_1$ so that $p, p' \in F_w \bar{V}_1$ for some $w \in W_1$, and thus

$$\mathcal{D}^{(2)}(f) \geq r_w^{-1} \mathcal{D}^{(1)}(f \circ F_w) \geq c_1 (f(p) - f(p'))^2,$$

for some $c_1 > 0$. Also, we define q' in the same manner. Note that $\mathcal{D}^{(2)}(f) \geq \mathcal{D}^{(1)}(f) \geq c_2 (f(p') - f(q'))^2$ for some $c_2 > 0$, it then follows that

$$\begin{aligned} \mathcal{D}^{(2)}(f) &\geq \min\{c_1, c_2\} \left((f(p) - f(p'))^2 + (f(p') - f(q'))^2 + (f(q') - f(q))^2 \right) \\ &\geq c_3 (f(p) - f(q))^2, \end{aligned}$$

for some $c_3 > 0$. (RF4) follows immediately. Thus, we have proved that $(\mathcal{D}^{(2)}, \mathcal{F}^{(2)})$ is a resistance form on \bar{V}_2 . The claim that $R_2(p, q) = R_1(p, q)$ for $p, q \in \bar{V}_1$ is obvious. The same arguments can be used inductively for $m \geq 3$. □

In some situations, it is convenient to involve words in \tilde{W}_* .

LEMMA 6.3. *Let $w \in \tilde{W}_*$ and m be the number of 0 's in w . Then we have*

$$\mathcal{D}^{(1)}(f \circ F_w) \leq r_w \mathcal{D}^{(m+1)}(f),$$

for any $f \in \mathcal{F}^{(m+1)}$. As a consequence, there is a constant $c > 0$ such that, for any $p, q \in F_w \bar{V}_1$, we have

$$R_{m+1}(p, q) \leq c d(p, q)^\theta.$$

Proof. Noticing that $\{w\tau : \tau \in W_1\} \subset W_{m+1}$, the first statement follows. The second statement follows from the first statement and proposition 3.3: for any $p, q \in F_w \bar{V}_1$,

$$R_{m+1}(p, q) \leq r_w R_1(F_w^{-1}p, F_w^{-1}q) \leq cr_w d(F_w^{-1}p, F_w^{-1}q)^\theta = c d(p, q)^\theta,$$

holds for some constant $c > 0$, where the first inequality follows from the first statement, and the second inequality follows from proposition 3.3. □

Using lemmas 6.2 and 6.3, we have the following estimate of the resistance metric.

LEMMA 6.4. *For $m \geq 0$ and $p, q \in \bar{V}_m$, define $\tilde{R}(p, q) = R_m(p, q)$. Then $\tilde{R}(p, q)$ is well defined on $(\bigcup_{m \geq 0} \bar{V}_m) \times (\bigcup_{m \geq 0} \bar{V}_m)$, and we have $\tilde{R}(p, q) \leq c d(p, q)^\theta$ for some $c > 0$.*

Proof. First, we claim that there is a constant $c_1 > 0$ such that

$$\tilde{R}(p, q) \leq c_1 \rho_w^\theta, \quad \forall w \in \tilde{W}_*, \forall p, q \in F_w \mathcal{G} \cap \left(\bigcup_{m \geq 0} \tilde{V}_m \right).$$

We first consider the case $q \in F_w \tilde{V}_1$. Assume that $p \in F_w \tilde{V}_n$ for some $n \geq 1$, then we can find $\tau \in W_{n-1}$ such that $p \in F_w F_\tau \tilde{V}_1$. We can then find a sequence

$$q = p_0, p_1, \dots, p_{|\tau|+1} = p,$$

such that $p_i \in F_w F_{[\tau]_{i-1}} \tilde{V}_1 \cap F_w F_{[\tau]_i} \tilde{V}_1$ for $1 \leq i \leq |\tau|$. As a consequence, by using lemma 6.3, we see that

$$\tilde{R}(p, q) \leq \sum_{i=0}^{|\tau|} c_2 d(p_i, p_{i+1})^\theta \leq \sum_{i=0}^{|\tau|} c_2 (\rho_w \rho^i)^\theta \leq \frac{c_2}{1 - \rho^\theta} \rho_w^\theta,$$

where c_2 is the same constant in lemma 6.3. For general q , we only need to set $c_1 = \frac{2c_2}{1 - \rho^\theta} \rho_w^\theta$. Now, let $p, q \in \bigcup_{m=0}^\infty \tilde{V}_m$. We choose $w, w' \in \tilde{W}_*$ such that $p \in F_w \mathcal{G}$, $q \in F_{w'} \mathcal{G}$ and

$$\rho d(p, q) \leq \rho_w, \rho_{w'} < \rho^{-1} d(p, q).$$

In addition, we can find a chain

$$w = w^{(0)}, w^{(1)}, \dots, w^{(k)} = w'$$

such that $\min\{\rho_w, \rho_{w'}\} \leq \rho_{w^{(i)}} < \rho^{-2} \min\{\rho_w, \rho_{w'}\}$ of length at most c_3 , where c_3 is a constant independent of p, q . By choosing a sequence $p = p_0, p_1, \dots, p_{k+1} = q$ such that $p_i \in F_{w^{(i-1)}} \tilde{V}_1 \cap F_{w^{(i)}} \tilde{V}_1$, $1 \leq i \leq k$, we get the desired estimate as above. \square

Now, we can show that $(\mathcal{E}, \mathcal{F})$ is a good form.

THEOREM 6.5. *$(\mathcal{E}, \mathcal{F})$ defined in definition 6.1 is a strongly local regular resistance form on \mathcal{G} .*

Proof. First, we claim that $(\mathcal{E}, \mathcal{F})$ is a resistance form on $\bigcup_{m \geq 0} \tilde{V}_m$. (RF1) and (RF5) are obvious given lemma 6.2. Observing that by iterating the minimal energy extension, we can extend any $f \in \mathcal{F}^{(m)}$ to $f \in \mathcal{F}$ thanks to the upper-bound estimate of the resistance metric in lemma 6.4. (RF2), (RF3) and (RF4) are then easy to show with lemma 6.2. In addition, we see that

$$\tilde{R}(p, q) = R(p, q) := \sup_{f \in \mathcal{F}} \frac{|f(p) - f(q)|^2}{\mathcal{E}(f)}, \quad \forall p, q \in \bigcup_{m \geq 0} \tilde{V}_m.$$

Next, to prove that $(\mathcal{E}, \mathcal{F})$ is a resistance form on \mathcal{G} , we need to show that \mathcal{F} separates points in \mathcal{G} , just like in proposition 3.3. It suffices to prove that \mathcal{F} is dense in $C(\mathcal{G})$. Let $u \in C(\mathcal{G})$, we fix N large enough so that $|u(x) - u(y)| < \varepsilon$ if $x, y \in F_w K$ and $|w| \geq N$. We can apply proposition 4.3 to create $f \in \mathcal{F}$ such that $\|f - u\|_{L^\infty(\mathcal{G})} < 2\varepsilon$. First, we find $f_1 \in \mathcal{F}^{(1)}$ such that $1. \|f_1 - u\|_{\tilde{V}_1} < \varepsilon;$

2. $f_1(p) = u(p)$ for any $p \in \bigcup_{n=1}^N \bigcup_{w \in W_{1,n}} F_w V_0$. Then we apply harmonic extension to f_1 on $\bar{V}_2 \setminus \bigcup_{n=1}^N \bigcup_{w \in W_{1,n}} F_w \bar{V}_1$. On the cells $F_w \bar{V}_1$ with $|w| < N$, we apply the same construction to get f_2 , but with $N - 2$ replacing N this time. After $k = [N/2] + 1$ times, we get $f_k \in \mathcal{F}^{(k)}$ such that $\|f_k - u|_{\bar{V}_k}\|_{L^\infty} < 2\varepsilon$. Since all cells have size smaller than ρ^N , by harmonically extending, we get $f \in \mathcal{F}$ such that $\|f - u\|_{L^\infty(\mathcal{G})} < 2\varepsilon$. Thus, $(\mathcal{E}, \mathcal{F})$ is regular resistance form on \mathcal{G} . It remains to show that the form is strongly local. Let $f, g \in \mathcal{F}$ with $\text{supp}(f) \cap \text{supp}(g) = \emptyset$, then there exists $\varepsilon > 0$ such that $d(\text{supp}(f), \text{supp}(g)) > \varepsilon$. Thus, we have $\mathcal{D}^{(n)}(f, g) = 0$ for large n , because the supports of f and g are suitably separated by n -cells for large n . By taking the limit, we see that $\mathcal{E}(f, g) = 0$. Clearly $1 \in \mathcal{F}$ with $\mathcal{E}(1) = 0$, and it follows that the form is strongly local. \square

In the remaining part of this section, we would like to characterize $(\mathcal{E}, \mathcal{F})$ as the unique self-similar form associated with the infinite IFS $\{F_w\}_{w \in W_1}$.

THEOREM 6.6. *The resistance form $(\mathcal{E}, \mathcal{F})$ satisfies the following properties:*

- (a) $\mathcal{F} \subset C(\mathcal{G})$.
- (b) For each $f \in \mathcal{F}$, we have $f \circ F_w \in \mathcal{F}$ for all $w \in W_1$, and in addition,

$$\mathcal{E}(f) = \sum_{w \in W_1} \rho_w^{-\theta} \mathcal{E}(f \circ F_w).$$

- (c) Conversely, let $f \in C(\mathcal{G})$, if $f \circ F_w \in \mathcal{F}$ for all $w \in W_1$, and $\sum_{w \in W_1} \rho_w^{-\theta} \mathcal{E}(f \circ F_w) < \infty$, then $f \in \mathcal{F}$.

Moreover, $(\mathcal{E}, \mathcal{F})$ (up to constants) and θ are uniquely determined by the above properties.

Proof. The claimed properties of $(\mathcal{E}, \mathcal{F})$ are immediate consequences of the construction.

The uniqueness follows by a well-known argument, but in the infinite graph version. Let $(\mathcal{E}', \mathcal{F}')$ be another form satisfying the above properties with θ' replacing θ . Define \mathcal{D}' to be the trace of \mathcal{E}' onto V_0 , and write $r'_w = \rho_w^{\theta'}$, $r' = \rho^{\theta'}$. For any $u \in l(V_0)$, let h_u be the harmonic extension of u to \mathcal{F}' , then we can see that

$$\begin{aligned} \mathcal{D}'(u) &= \mathcal{E}'(h_u) = \sum_{w \in W_1} r'^{-1}_w \mathcal{E}'(h_u \circ F_w) \geq \sum_{w \in W_1} r'^{-1}_w \mathcal{D}'((h_u \circ F_w)|_{V_0}) \\ &\geq r'^{-2} \mathcal{R}_{r'} \mathcal{D}'(u), \end{aligned}$$

where $\mathcal{R}_{r'}$ is the renormalization map introduced in definition 4.1, and we use properties (a) and (b) in the inequalities.

On the other hand, we can perform the harmonic extension of u in two steps: first, we extend u to $f_1 \in C(\bar{V}_1)$ so that f_1 minimizes $\Psi_{r'} \mathcal{D}'$, then we take harmonic extension of f_1 on each cell $F_w \mathcal{G}$, $w \in W_1$, to $f \in C(\mathcal{G})$, by using property (a) and the Markov property (RF5). In addition, $f \in \mathcal{F}'$ by the property (c). Then, by

property (b),

$$r'^{-2}\mathcal{R}_{r'}\mathcal{D}'(u) = r'^{-2}\Psi_{r'}\mathcal{D}'(f_1) = \sum_{w \in W_1} r'_w{}^{-1}\mathcal{E}'(f \circ F_w) = \mathcal{E}'(f) \geq \mathcal{D}'(u).$$

Thus, we get $\mathcal{R}_{r'}\mathcal{D}' = r'^2\mathcal{D}'$, which implies that $\mathcal{D}' = \mathcal{D}$ and $\theta' = \theta$ by theorem 5.1. Finally, by a similar argument, one can easily find that the restriction of \mathcal{E}' to \bar{V}_m is $\mathcal{D}^{(m)}$, and the claim that $\mathcal{E}' = \mathcal{E}$ follows immediately by taking the limit. \square

Finally, the form $(\mathcal{E}, \mathcal{F})$ is decimation invariant with respect to the graph-directed construction in definition 2.2.

DEFINITION 6.7. Using the same notation as in definition 2.2, let $(\mathcal{E}_1, \mathcal{F}_1) = (\mathcal{E}, \mathcal{F})$, and define $(\mathcal{E}_2, \mathcal{F}_2)$ as follows:

$$\begin{cases} \mathcal{E}_2(f, g) = \sum_{w \in W_1, F_w\mathcal{G} \subset K_2} \rho_w^{-\theta} \mathcal{E}(f \circ F_w, g \circ F_w), \\ \mathcal{F}_2 = \{f \in C(K_2) : f \circ F_w \in \mathcal{F}, \forall w \in W_1 \text{ such that } F_w\mathcal{G} \subset K_2, \mathcal{E}_2(f) < \infty\}. \end{cases}$$

It is not hard to verify that $(\mathcal{E}_2, \mathcal{F}_2)$ is a resistance form on K_2 . Moreover, we have

THEOREM 6.8. Recall the notation of definition 2.2, and write ρ_{e_j} for the similarity ratio of ψ_{e_j} , $1 \leq j \leq 6$. Let $(\mathcal{E}_i, \mathcal{F}_i), i = 1, 2$ be defined as in definition 6.7. Then, for $f_i \in \mathcal{F}_i, i = 1, 2$, we have $f_{e_{j,1}} \circ \psi_{e_j} \in \mathcal{F}_{e_{j,2}}$ for $1 \leq j \leq 6$ and

$$\mathcal{E}_1(f_1) = \sum_{j=1}^2 \rho_{e_j}^{-\theta} \mathcal{E}_{e_{j,2}}(f_1 \circ \psi_{e_j}), \quad \mathcal{E}_2(f_2) = \sum_{j=3}^6 \rho_{e_j}^{-\theta} \mathcal{E}_{e_{j,2}}(f_2 \circ \psi_{e_j}).$$

Conversely, let $f_1 \in C(K_1)$, if $f_1 \circ \psi_{e_j} \in \mathcal{F}_{e_{j,2}}$ for $j = 1, 2$, then $f_1 \in \mathcal{F}_1$. The same holds for $(\mathcal{E}_2, \mathcal{F}_2)$.

REMARK. At the end of this section, we remark that a same construction can be applied to get some non-standard self-similar forms on \mathcal{G} with respect to the infinite IFS $\{F_w\}_{w \in W_1}$, by starting with any solution $R_{r'}\mathcal{D}' = \lambda(r')\mathcal{D}'$. Theorems 6.5 and 6.8 still hold for the forms, with slight changes of the renormalization factors. Nevertheless, the good heat kernel estimate (theorem 7.4) will not hold, but it is possible to get a heat kernel estimate in the form of Hambly and Kumagai's on PCF self-similar sets [18].

7. Transition density estimate

Let μ_H be the normalized Hausdorff measure on \mathcal{G} . $(\mathcal{E}, \mathcal{F})$ becomes a local regular Dirichlet form on $L^2(\mathcal{G}, \mu_H)$ ($L^2(\mathcal{G})$ for short) in a standard way (see [22, theorem 2.4.1]). By the celebrated result [15, theorem 7.2.1], there is a Hunt process $X =$

$(\mathbb{P}^x, x \in \mathcal{G}, X_t, t \geq 0)$ associated with $(\mathcal{E}, \mathcal{F})$ such that

$$\mathbb{E}^x[f(X_t)] = P_t f(x), \quad \text{a.e. } x \in \mathcal{G},$$

where $(P_t)_{t \geq 0}$ is the associated semigroup. In this last section, we will show that X is a fractional diffusion. We recall from Barlow’s book [3, § 3], for the definition of this fractional diffusion.

DEFINITION 7.1. A Markov process $X = (\mathbb{P}^x, x \in \mathcal{G}, X_t, t \geq 0)$ is a fractional diffusion on the fractional metric space $(\mathcal{G}, d_g, \mu_H)$ (see §2) if (a). X is a conservative Feller diffusion with state space \mathcal{G} ; (b). X is μ_H -symmetric; (c). X has a symmetric transition density $p(t, x, y) = p(t, y, x)$, $t > 0$, $x, y \in \mathcal{G}$, which satisfies the Chapman–Kolmogorov equations and is jointly continuous for $t > 0$; (d). There exist a constant β and $c_1 - c_4 > 0$, such that for $0 < t \leq 1$,

$$\begin{aligned} c_1 t^{-d_H/\beta} \exp\left(-c_2 \left(\frac{d_g(x, y)^\beta}{t}\right)^{\frac{1}{\beta-1}}\right) &\leq p(t, x, y) \\ &\leq c_3 t^{-d_H/\beta} \exp\left(-c_4 \left(\frac{d_g(x, y)^\beta}{t}\right)^{\frac{1}{\beta-1}}\right), \end{aligned}$$

where d_H is the Hausdorff dimension of \mathcal{G} .

Since $d_g \asymp d$ by lemma 2.3, it suffices to consider the Euclidean metric d in the following.

We will closely follow Barlow’s book [3] and Hambly and Kumagai’s paper [18]. We only provide some essential estimates, including a Nash inequality and an estimate of the resistance metric R .

For convenience, for $0 < s < 1$, we write $\tilde{W}_s = \{w \in \tilde{W}_* : \rho_w \leq s < \rho_{([w]_{|w|_{-1}})}\}$, and by identifying words representing the same cells, we get a quotient class \hat{W}_s .

PROPOSITION 7.2 Nash inequality. Let $d_S = \frac{2d_H}{d_H + \theta}$ with $\theta = \frac{\log r}{\log \rho}$, and $f \in \mathcal{F}$, we have

$$\|f\|_{L^2(\mathcal{G})}^{2+4/d_S} \leq c \left(\mathcal{E}(f) + \|f\|_{L^2(\mathcal{G})}^2\right) \|f\|_{L^1(\mathcal{G})}^{4/d_S},$$

for some constant $c > 0$ independent of f .

Proof. The proof is essentially the same as that for PCF self-similar sets [18]. We reproduce it here for convenience of readers. First, we claim that for any $f \in \mathcal{F}$, $\|f\|_{L^2(\mathcal{G})}^2 \leq c_1(\mathcal{E}(f) + \|f\|_{L^1(\mathcal{G})}^2)$ for some constant $c_1 > 0$ independent of f . In fact, let $\bar{f} = \int_{\mathcal{G}} f d\mu_H$ and $g = f - \bar{f}$, it suffices to check that $\|g\|_{L^2(\mathcal{G})}^2 \leq c_2 \mathcal{E}(f)$ for some $c_2 > 0$, which follows from

$$\begin{aligned} \|g\|_{L^2(\mathcal{G})}^2 &= \frac{1}{2} \int_{\mathcal{G}} \int_{\mathcal{G}} (g(x) - g(y))^2 d\mu_H(x) d\mu_H(y) \\ &= \frac{1}{2} \int_{\mathcal{G}} \int_{\mathcal{G}} (f(x) - f(y))^2 d\mu_H(x) d\mu_H(y) \leq c_2 \mathcal{E}(f), \end{aligned}$$

where the last inequality is due to (RF4) and proposition 3.3. Next, write $f_w = f \circ F_w$ for $w \in \tilde{W}_*$ for short. Then for $0 < s < 1$,

$$\begin{aligned} \|f\|_{L^2(\mathcal{G})}^2 &\leq \sum_{w \in \tilde{W}_s} \rho_w^{d_H} \|f_w\|_{L^2(\mathcal{G})}^2 \leq c_1 \sum_{w \in \tilde{W}_s} \rho_w^{d_H} \left(\mathcal{E}(f_w) + \|f_w\|_{L^1(\mathcal{G})}^2 \right) \\ &\leq c_3 s^{d_H + \theta} \sum_{w \in \tilde{W}_s} \rho_w^{-\theta} \mathcal{E}(f_w) + c_4 s^{-d_H} \sum_{w \in \tilde{W}_s} (\rho_w^{d_H} \|f_w\|_{L^1(\mathcal{G})})^2 \\ &\leq c_5 \left(s^{d_H + \theta} \mathcal{E}(f) + s^{-d_H} \|f\|_{L^1(\mathcal{G})}^2 \right), \end{aligned}$$

for some $c_3 - c_5 > 0$, where in the last inequality, we use the observation that $\sum_{w \in \tilde{W}_s} \rho_w^{-\theta} \mathcal{E}(f_w) \leq c' \mathcal{E}(f)$ for some $c' \geq 1$. In the case that $\mathcal{E}(f) > \|f\|_{L^1(\mathcal{G})}^2$, we choose s such that $s^{2d_H + \theta} \mathcal{E}(f) = \|f\|_{L^1(\mathcal{G})}^2$, then $\|f\|_{L^2(\mathcal{G})}^2 \leq 2c_5 \mathcal{E}(f)^{\frac{d_H}{2d_H + \theta}} \|f\|_{L^1(\mathcal{G})}^{\frac{2d_H + 2\theta}{2d_H + \theta}}$, and so the desired result follows immediately. In the case that $\mathcal{E}(f) \leq \|f\|_{L^1(\mathcal{G})}^2$, we have $\|f\|_{L^2(\mathcal{G})}^2 \leq c_1 (\mathcal{E}(f) + \|f\|_{L^1(\mathcal{G})}^2) \leq 2c_1 \|f\|_{L^1(\mathcal{G})}^2$, and the result still follows. \square

The Nash inequality provides an upper-bound estimate $p(t, x, y) \leq c_1 t^{-d_S/2}$. In addition, $|p(t, x, y) - p(t, x, y')| \leq c_2 t^{-1-d_S/2} R(y, y'), \forall 0 < t \leq 1, x, y, y' \in \mathcal{G}$. See [11] for a proof.

PROPOSITION 7.3. *Let $R(\cdot, \cdot)$ be the resistance metric associated with $(\mathcal{E}, \mathcal{F})$ on \mathcal{G} . Then there exist $c_1, c_2 > 0$ such that*

$$c_1 d(p, q)^\theta \leq R(p, q) \leq c_2 d(p, q)^\theta, \quad \forall p, q \in \mathcal{G}.$$

In addition, for $p \in \mathcal{G}$ and $A \subset \mathcal{G}$, define $R(p, A) = \sup\{\mathcal{E}(f)^{-1} : f \in \mathcal{F}, f(p) = 1, f|_A = 0\}$. Then there exists $c_3, c_4 > 0$ such that

$$c_3 s^\theta \leq R(p, B_s^c(p)) \leq c_4 s^\theta,$$

where $B_s(p) = \{q \in \mathcal{G} : d(p, q) < s\}$ with $p \in \mathcal{G}$ and $0 < s < 1$, and $B_s^c(p)$ is the complement of $B_s(p)$ in \mathcal{G} .

Proof. We already have the estimate $R(p, q) \leq c_2 d(p, q)^\theta$ from lemma 6.4 and theorem 6.5. Now we show $R(p, B_s^c(p)) \geq c_3 s^\theta$ for $p \in \mathcal{G}$ and $0 < s < 1$. Define

$$\begin{aligned} U_{p,s,0} &= \bigcup_{w \in \tilde{W}_{p,s,0}} F_w \mathcal{G} \text{ with } \hat{W}_{p,s,0} = \{w \in \hat{W}_{s\rho^2} : p \in F_w \mathcal{G}\}, \\ U_{p,s,1} &= \bigcup_{w \in \tilde{W}_{p,s,1}} F_w \mathcal{G} \text{ with } \hat{W}_{p,s,1} = \{w \in \hat{W}_{s\rho^2} : F_w \mathcal{G} \cap U_{p,s,0} \neq \emptyset\}. \end{aligned}$$

Clearly, we have $U_{p,s,0} \subset U_{p,s,1} \subset B_s(p)$. Since $(\mathcal{E}, \mathcal{F})$ is regular, there exists $f_{p,s} \in \mathcal{F}$ so that $f_{p,s}|_{U_{p,s,1}^c} = 0$ and $f_{p,s}|_{U_{p,s,0}} = 1$. As \mathcal{G} satisfies the finite type property, there exists a finite class $\{(p_i, s_i)\}_{i=1}^N$ such that for any $p \in \mathcal{G}$ and $0 < s < 1$, there exists $1 \leq i \leq N$ and an affine map ψ such that $\psi : U_{p,s,l} \rightarrow U_{p_i,s_i,l}$ for $l = 0, 1$,

which maps cells corresponding to $\hat{W}_{p,s,l}$ to those corresponding to $\hat{W}_{p_i,s_i,l}$. In addition, we require that ψ maps the boundary of $U_{p,s,l}$ to the boundary of $U_{p_i,s_i,l}$, which only depend on how the outside cells of approximately same size intersect $U_{p_i,s_i,1}$. Thus, we can assume that

$$f_{p,s}(q) = \begin{cases} f_{p_i,s_i} \circ \psi(q), & \text{if } q \in U_{p,s,1}, \\ 0, & \text{if } q \in U_{p,s,1}^c. \end{cases}$$

By a similar observation as in lemma 6.3, there exists $m \in \mathbb{Z}$ such that

$$\mathcal{D}^{(n)}(f_{p_i,s_i}) \leq \rho_\psi^\theta \mathcal{D}^{(n+m)}(f_{p,s}),$$

where ρ_ψ is the similarity ratio of ψ . So we have $\mathcal{E}(f_{p,s}) = \rho_\psi^{-\theta} \mathcal{E}(f_{p_i,s_i}) \leq c_3^{-1} s^{-\theta}$ for some constant c_3 independent of p, s, i . Since $f_{p,s}|_{B_s^c(p)} = 0$ and $f_{p,s}(p) = 1$, we get the estimate $R(p, B_s^c(p)) \geq c_3 s^\theta$.

Finally, the estimates $R(p, q) \geq c_1 d(p, q)^\theta$ follows from the fact that $R(p, q) \geq R(p, B_{d(p,q)}^c(p)) \geq c_3 d(p, q)^\theta$, and $R(p, B_s^c(p)) \leq c_4 s^\theta$ follows from the fact that $R(p, B_s^c(p)) \leq R(p, q) \leq c_2 s^\theta$ for some $q \in \mathcal{G}$ satisfying $d(p, q) = s$. □

By the resistance metric estimate in proposition 7.3, the Ahlfors regularity of the measure μ_H (lemma 2.3) and the resulted estimates from the Nash inequality, there exist a lower-bound estimate $p(t, x, y) \geq c_3 t^{-d_S/2}$ and an estimate of the hitting time $c_4 s^{\theta+d_H} \leq \mathbb{E}^x \tau(x, s) \leq c_5 s^{\theta+d_H}$, where $\tau(x, s) = \inf\{t \geq 0 : X_t \notin B_s(x)\}$. See [3, § 8] for details. Finally, by [3, theorem 3.1.1] of Barlow or by following [18], we can finally find that our diffusion is a fractional diffusion.

THEOREM 7.4. *The Hunt process $X = (\mathbb{P}^x, x \in \mathcal{G}, X_t, t \geq 0)$ associated with the form $(\mathcal{E}, \mathcal{F})$ on $L^2(\mathcal{G}, d_H)$ is a fractional diffusion, with $\beta = \theta + d_H$, in the sense of definition 7.1.*

Acknowledgements

The research of Qiu was supported by the National Natural Science Foundation of China, grant 12071213, and the Natural Science Foundation of Jiangsu Province in China, grant BK20211142.

Conflicts of interest

The Authors declare that there is no conflict of interest.

Data availability statement

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- 1 T. Aougab, C. S. Dong and R. S. Strichartz. Laplacians on a family of quadratic Julia sets II. *Commun. Pure Appl. Anal.* **12** (2013), 1–58.
- 2 C. Bandt and H. Rao. Topology and separation of self-similar fractals in the plane. *Nonlinearity* **20** (2007), 1463–1474.

- 3 M. T. Barlow. *Diffusions on fractals*. Lectures on Probability Theory and Statistics, pp. 1–121 (Saint-Flour, 1995), Lecture Notes in Math., vol. 1690 (Berlin: Springer, 1998).
- 4 M. T. Barlow and R. F. Bass. The construction of Brownian motion on the Sierpinski carpet. *Ann. Inst. Henri Poincaré* **25** (1989), 225–257.
- 5 M. T. Barlow and R. F. Bass. Transition densities for Brownian motion on the Sierpinski carpet. *Probab. Theory Relat. Fields* **91** (1992), 307–330.
- 6 M. T. Barlow and R. F. Bass. Brownian motion and harmonic analysis on Sierpinski carpets. *Can. J. Math.* **51** (1999), 673–744.
- 7 M. T. Barlow, R. F. Bass, T. Kumagai and A. Teplyaev. Uniqueness of Brownian motion on Sierpinski carpets. *J. Eur. Math. Soc.* **12** (2010), 655–701.
- 8 M. T. Barlow and E. A. Perkins. Brownian motion on the Sierpinski gasket. *Probab. Theory Relat. Fields* **79** (1988), 543–623.
- 9 S. Cao and H. Qiu. Resistance forms on self-similar sets with finite ramification of finite type. *Potential Anal.* **54** (2021), 581–606.
- 10 S. Cao, M. Hassler, H. Qiu, E. Sandine and R. S. Strichartz. Existence and uniqueness of diffusions on the Julia sets of Misiurewicz-Sierpinski maps. *Adv. Math.* **389** (2021), 107922.
- 11 E. A. Carlen, S. Kusuoka and D. W. Stroock. Upper bounds for symmetric Markov transition functions. *Ann. Inst. H. Poincaré Probab. Stat.* **23** (1987), 245–287.
- 12 J. M. Fraser. *Assouad dimension and fractal geometry*. Cambridge Tracts in Mathematics, vol. 222 (Cambridge: Cambridge University Press, 2021).
- 13 P. J. Fitzsimmons, B. M. Hambly and T. Kumagai. Transition density estimates for Brownian motion on affine nested fractals. *Commun. Math. Phys.* **165** (1994), 595–620.
- 14 T. C. Flock and R. C. Strichartz. Laplacians on a family of quadratic Julia sets I. *Trans. Am. Math. Soc.* **364** (2012), 3915–3965.
- 15 M. Fukushima, Y. Oshima and M. Takeda. *Dirichlet forms and symmetric Markov processes*. Second revised and extended edn. De Gruyter Studies in Mathematics, vol. 19 (Berlin: Walter de Gruyter & Co., 2011).
- 16 S. Goldstein. *Random walks and diffusions on fractals*. Percolation theory and ergodic theory of infinite particle systems (Minneapolis, Minn., 1984–1985), IMA Vol. Math. Appl., vol. 8, pp. 121–129 (New York: Springer, 1987).
- 17 A. Grigor'yan, J. Hu and K.-S. Lau. Heat kernels on metric measure spaces and an application to semilinear elliptic equations. *Trans. Am. Math. Soc.* **355** (2003), 2065–2095.
- 18 B. M. Hambly and T. Kumagai. Transition density estimates for diffusion processes on post critically finite self-similar fractals. *Proc. London Math. Soc.* **78** (1999), 431–458.
- 19 B. M. Hambly and S. O. G. Nyberg. Finitely ramified graph-directed fractals, spectral asymptotics and the multidimensional renewal theorem. *Proc. Edinb. Math. Soc.* **46** (2003), 1–34.
- 20 J. Kigami. A harmonic calculus on the Sierpinski spaces. *Jpn. J. Appl. Math.* **6** (1989), 259–290.
- 21 J. Kigami. A harmonic calculus on PCF self-similar sets. *Trans. Am. Math. Soc.* **335** (1993), 721–755.
- 22 J. Kigami. *Analysis on fractals*. Cambridge Tracts in Mathematics, vol. 143 (Cambridge: Cambridge University Press, 2001).
- 23 T. Kumagai. Estimates of transition densities for Brownian motion on nested fractals. *Probab. Theory Relat. Fields* **96** (1993), 205–224.
- 24 S. Kusuoka. A diffusion process on a fractal. In *Probabilistic methods in mathematical physics, pro. taniguchi intern. symp. (Katata/Kyoto, 1985)* (eds. K. Ito and N. Ikeda), pp. 251–274 (Boston: Academic Press, 1987).
- 25 S. Kusuoka and X. Y. Zhou. Dirichlet forms on fractals: Poincaré constant and resistance. *Probab. Theory Relat. Fields* **93** (1992), 169–196.
- 26 K.-S. Lau and S.-M. Ngai. A generalized finite type condition for iterated function systems. *Adv. Math.* **208** (2007), 647–671.
- 27 T. Lindstrøm. Brownian motion on nested fractals. *Mem. Am. Math. Soc.* **83** (1990), iv+128.
- 28 D. Mauldin and S. Williams. Hausdorff dimension in graph directed constructions. *Trans. Am. Math. Soc.* **309** (1998), 811–829.

- 29 V. Metz. Hilbert's projective metric on cones of Dirichlet forms. *J. Funct. Anal.* **127** (1995), 438–455.
- 30 M. Moran. Hausdorff measure of infinitely generated self-similar sets. *Monatsh. Math.* **122** (1996), 387–399.
- 31 S.-M. Ngai and Y. Wang. Hausdorff dimension of self-similar sets with overlaps. *J. London Math. Soc.* **63** (2001), 655–672.
- 32 H. Rao and Z.-Y. Wen. A class of self-similar fractals with overlap structure. *Adv. Appl. Math.* **20** (1998), 50–72.
- 33 L. G. Rogers and A. Teplyaev. Laplacians on the Basilica Julia sets. *Commun. Pure Appl. Anal.* **9** (2010), 211–231.
- 34 C. Sabot. Existence and uniqueness of diffusions on finitely ramified self-similar fractals (English, French summary). *Ann. Sci. École Norm. Sup.* **30** (1997), 605–673.
- 35 A. Teplyaev. Harmonic coordinates on fractals with finitely ramified cell structure. *Can. J. Math.* **60** (2008), 457–480.