

NOTES FOR AUTHORS

The *Bulletin of Entomological Research* publishes original research papers concerning insects, mites, ticks or other arthropods of economic importance in agriculture, forestry, stored products, biological control, medicine, animal health and natural resource management. The geographical scope of the *Bulletin* is worldwide but with emphasis on the tropics. Taxonomic papers are accepted if relevant. Short review papers, although normally by invitation, will also be considered for publication.

Page Format. The *Bulletin* is printed in a two-column format (column width of 80 mm) with a text area of 170×225 mm.

Text. Papers should be typed, on one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side and with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used. Guidelines for taxonomic papers are available.

When the paper has been accepted word-processed text stored on floppy disk is encouraged, providing the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with fewer type-setting errors.

Abstract. Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words. A short title should be provided as a running head.

Tables. Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or landscape (if absolutely necessary) format. Table captions should be typed on a separate sheet.

Illustrations. Copies only of artwork should be submitted. The original illustrations should accompany the paper after acceptance and revision. Text figures, line drawings, computer-generated figures and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Half-tone photographs are acceptable where they are a real contribution to the text. Figure and captions should be typed on a separate sheet in

the following format:

Figs 23–26. Figs 23–24, *Urophora* eggs: 23, *U. hispanica*; 24, *U. stigma*. Figs 25–26, spermathecae: 25, *U. maura*; 26, *U. stigma*; scale lines = 0.05 mm.

Voucher specimens. The deposition of voucher specimens should be considered where appropriate.

References. References must be based on the name and year system, give full journal titles and conform to the following styles:

Powell, W. (1986) Enhancing parasitoid activity in crops. pp. 319–340 in Waage, J. & Greathead, D. (Eds) *Insect parasitoids*. London, Academic Press (Symposium, Royal Entomological Society of London No. 13).

Southwood, T.R.E. (1978) *Ecological methods with particular reference to the study of insect populations*. 2nd edn. 524 pp. London, Chapman & Hall.

Zhou, X., Carter, N. & Mumford, J. (1989) A simulation model describing the population dynamics and damage potential of the rose grain aphid, *Metopolophium dirhodum* (Walker) (Hemiptera: Aphididae), in the UK. *Bulletin of Entomological Research* 79, 373–380.

Citation of authors in the text should appear in the form: Polaszek (1990) or (Polaszek, 1990). More than one author should be cited in chronological order as: (Holloway *et al.*, 1987; Walker & Huddleston, 1988).

Offprints. 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

Manuscripts. Three copies of the manuscript and artwork should be submitted to:

The Editors
Bulletin of Entomological Research
International Institute of Entomology
56, Queen's Gate
London
SW7 5JR, UK.

Bulletin of Entomological Research

Guest Editorial: Matteson, P. C. 'Farmer First' for establishing IPM	293
Guest Editorial Response: Ferrar, P. Classical biological control and social equity. A reply to Altieri	297
Blackwell, A., Mordue (Luntz), A. J., Young, M. R. & Mordue, W. Bivoltinism, survival rates and reproductive characteristics of the Scottish biting midge, <i>Culicoides impunctatus</i> (Diptera: Ceratopogonidae) in Scotland	299
Cross, A. E. & Morre, D. Developmental studies on <i>Anagryrus mangicola</i> (Hymenoptera: Encyrtidae), a parasitoid of the mealybug <i>Rastrococcus invadens</i> (Homoptera: Pseudococcidae)	307
Gagné, R. J. & Orphanides, G. M. The pupa and larva of <i>Asphondylia gennadii</i> (Diptera: Cecidomyiidae) and taxonomic implications	313
Grout, T. G. & Richards, R. L. The dietary effect of windbreak pollens on longevity and fecundity of a predacious mite <i>Euseius addoensis addoensis</i> (Acari: Phytoseiidae) found in citrus orchards in South Africa	317
Hargrove, J. W. & Brady, J. Activity rhythms of tsetse flies (<i>Glossina</i> spp.) (Diptera: Glossinidae) at low and high temperatures in nature	321
Hattingh, V. & Samways, M. J. Prey choice and substitution in <i>Chilocorus</i> spp. (Coleoptera: Coccinellidae)	327
Hill, J. K. & Gatehouse, A. G. Effects and temperature and photoperiod on development and pre-reproductive period of the silver Y moth <i>Autographa gamma</i> (Lepidoptera: Noctuidae)	335
Hoffmann, J. H. & Moran, V. C. Oviposition patterns and the supplementary role of a seed-feeding weevil, <i>Rhysomatus marginatus</i> (Coleoptera: Curculionidae), in the biological control of a perennial leguminous weed, <i>Sesbania punicea</i>	343
Kubasu, S. S. The ability of <i>Rhipicephalus appendiculatus</i> (Acarina: Ixodidae) stocks in Kenya to become infected with <i>Theileria parva</i>	349
Kumar, H. Oviposition, larval arrest and establishment of <i>Chilo partellus</i> (Lepidoptera: Pyralidae) on maize genotypes during anthesis	355
Mihok, S., Otieno, L. H. & Tarimo, C. S. Trypanosome infection rates in tsetse flies (Diptera: Glossinidae) and cattle during tsetse control operations in the Kagera River region of Rwanda	361
Nakache, J., Dunkelblum, E., Kehat, M., Anshelevich, L. & Harel, M. Mating disruption of the spiny bollworm, <i>Earias insulana</i> (Lepidoptera: Noctuidae), with Shin Etsu twist-tie ropes in Israel	369
Polaszek, A., Evans, G. A. & Bennett, F. D. <i>Encarsia</i> parasitoids of <i>Bemisia tabaci</i> (Hymenoptera: Aphelinidae, Homoptera: Aleyrodidae): a preliminary guide to identification	375
Ross, K. T. A. & Anderson, M. Larval responses of three vegetable root fly pests of the genus <i>Delia</i> (Diptera: Anthomyiidae) to plant volatiles	393
Scott, J. K. Biology and climatic requirements of <i>Perapion antiquum</i> (Coleoptera: Apionidae) in southern Africa: implications for the biological control of <i>Emex</i> spp. in Australia	399
Shufran, K. A., Margolies, D. C. & Black, W. C. IV. Variation between biotype E clones of <i>Schizaphis graminum</i> (Homoptera: Aphididae)	407
Torr, S. J. & Hall, M. J. R. Odour-baited targets to control New World screwworm, <i>Cochliomyia hominivorax</i> (Diptera: Calliphoridae): a preliminary study	417
Way, M. J., Cammell, M. E. & Paiva, M. R. Studies on egg predation by ants (Hymenoptera: Formicidae) especially on the eucalyptus borer <i>Phoracantha semipunctata</i> (Coleoptera: Cerambycidae) in Portugal	425
Book Reviews	433